



CMES 2024 Symposium is supported by TUBITAK



BOOK OF ABSTRACTS

ISBN-77733



**8th International Conference on
Computational Mathematics
and Engineering Sciences**

17 – 19 May 2024,
Şanlıurfa – Türkiye

Publish date: 16.05.2024

THE EIGHTH INTERNATIONAL CONFERENCE ON COMPUTATIONAL MATHEMATICS AND ENGINEERING SCIENCES (CMES- 2024), ŞANLIURFA/TÜRKİYE, MAY 17-19, 2024

The Eighth International Conference on Computational Mathematics and Engineering Sciences (CMES-2024) will be held in Harran University from 17- to 19 May 2024 in Şanlıurfa, Türkiye. It provides an ideal academic platform for researchers and professionals to discuss recent developments in both theoretical, applied mathematics and engineering sciences. This event also aims to initiate interactions among researchers in the field of computational mathematics and their applications in science and engineering, to present recent developments in these areas, and to share the computational experiences of our invited speakers and participants.

The Organizing Committee

©All Rights Reserved. This conference is organized by a cooperation of several international organizations including Fırat University, Fırat International University, Moulay Ismail University, Private University of Fes, Harran University, Van Yüzüncü Yıl University, Manas University, Ordu University, Erzurum Teknik University and İnönü University. No part of this book can be reproduced or utilized in any forms or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval systems, without permission from the authors.

© The informations provided in the papers published in this book are under the responsibility of their author(s).

Committee Chairs

Hasan Bulut, Fırat University, Türkiye
Zakia Hammouch, Ecole Normale Supérieure de Meknes, Moulay Ismail University, Morocco
Ercan Çelik, Kyrgyz-Turkish Manas University, Kyrgyzstan
Hacı Mehmet Baskonus, Harran University, Türkiye

MESSAGE FROM THE GENERAL CHAIRS



Dear Conference Attendees,

We are honored to welcome you to the **Eighth International Conference on Computational Mathematics and Engineering Sciences (CMES-2024)** at Harran University from 17 to 19 May 2024 in Şanlıurfa City, Türkiye.

CMES, founded in 2016 at Faculty of Science and Techniques Errachidia Moulay Ismail University Morocco is an annual international conference, which was very successful in the past years by providing opportunities to the participants in sharing their knowledge and informations and promoting excellent networking among different international universities. This year, the conference includes 200 extended abstracts, several submissions were received in response to the call for papers, selected by the Program Committee. The program features keynote talks by distinguished speakers such as:

Dumitru Baleanu from Institute of Space Sciences, Magurele-Bucharest, Romania; **Yusif Gasimov** from Azerbaijan University, Azerbaijan; **Naim L. Braha** from University of Prishtina, Kosovo; **Ekrem Savas** from Usak University, Türkiye; **Mehmet Emir Köksal** from Ondokuz Mayıs University, Türkiye; **Amdulla O. Mekhrabov** from Azerbaijan Technical University, Azerbaijan. The conference also comprises contributed sessions, posters sessions and various research highlights.

We would like to thank the Program Committee members and external reviewers for volunteering their time to review and discuss submitted abstracts. We would like to extend special thanks to the Honorary, Scientific and Organizing Committees for their efforts in making CMES-2024 a successful event. We would like to thank all the authors for presenting their research studies during our conference. We hope that you will find CMES-2024 interesting and intellectually stimulating, and that you will enjoy meeting and interacting with researchers around the world.

Hasan Bulut,

Firat University, Elazig, Türkiye.

Zakia Hammouch,

ENS Meknes, Moulay Ismail University Morocco

TOPICS

Control Theory,
Game Theory,
Applied Mathematics,
Financial Mathematics,
Artificial Intelligence,
Education Sciences,
Engineering Sciences,
Computer Science,
Information Technology,
Geometry and Its Applications,
Analysis and Its Applications,
Statistics and Its Applications,
Algebra and Its Applications,
Topology and Its Application,
Chaos and Dynamical Systems,
Cryptography and its Applications,
Fractional Calculus and Applications,
Economics and Econometric Studies,

Electrical and Electronic Engineering,
Defense industry and applications,
Mathematical Biology,
Computational Epidemiology,
Mathematical Chemistry,
Mathematics Education and Its Applications,
Numerical Methods and Scientific
Programming,
Linear and Nonlinear programming and
Dynamics,
Modeling of Bio-systems for Optimization
and Control,
Ordinary, Partial, Stochastic and Delay
Differential Equations,
Computational Fluids mechanics. Heat and
Mass Transfers,
Earth Sciences,
Applied Sciences

COMMITTEE CHAIRS

Prof.Dr.Hasan Bulut, Firat University, Elazığ, Türkiye

Prof.Dr.Zakia Hammouch, ENS Meknes, Moulay Ismail University, Morocco

Prof. Dr. Ercan Çelik, Kyrgyz-Turkish Manas University, Kyrgyzstan

Prof.Dr.Haci Mehmet Baskonus, Harran University, Türkiye

COMMITTEE CO-CHAIRS

Prof.Dr. Carlo Cattani, Tuscia University, Viterbo, Italy

Prof.Dr. Mohammed Ouazzani Jamil, University Privée de Fes, Morocco

Prof. Dr. Hassan Qjidaa, Sidi Mohammed Ben Abdellah University, Fes Morocco

Prof. Dr. El Mehdi El Khattabi, Moulay Ismail University Morocco

Assoc. Prof. Dr. Tolga Aktürk, Ordu University, Türkiye

HONORARY COMMITTEE

- Prof. Dr. Mehmet Tahir Güllüoğlu, Rector of Harran University, Sanliurfa, Türkiye
Prof. Dr. Fahrettin Gökteş, Rector of Firat University, Elazığ, Türkiye
Prof. Dr. Alpaslan Ceylan, Rector of Kyrgyz-Turkish Manas University, Kyrgyzstan
Prof. Dr. Vilayet M. Veliyev, Rektör of Azerbaijan Technical University, Baku, Azerbaijan
Prof. Dr. Ahmed Mouchtachi, Rector of Moulay Ismail University, Meknes Morocco
Prof. Dr. Mohammed Aziz Lahlou, President of Universite Privee of Fes, Morocco
Prof. Dr. Abdelmjid Abourriche, Vice Rector of Moulay Ismail University, Meknes Morocco
Prof. Dr. Ouazzani Jamil Mohamed, Vice President of Universite Privee of fez, Morocco
Dr. Saadat Aliyeva, Rector of Azerbaijan University, Baku, Azerbaijan
Prof. Dr. Hamdullah Şevli, Rector of Van Yüzüncü Yıl University, Türkiye
Prof. Dr. Hüseyin Yaratın, Rector of Final International University, Gime, Cyprus
Prof. Dr. Ekrem Savaş, Rector of Uşak University, Türkiye
Prof. Dr. Bülent Çakmak, Rector of Erzurum Technical University, Erzurum, Türkiye
Prof. Dr. Orhan Baş, Rector of Ordu University, Ordu, Turkey
Prof. Dr. Subhan N. Namazov, Vice-rector .Azerbaijan Technical University, Baku, Azerbaijan
Prof. Dr. Etibar Penahh, Baku State University, Baku, Azerbaijan
Prof. Dr. Necdet Bildik, Retired Faculty Member
Prof. Dr. Mahmut Ergüt, Namık Kemal University, Tekirdağ, Türkiye
Prof. Dr. Yusif Gasimov, Azerbaijan University, Baku, Azerbaijan
İbrahim Taşel, Final International University, Gime, Cyprus
Şevket Ertem, Final International University, Gime, Cyprus
Veysel Demirci, (Chairman of the board of Ziver Holding)
İzzettin Toraman, (Chairman of the board of Toraman Tekstil)
Selahattin Yıldız, (Chairman of the board of Hasyil Energy)

ORGANIZING COMMITTEE

Prof. Dr. Hasan Bulut, (Chair), Fırat University, Türkiye
Prof. Dr. Zakia Hammouch, (Chair), China Medical University, Taiwan
Prof. Dr. Ercan Çelik, (Chair), Kyrgyz-Turkish Manas University, Kyrgyzstan
Prof. Dr. Hacı Mehmet Baskonus, (Chair), Harran University, Türkiye
Prof. Dr. Ouazzani Jamil Mohammed, (Co-Chair), Universite Privee de Fes Morocco
Prof. Dr. Hassan Qjidaa, (Co-Chair), Sidi Mohammed Ben Abdellah University, Fes Morocco
Prof. Dr. Carlo Cattani, (Co-Chair), Tuscia University, ItalyAssoc.
Prof. Dr. El Mehdi El Khattabi, (Co-Chair), Moulay Ismail University Morocco
Assoc. Prof. Dr. Tolga Aktürk, (Co-Chair), Ordu University, Türkiye
Prof. Dr. Amdulla O. Mekhrabov, Director, Azerbaijan Technical University, Azerbaijan
Prof. Dr. Mikail Et, Fırat University, Türkiye
Prof. Dr. Mahmut Işık, Harran University, Türkiye
Prof. Dr. Vedat Asil, Fırat University, Türkiye
Prof. Dr. Fevzi Erdoğan, Yüzüncüyıl University, Türkiye
Prof. Dr. Alaattin Esen, İnönü University, Türkiye
Prof. Dr. Sevilay Kırıcı Serenbay, Harran University, Türkiye
Assoc. Prof. Dr. Mahmut Modanlı, Harran University, Türkiye
Assoc. Prof. Dr. Hacer Şengül Kandemir, Harran University, Türkiye
Assis. Prof. Dr. Mustafa Çağrı GÜRBÜZ, Harran University, Türkiye
Assis. Prof. Dr. Işıl BOZKURT, Harran University, Türkiye
Assoc. Prof. Dr. Fatih Özbağ, Harran University, Türkiye
Assis. Prof. Dr. N. Feyza Yalçın, Harran University, Türkiye
Assoc. Prof. Dr. Kemal Toker, Harran University, Türkiye
Res. Assist. Ömer Faruk BOYUN, Harran University, Türkiye
Assoc. Prof. Dr. Gülnur Yel, Final International University, Girne, Cyprus
Assoc. Prof. Dr. Gülay Oğuz, Harran University, Türkiye
Dr. Faculty Mem. Muhammed Yiğider, Erzurum Technic University, Türkiye

LANGUAGE EDITORIAL COMMITTEE

Suheyra Demirkol Orak (Fırat University, Elazığ, Türkiye)

International Communications Consultant

Rengin AK, Kırklareli University, Kırklareli, Türkiye
ikmet Kemaloğlu, Fırat University, Elazığ, Türkiye
İlkay Açıkgoz Erkaya, Ahi evran University, Türkiye
M. Evren Aydın, Fırat University, Elazığ, Türkiye

PROCEEDINGS

Full version of submitted papers will be published in Special Volumes of reputed journals. Procedure, Guidelines and Checklist for the preparation and submission of papers to the Proceedings of CMES-2024 can be found in the journals websites. The journals in which selected and peer-reviewed full papers of CMES-2024 will be published are as follows:

1. BOOK OF ABSTRACTS [Free of charge]

If Authors submit ABSTRACT TEXTS, then, after getting referees evaluations for these abstracts, they will be published in ABSTRACT PROCEEDING BOOK of CMES-2024. For FULL TEXT PAPERS, Authors have to submit their FULL TEXT PAPERS online via submission system of CMES-2024. These FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2024 after getting at least two positive reports.

2. CONFERENCE PROCEEDINGS [Free of charge]

At the beginning, if Authors submit FULL TEXT PAPERS, then, after getting at least two positive referee reports, FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2024 with ISBN:77733 number. Therefore, Abstracts of these FULL TEXT PAPERS will **NOT** be published in ABSTRACT PROCEEDING BOOK of CMES-2024.

3. FRACTAL AND FRACTIONAL JOURNAL [SCI-E], Selected papers from CMES-2024 will be published in a special issue dedicated to the Conference entitled "Feature Papers for Mathematical Physics Section".

https://www.mdpi.com/journal/fractalfract/special_issues/1TAP5BBZ45

This journal is indexed by SCI-E.

4. PROCEEDINGS OF THE INSTITUTE OF MATHEMATICS AND MECHANICS [E-SCI]

Selected papers from CMES-2024 will be published by <https://proc.imm.az/special/>

This journal is indexed by E-SCI.

5. TURKISH JOURNAL OF SCIENCE, [FREE]

Participants of CMES 2024 can submit their good quality papers to Turkish Journal of Science. After the peer review process, the papers will be published at TJOS. The authors must write "CMES 2024" as comments to the editor.

(Editor in Chief: Dr. Ahmet Ocak AKDEMİR) For online submission: <https://dergipark.org.tr/pub/tjos>

6. TURKISH JOURNAL OF INEQUALITIES, [FREE]

"Participants of CMES 2024 can submit their good quality papers to Turkish Journal of Inequalities. Selected papers will be published at TJI after the peer review process. The participants can send their papers to erhanet@tjinequality.com. The authors must write "CMES 2024" as the subject.

(Editor in Chief: Prof. Dr. Erhan SET)

<http://tjinequality.com/>

7. MATHEMATICS IN NATURAL SCIENCE (MNS)

Authors can submit their full text paper directly to the journal by using the following link

<https://www.isr-publications.com/mns>

8. MATHEMATICS IN ENGINEERING, SCIENCE AND AEROSPACE (MESA), [FREE, SCOPUS]

"Selected papers will be published after peer review in the Journal of Mathematics in Engineering, Science and Aerospace (MESA)"

(Editor in Chief: Prof. Seenith Sivasundaram)

<http://nonlinearstudies.com/index.php/mesa>

9. APPLIED MATHEMATICS AND NONLINEAR SCIENCES, [SCOPUS]

Participants of CMES 2024 can submit their high quality full text papers to Applied Mathematics and Nonlinear Sciences by selecting CMES-2024 under the Select Article Type Menu.

<https://www.editorialmanager.com/amns/default.aspx>

10. MATHEMATICAL MODELLING AND NUMERICAL SIMULATION WITH APPLICATIONS (MMNSA), [TR DİZİN]

The Special Issue on "Advanced Methods of Modelling and Numerical Computation in Science and Engineering". Authors can submit their full text paper directly to the journal by using the information provided in the following link

https://mmnsa.org/index.php/mmnsa/special_issues/SI-CMES2023

11. SYMMETRY [SCI-E] ; SPECIAL ISSUE "ADVANCES IN MATRIX TRANSFORMATIONS, OPERATORS AND SYMMETRY"

Authors can submit their full text paper directly to the journal by using the following link

https://www.mdpi.com/journal/symmetry/special_issues/Advances_Matrix_Transformations_Operators_Symmetry

12. YUZUNCU YIL UNIVERSITY JOURNAL OF THE INSTITUTE OF NATURAL AND APPLIED SCIENCES (TR-Dizin)

Authors can submit their full text paper directly to the journal by using the following link

<https://dergipark.org.tr/tr/pub/yyufbed>

13. PEDAGOGICAL PERSPECTIVE (PEDPER)

Pedagogical Perspective (PedPer) is an international, double blind reviewing, non-profit, professional scientific journal. PedPer is a journal that accepts manuscripts related to pedagogy and education. <http://pedagogicalperspective.com/>

PLENARY & INVITED TALKS



Generalised fractional operators with some applications

Dumitru Baleanu

Lebanese American University, Department of Computer Science and Mathematics, Beirut,
Lebanon

Institute of Space Sciences, Magurele-Bucharest, Romania

dumitru.baleanu@lau.edu.lb

Abstract: We know that fractional calculus deals with the study of so-called fractional order integral and derivative operators over real or complex domains, and their applications. However, a clear definition of a generalized fractional operator is needed. In this talk I will concentrate on solving this important issue and provide some real-world applications.

Keywords: fractional calculus, generalised fractional operators

References

- [1] Al-Refai, M, Baleanu, D (2022), On an extension of the operator with Mittag-Leffler kernel, *Fractals*, 30(5): 2240129.
- [2] Anwar A, Baleanu D (2023), On two backward problems with Dzherbashian-Nersesian operator, 8(1): 887-904, *AIMS Mathematics*.



ON SOME INVERSE PROBLEMS IN UNTRADITIONAL FORMULATION

Yusif Gasimov

Azerbaijan University, Baku, Azerbaijan

yusif.gasimov@au.edu.az

The talk is devoted to the solution of some type of inverse problems. Usually, when solving inverse problems one has to recover the equation or boundary conditions describing the process using given additional conditions. As such conditions usually some signals received from the object may be taken. These signals in mathematical formulation are called spectral data that must satisfy some conditions. The searched objects are some functions, coefficients in the equations or in the boundary conditions.

The problems considered in the talk are different from the traditional ones. Here we consider the inverse problems for some operators and the searched object are not functions as usual but are domains. We try to identify the domain where the process is going on. To solve such problems one meets some serious mathematical problems. The first problem is the choice of additional conditions – spectral data that satisfies all necessary conditions and allows to find the domain. The second problem is to construct a constructive mathematical apparatus that allows to work with functionals of the domains. To do this the space of the domains should be developed with all necessary mechanisms.

In the work the space of the convex bounded domains is constructed and a scalar product is defined there. Then the definition of the s-functions expressed by the spectral data of the Schrodinger operator is given. A scheme is proposed to solve the following inverse spectral problem with respect to domain: Define a domain on the boundary of which the s-functions of the Schrodinger operator are equal to the given functions.

REFERENCES

1. Niftiyev A.A. Gasimov Y.S. (2004). Control by Boundaries and Eigenvalue Problems with Variable Domain. Baku University Publishing, 185p.
2. Gasimov, Y.S. (2008). Some shape optimization problems for eigenvalues. *Journal of Physics A: Mathematical and Theoretical*, 41(5), 055202.
3. Gasimov, Y.S. (2013). On a shape design problem for one spectral functional. *Journal of Inverse and Ill-Posed Problems*, 21(5), 629-637.



THE SECRET BEHIND WESTERN CIVILIZATION: ISLAMIC SCIENCE

Ekrem Savas¹

¹ Department of Mathematics, Usak University, Usak, Turkey

ekremsavas@yahoo.com,

Abstract

In this study; what is the place of the Islamic Cultural world in the history of sciences? I will try to explain this. I will also explain that Western civilization is the continuation of Islamic civilization under different geographical and economic conditions.

Keywords: Islamic culture; Western civilization

REFERENCES

1. Fuat Sezgin, İslam Bilim tarihi, Timaş yayınları, 2015.



Fractional Order Thinking and Proportional-Integral-Derivative (PID) Control

Mehmet Emir KÖKSAL

Department of Mathematics, Ondokuz Mayıs University, 55139 Atakum, Samsun, Turkey

mekoksal@omu.edu.tr

Abstract: The subject of fractional calculus has become very well-known and popular in recent decades. This is because fractional-order models simulate the properties of real systems better than whole order models. Therefore, fractional calculus is used as a powerful and important tool for defining, investigating, analyzing, solving, and understanding many different chemical, engineering, mathematical, physical, statistical, and social problems in real life. In this lecture, the basic concepts of fractional calculus and various common definitions of fractional integration and differentiation are introduced. Various applications in science and engineering are mentioned. In particular, the design of fractional-order proportional integral derivative controllers is emphasized. Mathematical formulations of five design specifications corresponding to the 3D drawing are presented with program implementations. The system design specifications of phase margin, gain margin, phase flatness, low-frequency output noise suppression, and high-frequency noise suppression are considered for designing controllers using the presented 3D graphical method. Each specification is represented by some surfaces that define the boundaries of the permissible parameters of PID control coefficients. The requirements are mapped in the 3D Euclid space by 3D surfaces and/or lines so that the proportional, integral, derivative control coefficients can be optimally chosen to meet the given specifications in an optimum way and to allow trade-off or compromise.

Keywords: Fractional calculus, Fractional order modeling, PID controller, FOPID controller, 3D plots.

References:

1. M.E. Koksall, Time and frequency responses of non-integer order RLC circuits, *AIMS Mathematics*, 4 (1) 61-75, 2019
2. M.E. Koksall, Stability analysis of fractional differential equations with unknown parameters, *Nonlinear Analysis: Modeling and Control*, 24 (2) 224-240, 2019
3. M.E. Koksall, Explicit commutativity conditions for second-order linear time-varying systems with non-zero initial conditions, *Archives of Control Sciences*, 29 (3) 413-432, 2019



Design and Development of Advanced Magnetic Materials via Computational Material Science for Technological Applications

Amdulla O. Mekhrabov^{1*} and M. Vedat Akdeniz²

¹Novel Materials and Nanotechnologies Institute, Azerbaijan Technical University (AzTU), Az1073, Baku, Azerbaijan, *emdulla.mehrabov@aztu.edu.az

²Novel Alloys Design and Development Lab (NOVALAB), Department of Metallurgical and Materials Engineering (Met E), Middle East Technical University (METU), 06800-Ankara, Turkey

Abstract: The presentation will be an overview of the main research thrusts at the “Novel Alloys Design and Development Lab” (NOVALAB) of MetE-METU and at "Novel Materials and Nanotechnologies" Institute of Azerbaijan Technical University (AzTU) in the designing, development and utilizations of advanced multicomponent magnetic materials for technological applications. Fundamental principles and main aspects of *Computational Materials Science (CMS)* for *modeling and simulation based “alloy design”* which has been developed over 45 years by Prof. Mekhrabov, will be presented.

Keywords: Modeling, Simulation, Soft Magnetic Materials, Metallic Glasses, Nanostructured alloys, Glass Formation Ability, Monte Carlo, Reverse Monte Carlo, Molecular Dynamics

REFERENCES

1. Aykol M., Mekhrabov A.O. and Akdeniz M.V., Nano-scale Phase Separation in Amorphous Fe-B Alloys: Atomic and Cluster Ordering, Acta Mater., vol. 57, 171- 81, 2009.
2. Aykol M., Akdeniz M.V. and Mekhrabov A.O., Solidification behavior, glass forming ability and thermal characteristics of soft magnetic Fe-Co-B-Si-Nb-Cu bulk amorphous alloys, Intermetallics, vol. 19, 1330-1337, 2011.
3. M.V. Akdeniz and A.O. Mekhrabov, Size dependent stability and surface energy of amorphous FePt nanoalloy, J. of Alloys and Comp., vol. 788, 787-798, 2019.



Approximation by modified (p, q) -gamma-type operators

Naim Latif Braha

nbraha@yahoo.com

Department of Mathematics and Computer Sciences, University of
Prishtina, Avenue Mother Teresa,
No-5, Prishtine, 10000, Kosova and
ILIRIAS Research Institute, Janina, No-2, Ferizaj, 70000, Kosovo

Abstract

The main object of this paper is to construct a new class of modified (p, q) -Gamma-type operators. For this new class of operators, in section one, the general moments are found; in section two, the Korovkin-type theorem and some direct results are proved by considering the modulus of continuity and modulus of smoothness and their behavior in Lipschitz-type spaces. In section three, some results in the weighted spaces are given, and in the end, some shape-preserving properties are proven.

Keywords: Modified (p, q) -Gamma-type operators; Modulus of continuity; Shape-preserving approximation

References

1. Altomare, F., Campiti, M.: Korovkin-Type Approximation Theory and Its Application. Walter de Gruyter Studies in Math., vol. 17. de Gruyter & Co., Berlin (1994)
2. Atlihan, O.G., Unver, M., Duman, O.: Korovkin theorems on weighted spaces: revisited. Period. Math. Hung. 75(2), 201–209 (2017)

TYPE	AUTHORS	TITLES	PAGE	TOPICS
Oral	Sıdıka Şule Şener Kılıç, Adem Irmak, Arzu Aykut	Ritz Method for the Numerical Solution of the Heat Equation	1	Applied Mathematics
Oral	Ambreen Siyal, Kashif Ali Abro	Repel Effects of Heat Transference from Brinkman Fluid under Ferromagnet via Non-Singularized Differentials	2	Computational Fluids mechanics
Oral	Abd Essamed Guettouche, Chaabane Djama	Optimizing a linear function over the set of efficient solutions: Case of the stochastic set-covering problem.	3	Applied Mathematics
Online	Hakkı Güngör	The Novel Numerical Solutions Of Conformable Fractional Benjamin–Bona– Mahony Equation By Using The Robust Conformable Method	4	Applied Mathematics
Online	Hacer Bilgin Ellidokuzoğlu, Serkan Demiriz	q- Paranormed Difference Sequence Spaces	5	Analysis and Its Applications
Online	Hacer Bilgin Ellidokuzoğlu, Serkan Demiriz	Paranormed Narayana Sequence Spaces	6	Analysis and Its Applications
Oral	Hasan Karaçalı, Orhan Özdemir	On the Oscillation of a Second Order Differential Equation With a Superlinear Neutral Term	7	Applied Mathematics
Online	Hayatem Hamal	Estimates of Bivariate New Kantorovich Type of the Balázs–Szabados Operators Based on q-integers	8	Analysis and Its Applications
Oral	Melik Şenyuva, Özlem Kirci, Hasan Bulut	New Exact Wave Solutions of the New Hamiltonian Amplitude Equation Through $(m + 1/G')$ -Expansion Method	9	Applied Mathematics
Oral	Orhan Özdemir	Oscillation of Second Order Neutral Emden–Fowler Differential Equations	10	Applied Mathematics
Oral	Ercan Tunç	Improved oscillation criteria for third-order half-linear delay differential equations via canonical transform	11	Applied Mathematics
Online	Süleyman Cengizci	Applications of the SUPG-YZ finite element formulation: from mussel-algae interactions to Schnakenberg reaction models	12	Applied Mathematics
Online	Haci Mehmet Baskonus, Halil Şençiçek	Some New Analytical Solutions to the Nonlinear Modified Quantum Zakharov–Kuznetsov Equation	13	Applied Mathematics
Oral	Orhan Dalkılıç, Esin İlhan, Hasan Bulut	Comparative Analysis of Rankings and Preference Values for Fuzzy Decision-Making Approaches in Reducing Unnecessary Biopsies	14	Topology and Its Application
Online	Artion Kashuri, Rozana Liko	Estimations of Integral Majorization Inequality For Differentiable Convex Functions And Applications	15	Analysis and Its Applications
Oral	Gözde Karataş Baydoğan, Şahsene Altınkaya	A Survey On Different Statistical Distributions Using Python Programming	16	Computer Science
Oral	Gözde Karataş Baydoğan, Şahsene Altınkaya, Nahide Zeynep Cicekli	Exploring Machine Learning Techniques For Gender Voice Recognition Using Limited Speech Data	17	Computer Science
Online	Yaşar Çalışkan, Mikail Et	On Lacunary Statistical Boundedness of Sequences of Sets	18	Analysis and Its Applications
Online	Ayşe Eren, Mikail Et	On λ -Statistical Boundedness of Sequences of Sets	19	Analysis and Its Applications
Online	Murat Turan, Hülya Gün Bozok, Mahmut Ergüt	Inextensible Flows Of Curves With Quasiframe In Galilean Space G_3	20	Geometry and Its Applications

Online	Murat Temizkan, Hıfı Altınok	On Differences Of Bounded Variation Fuzzy Sequences	21	Analysis and Its Applications
Online	Mithat Kasap, Hıfı Altınok	On Lacunary Statistical Boundedness	22	Analysis and Its Applications
Oral	Tuğba Yavuz	On a Coefficient Problem For Functions Belongs to Certain Subclass of Univalent Functions	23	Analysis and Its Applications
Online	İlkay Koçoğlu, Hasan Bulut	New Wave Behaviors For Solutions Of The Truncated M-Fractional Variant Boussinesq System	24	Applied Mathematics
Oral	Neşe İşler Acar	A Collocation Method for Numerical Solution of Linear Integro-Differential Equations by Stancu Polynomials	25	Applied Mathematics
Oral	Metin Turgay	Approximation Properties of Kantorovich Type Sampling Series In Weighted Spaces of Functions	26	Analysis and Its Applications
Oral	Beyhan Kemaloglu, Gülnur Yel, Hasan Bulut	Analytical Solution of Hirota Equation by Rational SineGordon Method	27	Applied Mathematics
Online	Burak Çevik, Muharrem Tuncay Gençoğlu	Blockchain Applications In Medula System	28	Computer Science
Oral	Muhammed Huzeyfe Uzunyol, Berat Karaagac, Alaattin Esen	Crank-Nicolson Finite Difference Treatment of Time Fractional Klein Gordon Equation	29	Fractional Calculus and Applications
Oral	Gülay Oğuz, Abdülkadir Olcay	The Relations of Soft Topological Hyperstructures	30	Topology and Its Application
Oral	Gülay Oğuz, Ayhan Yüksel	Rough Approximation Operators on Algebraic Hyperstructures	31	Topology and Its Application
Oral	Shorish Omer Abdulla, Mahmut Modanlı	Analytic Solutions for Third-Order Fractional Partial Differential Equation Using Modified Double Laplace Transform Method	32	Fractional Calculus and Applications
Oral	Sadettin Kursun	Some New Results for Exponential-type Durrmeyer Sampling Series	33	Analysis and Its Applications
Oral	Ugur Bayrakcı, Seyma Tuluçe Demiray, Huseyin Yıldırım	New soliton solutions with generalized exponential rational function method	34	Applied Mathematics
Online	Merve Karaoğlu, Erdal Baş	Introduction To M-Sturm-Liouville Problem For Diffusion Operator	35	Fractional Calculus and Applications
Oral	M. Mustafa Beydağı, A. Fatih Özcan, İlhan İçen	Properties of Rough Subgrupoids	36	Topology and Its Application
Oral	Ulviye Demirbilek, Mehmet Şenol, Hasan Bulut	Solving Dynamic Complexity with Analytical Solution Techniques	37	Linear and Nonlinear programming and Dynamics
Oral	Semih Geçen, İlhan İçen, A. Fatih Özcan	Grupoid Atlases	38	Topology and Its Application
Oral	Arif Özkul, Tolga Aktürk, Hasan Bulut	Modified Kudryashov Method for Solving Van der Waals Gas System	39	Applied Mathematics
Oral	Sibel Özer, Yusuf Uçar, Damla Özçelik	A Study On Numerical Solution of the Regularized Long Wave Equation	40	Applied Mathematics
Online	Sezer Erdem, Serkan Demiriz	Domain Of Mersenne Matrix Operator In The Space of Convergent Sequences	41	Analysis and Its Applications
Online	Serkan Demiriz, Sezer Erdem	A Note On Almost Convergent Mersenne Sequence Space	42	Analysis and Its Applications

Oral	Bülent Oruç, Mustafa Berkay Doğan, Emir Balkan, İlkin Özsöz, Sunay Mutlu, Aybala Büşra Çalışkur	Gravity Modelling And Earthquake Analysis For East Anatolian Fault Zone And Surrounding Area	43	Earth Sciences
Online	Erdal Baş, Ali Selçuk	Generalized Fractional The Vertical Motion Of A Falling Body Problem	44	Applied Mathematics
Oral	Md. Nur Alam, Onur Alp İlhan, Md. Shahid Hasan, Uzzal Saha, F. Berna Benli	Some New Results Of The Nonlinear Conformable Model Arising In Plasma Physics	45	Applied Mathematics
Oral	Sebahattin Ertas, Hasan Bulut, Yusuf Pandir	New Exact Solutions of the (1+1) dimensional nonlinear Ostrovsky equation	46	Applied Mathematics
Online	Enise Kartal, Erdal Baş	Generalized Systems of Linear Equations With Local Derivative	47	Fractional Calculus and Applications
Oral	Gülden Mülayim	Model Order Reduction for Shigesada-KawasakiTeramoto Cross-Diffusion Systems	48	Numerical Methods and Scientific Programming
Online	Auwalu Sa'idu, Hikmet Kemaloğlu (Koyunbakan)	An Inverse Nodal Problem Of A Conformable Sturm-Liouville Problem By Retarded Constant	49	Analysis and Its Applications
Online	Emre Cıvgın, Numan Yalcın	Fundamental Algebraic And Topological Concepts In Geometric Analysis	50	Applied Mathematics
Oral	Ayşe Fidan, Erhan Pişkin	Blow up at finite time for sixth-order evolution equations with time dependent coefficient	51	Applied Mathematics
Oral	Hatice Karabenli, Yusuf Uçar, E. Nesligül Aksan, Alaattin Esen	A Comparative Study Of Finite Element Methods With Cubic And Quintic Basis Functions For The Smch Equation	52	Applied Mathematics
Online	Dilara Altan Koç, Yusuf Pandir, Hasan Bulut	A New Generalized Method For The Fractional Nonlinear Equation	53	Applied Mathematics
Oral	Natavan Allahverdiyeva, Yusif Gasimov	Some Properties of the Eigenfrequencies on the Domain of the Plate	54	Applied Mathematics
Online	Gülstan Butakin, Erhan Pişkin	Explosive Solutions for a Fourth-Order Reaction-Diffusion Equation in Variable Exponent Sobolev Spaces	55	Applied Mathematics
Oral	Nebi Yılmaz, Erhan Pişkin	Decay of Solutions for a Nonlinear Hyperbolic-type Equations with Variable Exponents	56	Applied Mathematics
Oral	Özlem Cerit, Durmuş Yarımpabuç	Forced Vibration Analysis Of Functionally Graded Rods By Pseudospectral Chebyshev Method	57	Applied Mathematics
Online	Meral Süer	Combinatorial Invariants Of Saturated Numerical Semigroups	58	Analysis and Its Applications
Oral	Oğuzhan Demirel, Durmuş Yarımpabuç	Thermal Analysis Of Functionally Graded 2d Plate	59	Applied Mathematics
Online	Nuket Aykut Hamal, Furkan Erkan	Existence And Uniqueness Results For Singular Fractional Differential Equations With P-Laplacian Operator	60	Fractional Calculus and Applications
Oral	Fatma Almaz	The Specific Energy And Specific Angular Momentum On Rotational Surfaces In Pseudo Euclidean 4-Space With Index 2	61	Geometry and Its Applications

Oral	Mahmut Ozusan , Hikmet Kemaloğlu	Expansion Theorem for Sturm–Liouville Problem including Local Derivative	62	Analysis and Its Applications
Online	Gülşah Belhan, Vedat Asil	Focal Curves According To The Alternative Frame	63	Geometry and Its Applications
Oral	Nuri Murat Yağmurlu, Selçuk Kutluay, Ali Sercan Karakaş	Cubic Hermite Collocation Method For The Equal Width Wave Equation	64	Applied Mathematics
Online	Sevilay Kırıcı Serenbay, Ecem Acar	On The Approximation By Nonlinear Operators Of Max–Product Kind	65	Analysis and Its Applications
Oral	Kübra Heredağ, Fatma Ayaz	Examination Of Mhd Effect and Fractional Derivative Model Between Porous Medium Parallel Plates In TimeDependent Flow	66	Fractional Calculus and Applications
Online	Emre Aydın, İnci Çilingir Süngü	On The Semi–Analytical And Hybrid Methods For The Drinfeld–Sokolov–Wilson System Modelling Dispersive Water Waves	67	Numerical Methods and Scientific Programming
Oral	Hasan Gündüz, Mesut Karabacak, Ercan Çelik	The Computation Of H^∞ –Norm Of Transfer Functions Of Linear Daes Via Two–Step Method	68	Control Theory
Online	Ahmed Abuhatim, Ebru Cavlak Aslan	Investigation of extended type a NLS equation using the extended direct algebraic method	69	Applied Mathematics
Oral	Muhteşem Demir, Erhan Pişkin	Growth Of Solution For Reaction Diffusion Equation With Kirchhoff Term And Multiple Nonlinearities	70	Applied Mathematics
Online	Betül Oğraş İkiz, Zühal Küçükarslan Yüzbaşı	Characterization Of Parametric Surfaces In Lie Groups Using Alternative Frame	71	Geometry and Its Applications
Oral	Gülşen Orucova Büyükoğlu, Hüseyin Haklı	Implementation Of Battle Royale Optimization Algorithm For 0–1 Knapsack Problem Using S–Shaped And V–Shaped Transfer Functions	72	Applied Mathematics
Oral	Bahadır Yüzbaşı	Housing Price Determinants: A Big Spatial Data Analysis	73	Economics and Econometric Studies
Oral	Enver Temo, Mehmet Eker, Durmuş Yarımpabuç	Pseudospectral Chebyshev Approach For Nonlinear Temperature Distributions In Functionally Graded Disks	74	Applied Mathematics
Oral	Derya Deniz, Ebru Cavlak Aslan	New Optical Soliton Solutions of the NLS Equation with Jacobi Elliptic Function Expansion Method	75	Applied Mathematics
Oral	Zühal Küçükarslan Yüzbaşı	Motion Of The Filament In Minkowski Space	76	Geometry and Its Applications
Oral	Mehmet Uçar, Aynur Yalçiner	On The Uniformly Parikh–Friendly Words	77	Analysis and Its Applications
Oral	Ömer Oruç	A Meshfree Method For Numerical Solutions Of Some Reaction–Diffusion Type Equations	78	Numerical Methods and Scientific Programming
Online	Dilara Yapişkan, Beyza Billur İskender Eroğlu	Optimal culling strategy for the fractional–order brucellosis transmission model	79	Control Theory
Online	Meltem Karaismailoğlu, Sezin Aykurt Sepet, Mahmut Ergüt	Pointwise Hemi–Slant Submersions From Cosymplectic Manifolds	80	Geometry and Its Applications
Online	Muhittin Evren Aydın, Aykut Has, Beyhan Yılmaz	Multiplicative Rectifying Curve In Multiplicative Euclidean Space	81	Geometry and Its Applications

Oral	Merve Zeynep Kaya, Ercan Çelik, Mesut Karabacak	Solution of Fractional Order Partial Differential Equations with Hosoya Neural Network	82	Applied Mathematics
Online	Muhammed Kerem Turkes, Yıldız Aydın	A New Facial Expression Recognition Methods Based On Hybrid Feature	83	Artificial Intelligence
Online	Süleyman Sarıkaya, Yavuz Altın	f-Statistical Convergence Of Double Sequences In Topological Groups	84	Topology and Its Application
Online	Hülya Gültekin Çitil, Fatma Gizem Özmen	An Investigation Of A Fuzzy Boundary Value Problem	85	Applied Mathematics
Online	Feride Tuğrul	Explaining Of Decision Making Processes With The Help Of Intuitionistic Fuzzy Sets	86	Applied Mathematics
Online	Pınar Zengin Alp	A New Paranormed Sequence Space Given By Jordan Totient Function	87	Analysis and Its Applications
Oral	Muhammed Veysi Güler, Muhammed Emre Çolak	Detecting Android Malware Using LightGBM: A Study on the TUANDROMD Dataset	88	Artificial Intelligence
Oral	Nurettin Bağırılmaz	On The Construction Of A Topology On A Rough Semigroup	89	Topology and Its Application
Oral	Ömer Miraç Kökçam, Muhammed Emre Çolak, Özal Yıldırım	Voting Classifier Based Explainable Artificial Intelligence Method For Detecting Glioma Grading Using Clinical And Mutation Features	90	Artificial Intelligence
Online	Zeynep Gülcen Kaya, Murat Şahin, Ayça Gülten	A Comparative Analysis Of Tree-Inspired Fractal Branchings Dendriform Structures, From The Bc To The L-System Based Structures	91	Fractional Calculus and Applications
Online	Mohammed Taleb, Nouredine En-Nahnahi, Nisrine Dad	Multi-category classification of inappropriate content on social media using Natural Language Processing techniques and Transformer Models	92	Artificial Intelligence
Online	Özge Akçay	Inverse Scattering Problem For Discontinuous Sturm-Liouville Operator	93	Applied Mathematics
Online	M. Ghebleh, A. Kanso, M. B. Khuzam	A Probabilistic Chaotic Image Encryption Scheme	94	Cryptography and its Applications
Online	Servet Akbaş, Bilgi Yılmaz	Exploring Divergence Measures: Concepts, Applications, and Advances across Disciplines	95	Applied Mathematics
Oral	Fatih Avşar	Fixed Points of multiplicative Zamfirescu Mapping in Multiplicative Metric Spaces	96	Analysis and Its Applications
Oral	Koray İbrahim Atabey, Muhammed Recai Türkmen, Mikail Et, Muhammed Çınar	q-Bell Statistical Convergence	97	Analysis and Its Applications
Online	Deniz Öztürk	Mathematical Analysis and Modeling of Biofouling in Urban Water Filtration Systems	98	Applied Mathematics
Oral	Tolga Aktürk	Effective Method for Analyzing Nonlinear Mathematical Model Behavior	99	Applied Mathematics
Oral	Funda Türk, Samet Erden	Ostrowski type inequalities via fractional integrals and related results	100	Analysis and Its Applications
Online	Semra Çelebi, İbrahim Türkoğlu	Personality Analysis Using Artificial Intelligence According To The Eye Descriptions In Marifetname	101	Artificial Intelligence

Oral	Funda Türk, Samet Erden, Burçin Gökkurt Özdemir	Exponential Inequalities Involving Riemann–Liouville Fractional Integral	102	Analysis and Its Applications
Online	Aslı Öner, Sertac Goktas, Büşra Barut	Conformable Sturm–Liouville Problem With Two Parameter	103	Fractional Calculus and Applications
Oral	Koray İbrahim Atabey, Murat Karakaş	q -Pell Sequence Spaces	104	Analysis and Its Applications
Online	Gülcan Atıcı Turan	On p -Statistical Convergence Defined By Modular Sequence Spaces Of Order	105	Analysis and Its Applications
Online	Imane El Mhamedi, Zakaria El Malk	Developing High-Efficiency Organic Solar Cells through Molecular Design Analysis of Novel D-A-Di-A-D Conjugated Compounds	106	Electrical and Electronic Engineering
Oral	Enes Ata, İ. Onur Kıymaz, Hacı Mehmet Başkonuş	A New Fractional Modelling Of Rc Electric Circuit	107	Applied Mathematics
Oral	Meltem Uzun	On Wave Structures Of Time Conformable Zakharov–Kuznetsov Equation	108	Applied Mathematics
Oral	Yasemin Bakır, Oya Mert, Gülay Karahanlı	On Using A New Approach To Determine The Root Of Nonlinear Equations	109	Applied Mathematics
Oral	Şeyma Firdevs Korkmaz, Hasan Bulut, Gülnur Yel	Modeling Epidemics Using Ising Model and Voronoi Tessellation: A Novel Study and Epidemiological Applications	110	Applied Mathematics
Oral	Zulqurnain Sabir, Ayse Nur Akkilic, Hasan Bulut	Designing a novel radial basis process for the nonlinear prey–predator system	111	Applied Mathematics
Online	Mevlüt Açar, Mustafa Yeneroğlu	Lie Algebra And Some Geodesic Properties	112	Geometry and Its Applications
Oral	Meltem Öğrenmiş	Curvatures Computation For Curves In Affine Space Using Fractional Calculus	113	Geometry and Its Applications
Oral	Meltem Öğrenmiş	Expanding Fractional Equiaffine Curvatures Of Plane Curves	114	Geometry and Its Applications
Online	Ali Olgun, Zekiye Rana Lüsna, Oğuz Yağcı	Several Integral Representations of the ${}_p-k$ Srivastava's triple hypergeometric functions	115	Applied Mathematics
Oral	Mehmet Aydın, Resat Yilmazer	Fractional Solutions of the General Class of Non-Fuchsian Differential Equations	116	Applied Mathematics
Oral	Kübra Elif Akbaş, Mahmut Işık	Weighted Statistical Convergence in Probability	117	Applied Mathematics
Oral	Ayşe Metin Karakaş, Sinan Çalık	The New Gompertz Distribution	118	Statistics and Its Applications
Oral	Elif Nur Yıldırım, Münevver Tuz	Analysis Of Mathematical Model Wave Solutions With The Exponential Function Method	119	Applied Mathematics
Online	Gökhan Dere, Melih Taş	Predicting Student Performance Using Statistical Learning Techniques	120	Statistics and Its Applications
Online	Keziban Taş	On The Inverse Problem For A Secondorder Differential Operator With A Matrix Potential	121	Applied Mathematics
Oral	Tuğçem Partal, Melike Kakşı	Comparison Of Deterministic And Stochastic Dynamics Of Sir Model	122	Applied Mathematics
Oral	Özlem Defterli, Ayşe Özmen	GPLM for Regression of Complex Systems	123	Applied Mathematics
Oral	Ece Atlan, Handan Öztekin	Lanar Congruent Curves According To Caputo Fractional Derivative	124	Geometry and Its Applications

Oral	Emin Beso, Senada Kalabusić, Esmir Pilav, Arzu Bilgin	Dynamics Of A Plant-Herbivore Model Subject To Allee Effects With Logistic Growth Of Plant Biomass	125	Chaos and Dynamical Systems
Online	Ömer Akgüller, Mehmet Ali Balcı	Decoding Structural Isomer: An Artificial Intelligence Approach To Cluster Detection	126	Artificial Intelligence
Online	Merve Ak, Senem Şahan Vahaplar, M. Hakkı Ersoy, Ahmet Feyzioğlu	Measuring And Assessing Organizational Data Maturity	127	Engineering Sciences
Oral	Aslı Alkan, Tolga Aktürk, Hasan Bulut	The Novel Numerical Solutions of the Cahn-Hilliard Equation via the Novel Hybrid Method	128	Applied Mathematics
Oral	Aslı Alkan, Tolga Aktürk, Hasan Bulut	The New Numerical Solutions of the Navier-Stokes Equation with the New Hybrid Method	129	Applied Mathematics
Oral	Aslı Alkan, Tolga Aktürk, Hasan Bulut	The Novel Numerical Solutions of the Rosenau-Hyman Equation via the Novel Hybrid Method	130	Applied Mathematics
Online	Derya Avcı, Sanem Sakarya	How To Determine The Optimal Strategies To Eliminate The Harmful Effects Of Technology Addiction?	131	Control Theory
Online	Derya Avcı, Aylin Yetim	An Optimal Training Policy To Reduce Criminal Behavior	132	Control Theory
Online	Nazlım Deniz Aral , Hacer Şengül Kandemir ,Mikail Et	On f-Statistical Convergence Via q- Calculus	133	Analysis and Its Applications
Oral	Yener Altun, Şakir İşleyen	A Research On The Qualitative Behavior Of Solutions Of Neutral Systems With Periodic Coefficients	134	Applied Mathematics
Online	Fatma Uzer, Resat Yilmazer	Explicit Solutions of the Schrödinger Equation Using Fractional Analysis	135	Applied Mathematics
Online	Mine Babaoglu	Dirac Systems That Contain Eigenvalue Dependent Boundary Conditions	136	Applied Mathematics
Online	Imane El mhamedi, Anass El karkri, Zakaria El malki, Mohammed bouachrine	Voting Classifier Based Explainable Artificial Intelligence Method For Detecting Glioma Grading Using Clinical And Mutation Features	137	Electrical and Electronic Engineering
Online	Asan Omuraliev, Ella Abylaeva	Numerical Solution of the Singularly Perturbed Cauchy Problem for an Ordinary Differential Equation	138	Applied Mathematics
Online	Gülcan Tokay, Emrah Yılmaz	Infectious Disease Models With Proportional Derivatives On Time Scales	139	Applied Mathematics
Oral	Nejla Gürefe	The Concretization Process of the "Pyramid" Concept;Deaf Student Example	140	Mathematics Education and Its Applications
Oral	Yusuf Gürefe	Modified Exponential Function Method for TwoDimensional Nonlinear Mathematical Model	141	Applied Mathematics
Online	E.Panakhov, F.Asadli	The Calculation of the Trace Formulas for Dirac Operator by Lax Method	142	Applied Mathematics
Online	E.Panakhov, I.Shikhaliyeva	The Calculation of the Regularization Trace of the Diffusion Equation by Lax's Method	143	Applied Mathematics
Oral	Cemil İnan	Examining The Relationship Between Integral Equations And Differential Equations	144	Applied Mathematics
Online	Asan Omuraliev, Peyil Esengul kyzy	Asymptotics Of Solutions To A First-Order Partial Differential Equation With A Power-Law Boundary Layer	145	Applied Mathematics
Online	Zehra Özdemir, Emrah Yılmaz	Armament Model And Its Analysis With Proportional Derivative On Time Scales	146	Analysis and Its Applications

Online	Hazal Yüksekaya	Blow Up At Infinite Time Of Solutions For The Viscoelastic Plate Equation With Distributed Delay And Source Terms	147	Applied Mathematics
Online	Hazal Yüksekaya	Blow Up Results At Finite Time For The Kirchhoff-Type Viscoelastic Equation With Time Delay And Source Term	148	Applied Mathematics
Online	Yılmaz Erol, Ummahan Acar	On Prime Subhypermoudles	149	Algebra and Its Applications
Online	Hülya Gültekin Çitil, Ayşe Nur Başar	An Approach To A Fuzzy Problem With Variable Coefficients	150	Applied Mathematics

Ritz Method for the Numerical Solution of the Heat Equation

Sıdıka Şule Şener Kılıç¹, Adem Irmak², Arzu Aykut³

¹Department of Mathematics, University of Atatürk, Erzurum, Turkey

²Vocational School of Social Sciences, University of Bayburt, Bayburt, Turkey

³Department of Mathematics, University of Atatürk, Erzurum, Turkey

senersule@atauni.edu.tr, airmak@bayburt.edu.tr, aaykut@atauni.edu.tr,

Abstract

In this paper, we give the numerical solution of the boundary value problem for the heat equation. The Ritz method has been used to obtain this solution. We solve the numerical examples and errors analysis of the approximation solutions.

Keywords: Ritz method, heat equation, error analysis

REFERENCES

1. Storch, J. and Strang, G., 1988. Paradox lost: natural boundary conditions in the Ritz-Galerkin method, *International Journal for Numerical Methods in Engineering*, 26 (10), 2255– 2266.
2. Axelsson, O. and Barker, V.A., 2001. *Finite Element Solution of Boundary Value Problems: Theory and Computation*. Society for Industrial & Applied Mathematics, U.S., 431 p, New York, United States.
3. Gander, M.J., and Wanner, G., 2012. From Euler, Ritz, and Galerkin to Modern Computing. *SIAM Review*, 54 (4), 627-666.
4. Delves, L.M. and Freeman, T.L., 1981. *Analysis of Global Expansion Methods*. Academic Press, 275 p, New York.
5. Larsson, S. and Thomee, V., 2003. *Partial Differential Equations with Numerical Methods*. Springer-Verlag Berlin Heidelberg, 262 p, Berlin Heidelberg.
6. Knabner, P. and Angermann, L., 2003. *Numerical Methods for Elliptic and Parabolic Partial Differential Equations*. Springer-Verlag New York, 426 p, New York.
7. Omodei, B.J. and Anderssen, R.S., 1975. Stability of the Rayleigh-Ritz Procedure for Nonlinear Two-Point Boundary Value Problems. *Numerische Mathematik*, 24 (1), 27-38.

Repel Effects of Heat Transference from Brinkman Fluid under Ferromagnet via Non-Singularized Differentials

Ambreen Siyal¹, Kashif Ali Abro^{1*}

¹Department of Basic Sciences and Related Studies, Mehran University of Engineering and Technology, Jamshoro, Pakistan
kashif.abro@faculty.muet.edu.pk,

Abstract

When Brinkman fluid exhibits a net and strong magnetization then such Brinkman fluid leads to ferro-magnetized Brinkman fluid with high permeabilities for magneto-strictive flow behavior. This manuscript aims to present the [magneto-resistive](#) analytical solutions for the governing equations of ferro-magnetized Brinkman fluid under an effective and powerful approach Atangana-Baleanu differential operator. The governing equations of ferro-magnetized Brinkman fluid have been constructed from classical to Atangana-Baleanu differential operators. The analytical solutions are emphasized in terms of magnetized domains for velocity field, temperature and concentration profiles. Integral transforms approach is invoked to tackle the fractional verses classical solutions and ferro-magnetized verses non-ferro-magnetized solutions. Our results investigate that the transfer of heat in the presence of a ferro-magnetic has generated ripples in Brinkman fluid showing several rheological similarities and rheological dis-similarities.

Keywords: Ferro-magnetized Brinkman fluid, Ripple heat transference, Analytical mathematical approach, rheology of Ferro-magnetized Brinkman fluid.

Optimizing a linear function over the set of efficient solutions: Case of the stochastic set-covering problem.

Abd Essamed Guettouche,¹ Chaabane Djamal¹

¹USTHB Algiers. Algeria

Authors Emails: aguettouche@usthb.dz, dchaabane@usthb.dz

I. ABSTRACT

The stochastic multi-objective set-covering problem (the probabilistic multi-objective set-covering problem) is very challenging to solve directly. Generating the set of all efficient solutions might be very expensive and unproductive for the decision-maker because, in order to meet their preferences, they have to choose the best-compromised solution from a large list. If their preference is expressed as a linear combination of decision variables, one has to optimize this function over the efficient set of the multi-objective set-covering problem. In this paper, we consider a stochastic environment, i.e., all the parameters are derived from a discrete probability law. Once the problem is converted into a deterministic model, we use the technique presented and developed by (Chaabane and Pirlot 2010). To the best of our knowledge, no similar study has been conducted yet.

Keywords: Stochastic multi-objective optimization; Combinatorial optimization; Non-linear optimization; Probabilistic set-covering problem; Efficient solution.

REFERENCES

1. Chaabane D, Pirlot M, A method for optimizing over the integer efficient set, J Ind Manag Optim, 2010.
2. Chaabane D, Mebrek F, Optimization of a linear function over the set of stochastic efficient solutions, Comput Manag Sci, 2014.
3. Prékopa A, 1995, Stochastic Programming, Kluwer Academic Publishers, 1995.
4. Charnes A, Cooper W W, Deterministic equivalents for optimizing and satisficing under chance constrained, Journal of Operations Research, 1963.
5. Benson H P, Existence of efficient solutions for vector maximization problems, Journal of Optimization Theory and Appl, 1978.
6. Balas E, A class of location, distribution and scheduling problems: modeling and solution methods, J.Wiley and Sons, New York, 1983.
7. Ceria S, all, A Lagrangian-based heuristic for large-scale set covering problems, Mathematical Programming-Springer, 1998.
8. Chvatal V, A greedy heuristic for the set-covering problem, Mathematics of Operations Research, 1979.

THE NOVEL NUMERICAL SOLUTIONS OF CONFORMABLE FRACTIONAL BENJAMIN–BONA–MAHONY EQUATION BY USING THE ROBUST CONFORMABLE METHOD

Hakkı Güngör¹

¹ Department of Computer Technology, Vocational School, Ufuk University, Ankara, Turkey

hakki.gungor@ufuk.edu.tr,

Abstract

The q-formable homotopy analysis transform method is employed to analyze the conformable fractional Benjamin-Bona-Mahony equation. The numerical solutions to this problem are graphed. The suggested approach has been demonstrated to be successful and consistent based on numerical simulations.

Keywords: Benjamin-Bona-Mahony equation, q-formable homotopy analysis transform method, formable transform.

REFERENCES

1. Miller, K.S.; Ross, B. An Introduction to the Fractional Calculus and Fractional Differential Equations; John Wiley and Sons: New York, NY, USA, 1993.
2. Podlubny, I. Fractional Differential Equations, Mathematics in Science and Engineering; Academic Press: New York, NY, USA, 1999.
3. Güngör, N. (2022). Solution of Convolution Type Linear Volterra Integral Equations with Formable Transform. *International Journal of Latest Technology in Engineering, Management & Applied Science*, 11(12), 1-4.
4. Saadeh, R. Z., & Ghazal, B. F. A. (2021). A new approach on transforms: Formable integral transform and its applications. *Axioms*, 10(4), 332.

q – PARANORMED DIFFERENCE SEQUENCE SPACES

Hacer Bilgin Ellidokuzoğlu¹ and Serkan Demiriz²

¹ Department of Mathematics, University of Recep Tayyip Erdoğan, Rize, Türkiye
hacer.bilgin@erdogan.edu.tr

² Department of Mathematics, University of Tokat Gaziosmanpaşa, Tokat, Türkiye
serkandemiriz@gmail.com

Abstract

The purpose of this work is to construct the extended versions of Maddox's original paranormed sequence spaces, denoted by the notations $c_0(\nabla_q^2, p)$ and $c(\nabla_q^2, p)$. The spaces $c_0(p)$ and $c(p)$ are linearly isomorphic to these spaces. The Schauder basis for these spaces must then be constructed. Then the topological properties of the $c_0(\nabla_q^2, p)$ and $c(\nabla_q^2, p)$ are studied. Finally, some matrix classes are described.

Keywords: q -calculus; q -second difference matrix; paranormed sequence spaces; matrix transformations

REFERENCES

1. A. Alotaibi, T. Yaying, and S. A. Mohiuddine. Sequence spaces and spectrum of q -difference operator of second order. *Symmetry*, 14(6):1155, 2022.
2. T. Yaying, B. Hazarika, B. Chandra Tripathy, and M. Mursaleen. The spectrum of second order quantum difference operator. *Symmetry*, 14(3):557, 2022.
3. H. Bilgin Ellidokuzoğlu and S. Demiriz. Some New Paranormed Sequence Spaces Derived by q -Second Difference Matrix. *JOURNAL OF MATHEMATICAL EXTENSION*, vol.17, no.10, 1-18, 2023.

*Abstract Submission should be prepared only **1 page**.

PARANORMED NARAYANA SEQUENCE SPACES

Hacer Bilgin Ellidokuzoğlu¹ and Serkan Demiriz²

¹Department of Mathematics, University of Recep Tayyip Erdoğan, Rize, Türkiye
hacer.bilgin@erdogan.edu.tr

²Department of Mathematics, University of Tokat Gaziosmanpaşa, Tokat, Türkiye
serkandemiriz@gmail.com

Abstract

In the present paper, we construct the extended versions of Maddox's original paranormed sequence spaces, denoted by the notations $c_0(N, p)$ and $c(N, p)$. We investigate the topological structures and establish α -, β - and γ - duals of these spaces. Furthermore, the matrix transformations between these spaces the basic sequence spaces $c_0(p)$ and $c(p)$ are characterized.

Keywords: Narayana sequence spaces; paranormed sequence spaces; matrix transformations

REFERENCES

1. J. Allouche, T. Johnson, *Narayana's cows and delayed morphisms' cahiers du GREYC*, Troisiemes Journées d'Informatique Musicale (JIM 96) 4.
2. Y. Soykan, On generalized Narayana numbers, *International Journal of Advances in Applied Mathematics and Mechanics* 7 (3) (2020) 43–56.
3. M. İlkhani, S. Demiriz, and E. E. Kara. A new paranormed sequence space defined by Euler totient matrix. *Karaelmas Science and Engineering Journal*, 9(2):277–282, 2019.

*Abstract Submission should be prepared only **1 page**.

ON THE OSCILLATION OF A SECOND ORDER DIFFERENTIAL EQUATION WITH A SUPERLINEAR NEUTRAL TERM

Hasan Karaçalı¹ and Orhan Özdemir²

¹ Institute of Graduate Studies, Tokat Gaziosmanpasa University, 60240, Tokat, Turkey

hasan.karacalli@yahoo.com,

² Department of Mathematics, Tokat Gaziosmanpasa University, 60240, Tokat, Turkey

orhanozdemir37@yahoo.com,

Abstract

In this work, we focus on a class of second-order functional differential equations containing a superlinear neutral term with both delayed and advanced arguments. By utilizing integral averaging technique, we construct some new oscillation criteria for the considered differential equation. The results obtained in this work generalize and extend some of the known results in the literature. Illustrative examples are also provided to show applicability of the results.

Keywords: Oscillation; second order; neutral differential equations; superlinear.

REFERENCES

1. Martin Bohner, Balakrishnan Sudha, Krishnan Tangavelu, Ethiraju Thandapani, Oscillation criteria for second-order differential equations with superlinear neutral term, Nonlinear Studies, Vol:26, No:2, 425–434, 2019.
2. Orhan Özdemir, Ayla Kılıç, Oscillation of second-order functional differential equations with superlinear neutral terms, Bulletin of the Malaysian Mathematical Sciences Society, Vol:45, 83–99, 2022.
3. Ravi P. Agarwal, Said R. Grace, Donal O'Regan, Oscillation criteria for certain nth order differential equations with deviating arguments, Journal of Mathematical Analysis and Applications, Vol:262, 601–622, 2001.

Estimates of Bivariate New Kantorovich Type of the Balázs-Szabados Operators Based on q -integers

Hayatem Hamal¹

¹ Department of Mathematics, Tripoli University, Tripoli 22131, Libya

hafraj@yahoo.com

Abstract

In this paper, we introduce the bivariate q -kantorovich Balázs-Szabados operators of the tensor product class as follows;

$$R_n^{q_1} R_m^{q_2} (f, x, y) = \sum_{k=0}^n \sum_{l=0}^m s_{n,k}(q_1, x) r_{m,l}(q_2, y) \int_0^1 f\left(\frac{[k]_{q_1} + q_1^k t}{b_n}, \frac{[l]_{q_2} + q_2^l t}{b_m}\right) d_{q_1, q_2} t, \quad (1)$$

$f: \mathbb{R}_+ \times \mathbb{R}_+ \rightarrow \mathbb{R}$ is a continuous function $(x, y) \in \mathbb{R}_+ \times \mathbb{R}_+$ and $b_n = [n]_{q_1}^{\beta_1}$, $b_m = [m]_{q_2}^{\beta_2}$, $a_n = [n]_{q_1}^{\beta_1-1}$, $a_m = [m]_{q_2}^{\beta_2-1}$ are sequences for all $n, m \in \mathbb{N}$ such that $0 < q_1 \leq 1$, $0 < q_2 \leq 1$ and

$0 < \beta_1 \leq 2/3$ and $0 < \beta_2 \leq 2/3$. As well,

$$s_{n,k}(q, x) = \frac{1}{(1 + a_n x)^n} \begin{bmatrix} n \\ k \end{bmatrix}_{q_1} (a_n x)^k \prod_{s=0}^{n-k-1} (1 + (1 - q_1)[s]_{q_1} a_n x)$$

and

$$r_{m,l}(q, y) = \frac{1}{(1 + a_m y)^m} \begin{bmatrix} m \\ l \end{bmatrix}_{q_2} (a_m y)^l \prod_{j=0}^{m-l-1} (1 + (1 - q_2)[j]_{q_2} a_m y).$$

Let $e_{ij}(x, y) = x^i y^j$ for $0 \leq i + j \leq 2$, $i, j \in \mathbb{N}$ be the test functions, we estimate the moments and central moments of these new operators $R_n^{q_1} R_m^{q_2}$.

Keywords: Bivariate q -integer; q -Balázs-Szabados operators; Kantorovich theorem

REFERENCES

1. Hamal. H, Sabancıgil, Some Approximation Properties of new Kantorovich type q -analogue of Balázs-Szabados Operators, Journal of Inequalities and Applications, Vol:159, 2020.

NEW EXACT WAVE SOLUTIONS OF THE NEW HAMILTONIAN AMPLITUDE EQUATION THROUGH $(m + 1/G')$ -EXPANSION METHOD

Melik Şenyuva¹, Özlem Kirci², Hasan Bulut³

^{1,3} Department of Mathematics, Firat University, Elazığ, Turkey

² Department of Mathematics, Kırklareli University, Kırklareli, Turkey

mlksnyv@gmail.com, ozlem.isik@klu.edu.tr, hbulut@firat.edu.tr,

Abstract

In this paper, the new Hamiltonian amplitude equation (NHAЕ) is considered to investigate the new exact wave solutions via the $(m + 1/G')$ -expansion method. This model has become very important due to being useful for understanding the behaviors of the traveling waves in the field of applied sciences including fluids, plasmas and optics. The proposed method enables to obtain new complex solutions to NHAЕ which are also verified by Mathematica. Finally, 3D and 2D graphs are presented for the obtained exact solutions. These solutions are also compared with the results reported in the literature before.

Keywords: $(m + 1/G')$ -expansion method; new Hamiltonian amplitude equation; Traveling wave solution.

REFERENCES

1. Zafar, A., Raheel, M., Ali, K. K., Razzaq, W., On optical soliton solutions of new Hamiltonian amplitude equation via Jacobi elliptic functions, The European Physical Journal Plus, Vol: 135, No: 8, 674, 2020.
2. Wang, K. L., New perspective on fractional hamiltonian amplitude equation, Optical and Quantum Electronics, Vol: 55, No: 12, 1033, 2023.
3. Wang, M., Li, X., Applications of F-expansion to periodic wave solutions for a new Hamiltonian amplitude equation, Chaos, Solitons & Fractals, Vol: 24, No: 5, 1257-1268, 2005.
4. Tuluçe Demiray, S., Bulut, H., New exact solutions of the new Hamiltonian amplitude equation and Fokas-Lenells equation, Entropy, Vol: 17, No: 9, 6025-6043, 2015.
5. Kumar, S., Singh, K., Gupta, R. K., Coupled Higgs field equation and Hamiltonian amplitude equation: Lie classical approach and (G'/G) -expansion method, Pramana, Vol: 79, 41-60, 2021.
6. Tarla, S., Ali, K. K., Yusuf, A., Exploring new optical solutions for nonlinear Hamiltonian amplitude equation via two integration schemes, Physica Scripta, Vol: 98, No:9, 2023.

OSCILLATION OF SECOND ORDER NEUTRAL EMDEN-FOWLER DIFFERENTIAL EQUATIONS

Orhan Özdemir¹ and Ercan Tunç¹

¹ Department of Mathematics, Tokat Gaziosmanpasa University, 60240, Tokat, Turkey

orhanozdemir37@yahoo.com,

ercantunc72@yahoo.com

Abstract

The study of the asymptotic and oscillatory behaviours of solutions of functional Emden-Fowler differential and dynamic equations on time scales is an active and important area of qualitative theory. This rapidly growing interest in examining such equations is motivated by their applications in engineering and natural sciences. In this work, we give some new oscillation criteria that guarantee to be oscillatory of all solutions of second-order neutral Emden-Fowler differential equations. The results obtained are based on the comparisons of a second-order neutral differential equation with the corresponding first-order functional differential inequality and first-order functional differential equation. Some illustrative examples are also provided to show applicability of the main results.

Keywords: Oscillation; second order; Emden-Fowler; neutral differential equations.

REFERENCES

1. Ravi P. Agarwal, Martin Bohner, Tongxing Li, Chenghui Zhang, Oscillation of second-order Emden-Fowler neutral delay differential equations, *Annali di Matematica Pura ed Applicata*, Vol:193, 1861–1875, 2014.
2. Ercan Tunç, Orhan Özdemir, On the oscillation of second-order half-linear functional differential equations with mixed neutral term, *Journal of Taibah University for Science*, Vol:13, No:1, 481–489, 2019.
3. Orhan Özdemir, Oscillation criteria for noncanonical neutral Emden-Fowler differential equations, *Quaestiones Mathematicae*, Vol:46, No:2, 1653–1668, 2023.

Improved oscillation criteria for third-order half-linear delay differential equations via canonical transform

Ercan Tunç¹

¹Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Turkey

ercantunc72@yahoo.com,

Abstract

This paper discusses the oscillatory behavior of solutions to a class of third-order half-linear delay differential equations via canonical transform. Sufficient conditions for all solutions to be oscillatory are established. The results obtained here are new, and they improve and complement many known oscillation criteria in the literature. Examples are provided to illustrate the results and suggestions for future research are included.

Keywords: Oscillation; half-linear differential equations; delay; third-order.

REFERENCES

1. G. E. Chatzarakis, J. Dzurina and I. Jadlovská, Oscillatory and asymptotic properties of third-order quasilinear delay differential equations, *Journal of Inequalities and Applications*, Vol:2019, No:23, 1—17, 2019.
2. S. R. Grace, R. P. Agarwal, R. Pavan and E. Thandapani, On the oscillation of certain third order nonlinear functional differential equations, *Applied Mathematics and Computation*, Vol:202, No:1, 102—112, 2008.
3. K. Saranya, V. Piramanantham, E. Thandapani and E. Tunç, Asymptotic behavior of semi-canonical third-order nonlinear functional differential equations, *Palestine Journal of Mathematics*, Vol:11, No:3, 433—442, 2022.

Applications of the SUPG- $YZ\beta$ finite element formulation: from mussel-algae interactions to Schnakenberg reaction models

Süleyman Cengizci¹

¹ Computer Programming, Antalya Bilim University, Antalya 07190, Turkey

suleyman.cengizci@antalya.edu.tr

Abstract

Coupled systems of reaction-convection-diffusion equations can model a wide range of phenomena from science, industry, and nature. Unfortunately, analytical solutions to such systems can rarely be obtained, and, as a result, their analysis usually necessitates the application of numerical techniques. However, when convection dominates the transport process, classical computational tools yield approximations polluted with physically meaningless oscillations, since solutions to such systems can exhibit rapid changes and may have boundary or inner layers. Therefore, this study utilizes a stabilized finite element formulation, the so-called streamline-upwind/Petrov–Galerkin (SUPG) formulation, to eliminate such numerical instabilities without sacrificing accuracy. In order to obtain better approximations around steep gradients, the $YZ\beta$ shock-capturing mechanism is then incorporated into the SUPG-stabilized formulation. To demonstrate the robustness of the proposed formulation, which we refer to as the SUPG- $YZ\beta$ formulation, a number of numerical test experiments are performed, encompassing cross-diffusion systems, the Schnakenberg reaction model, and mussel-algae interactions. The proposed formulation performs quite well without introducing excessive numerical dissipation, according to comparisons with reported studies.

Keywords: Reaction-convection-diffusion, Finite elements, Shock-capturing, SUPG- $YZ\beta$.

REFERENCES

1. Suleyman Cengizci, An enhanced SUPG-stabilized finite element formulation for simulating natural phenomena governed by coupled system of reaction-convection-diffusion equations, *Mathematical Modelling and Numerical Simulation With Applications*, Vol:3, No: 4, 297–317, 2023.
2. Yuri Bazilevs, Victor M. Calo, Tayfun E. Tezduyar, and Thomas J. R. Hughes, $YZ\beta$ discontinuity capturing for advection-dominated processes with application to arterial drug delivery, *International Journal for Numerical Methods in Fluids*, Vol: 54, 593–608, 2007.

SOME NEW ANALYTICAL SOLUTIONS TO THE NONLINEAR MODIFIED QUANTUM ZAKHAROV-KUZNETSOV EQUATION

Haci Mehmet Baskonus^{1*}, Halil Şenççek²

¹ Department of Mathematics and Science Education, Harran University, Şanlıurfa, Türkiye

² Department of Mathematics and Science Education, Harran University, Şanlıurfa, Türkiye

hmbaskonus@gmail.com, halilsencicek@gmail.com

*Corresponding Author

Abstract

In this paper, we obtain some new analytical solutions to the nonlinear modified Quantum Zakharov-Kuznetsov equation (MQZKE) by using several powerful analytical schemes. We obtain some important properties of the governing model. Various simulations are also plotted by using Computational package programs.

Keywords: Exponential method; modified Quantum Zakharov-Kuznetsov equation; Complex exponential solution.

REFERENCES

1. Areshi, M.; Seadawy, A.R.; Ali, A.; AlJohani, A.F.; Alharbi, W.; Alharbi, A.F. Construction of Solitary Wave Solutions to the (3+1)-Dimensional Nonlinear Extended and Modified Quantum Zakharov-Kuznetsov Equations Arising in Quantum Plasma Physics. *Symmetry* 2023, 15, 248. <https://doi.org/10.3390/sym15010248>
2. Kumar, S., Optimal system, dynamical behaviors and exact solution of a nonlinear transmission line model by applying the Lie symmetry method, *Indian J Phys* <https://doi.org/10.1007/s12648-022-02327-x>
3. G. Yel, H. M. Baskonus, H. Bulut, Novel archetypes of new coupled Konno-Oono equation by using sine-Gordon expansion method, *Opt. Quant. Electron.*, 49 (2017), 1–10.
4. H. Bulut, T. A. Sulaiman, H. M. Baskonus, New solitary and optical wave structures to the Korteweg-de Vries equation with dual-power law nonlinearity, *Opt. Quant. Electron.*, 48 (2016), 1–14.

Comparative Analysis of Rankings and Preference Values for Fuzzy Decision-Making Approaches in Reducing Unnecessary Biopsies

Orhan Dalkılıç¹, Esin İlhan², Hasan Bulut³

¹ Mersin City Hospital, Mersin, Turkey

² Faculty of Engineering and Architecture, University of Kırşehir Ahi Evran, Elazığ, Turkey

³ Department of Mathematics, University of Fırat, Elazığ, Turkey

orhandlk952495@hotmail.com

Abstract

Prostate cancer is a significant health concern, and accurate diagnosis plays a crucial role in determining appropriate treatment strategies. However, the high number of unnecessary biopsies in prostate cancer diagnosis poses challenges in terms of patient discomfort and healthcare costs. This study aims to address this issue by employing fuzzy decision-making approaches, including Fuzzy MOORA, Fuzzy TOPSIS, Fuzzy VIKOR and Fuzzy WASPAS, to reduce the number of unnecessary biopsies. The application of fuzzy set theory allows us to handle uncertainties and imprecisions inherent in prostate cancer diagnosis. Fuzzy MOORA is utilized to rank the criteria, fuzzy TOPSIS evaluates diagnostic alternatives, fuzzy VIKOR provides a compromise solution and fuzzy WASPAS determines the weights of criteria. Successful results have been achieved, demonstrating the effectiveness of these methods in reducing unnecessary biopsies. This research contributes to improving the efficiency and accuracy of prostate cancer diagnosis, leading to enhanced patient care, reduced healthcare costs and minimized patient discomfort.

Keywords: Fuzzy set; Fuzzy TOPSIS; Fuzzy VIKOR; Fuzzy MOORA; Fuzzy WASPAS; Multi-criteria decision-making.

REFERENCES

1. Mardani, A., Nilashi, M., Zakuan, N., Loganathan, N., Soheilrad, S., Saman, M. Z. M., & Ibrahim, O., A systematic review and meta-Analysis of SWARA and WASPAS methods: Theory and applications with recent fuzzy developments. *Applied Soft Computing*, Vol:57, 265-292, 2017
2. Nădăban, S., Dzitac, S., Dzitac, I., Fuzzy TOPSIS: a general view. *Procedia computer science*, Vol:91, 823-831, 2016.
3. Zadeh, L.A., Fuzzy sets. *Information and Control*, Vol:8, 338-353, 1965.

ESTIMATIONS OF INTEGRAL MAJORIZATION INEQUALITY FOR DIFFERENTIABLE CONVEX FUNCTIONS AND APPLICATIONS

Artion Kashuri¹, Rozana Liko²

¹Department of Mathematical Engineering, Polytechnic University of Tirana,
Tirana 1001, Albania

²Department of Mathematics, Faculty of Technical and Natural Sciences,
University Ismail Qemali, 9400 Vlora, Albania
a.kashuri@fimif.edu.al, rozana.liko@univlora.edu.al

Abstract

This study aims to estimate integral majorization inequalities by taking third-differentiable convex functions. By using Hölder, Power-mean and Jensen inequalities, we derive some new relations involving integral majorization. Additionally, our study explores various applications in information theory, including estimations for the Csiszár and Kullback–Leibler divergences, Shannon entropy, and Jeffrey’s distance.

Keywords: Convex function; Majorization inequality; Hölder's inequality; Power-mean inequality; Jensen's inequality; Information theory.

REFERENCES

1. Artion Kashuri, Rozana Liko, Some new Hermite-Hadamard type inequalities and their applications, *Studia Scientiarum Mathematicarum Hungarica*, Vol:56, 103-142, 2019.
2. Josip Pečarić, A companion inequality to Jensen–Steffensen’s inequality, *Journal of Approximation Theory*, Vol:44, 289–291, 1985.
3. Albert W. Marshall, Ingram Olkin, Barry C. Arnold, *Inequalities: Theory of majorization and its applications*, 2nd ed., Springer Series in Statistics, Springer, 2011.
4. Silvestru Sever Dragomir, Some majorization type discrete inequalities for convex functions, *Mathematical Inequalities and Applications*, Vol:7, 207–216, 2004.
5. Shan-He Wu, Muhammad Adil Khan, Hidayat Ullah Haleemzai, Refinements of majorization inequality involving convex functions via Taylor’s theorem with mean value form of the remainder, *Mathematics*, Vol:7, No:8, 663, 2019.
6. Abdul Basir, Muhammad Adil Khan, Hidayat Ullah, Yahya Almalki, Chanisara Metpattarahiran, Thanin Sitthiwiratttham, Improvements of integral majorization inequality with applications to divergences, *Axioms*, Vol:13, No:1, 21, 2024.

A SURVEY ON DIFFERENT STATISTICAL DISTRIBUTIONS USING PYTHON PROGRAMMING

Gözde Karataş Baydoğmuş¹, Şahsene Altinkaya²

¹Department of Computer Engineering, Marmara University, Istanbul, Türkiye

gkaratas@marmara.edu.tr,

²Department of Mathematics, Istanbul Beykent University, Istanbul, Türkiye

sahsenealtinkaya@beykent.edu.tr

Abstract

Number of 'data/information' is increasing rapidly around the world and is included in every aspect of our lives. Those interested in data science know the importance of statistical methods and aim to make improvements accordingly. In this study, the importance of 5 popular statistical distribution methods which are normal distribution, gaussian distribution, bernoulli distribution, binom distribution, and poisson distribution were examined. Accordingly, it was aimed to examine the working principles of the selected 5 statistical methods. Results obtained with traditional methods and the scipy library were evaluated using the Python programming language. In this study, run time and error rate of statistical distributions were examined on traditional methods and Python scipy libraries, and their differences were evaluated.

Keywords: Normal distribution, Gaussian distribution, Bernoulli distribution, Binom distribution, Poisson distribution.

REFERENCES

1. Fatma Hilal Yağın, Emek Göldoğan, Cemil Çolak, A Web-based Software for the Calculation of Theoretical Probability Distributions, The Journal of Cognitive Systems, Vol:6, No:1, 44-50, 2021.
2. Thomas Haslwanter, An Introduction to Statistics with Python. *Statistics and Computing*, Springer, Cham., 2022.

EXPLORING MACHINE LEARNING TECHNIQUES FOR GENDER VOICE RECOGNITION USING LIMITED SPEECH DATA

Gözde Karataş Baydoğan¹, Şahsene Altinkaya², Nahide Zeynep Cicekli³

¹Department of Computer Engineering, Marmara University, Istanbul, Türkiye

gkaratas@marmara.edu.tr,

²Department of Mathematics, Istanbul Beykent University, Istanbul, Türkiye

sahsenealtinkaya@beykent.edu.tr

³Nobel Ilac, Istanbul, Turkey

nzeynecicekli@gmail.com

Abstract

Voice recognition has gained popularity, leading to extensive research in the artificial intelligence field, primarily utilizing machine learning algorithms. However, determining effective voice recognition algorithms for small datasets remains really hard challenge. This study focuses on the significance of machine learning algorithms in gender voice recognition. The study examines the impact of machine learning algorithms on a dataset of number of 3168 data and 21 features. Machine learning algorithms used in this study are decision tree, xgboost, lightgbm, adaboost, gradient boosting, and k-nearest neighbor. These learning algorithms, especially boosting methods known for their success, were chosen for this study. Experimental results show that machine learning algorithms achieve successful performance rates particularly the Gradient Boosting algorithm. At the end study reached 98.26\% accuracy rate while deep learning approaches in existing literature achieved an accuracy rate of 97\%. As a result of the study, it was seen that a high performance rate was achieved by using machine learning algorithms in small-sized datasets.

Keywords: Voice recognition; Gender; Machine learning; Sound processing.

REFERENCES

1. Abeer Ali Alnuaim, Mohammed Zakariah, Chitra Shashidhar, Wesam Atef Hatamleh, Hussam Tarazi, Prashant Kumar Shukla, Rajnish, Speaker gender recognition based on deep neural networks and ResNet50, Wireless Communications and Mobile Computing, 2022, 1-13.

On Lacunary Statistical Boundedness of Sequences of Sets

Yaşar ÇALIŞKAN¹ and Mikail ET²

¹Fırat University, Elazığ, Turkey, yasar_2344@hotmail.com

²Fırat University, Elazığ, Turkey, mikail68@gmail.com

Abstract The concept of statistical convergence was introduced by Steinhaus [8] and Fast [2], and later reintroduced by Schoenberg [7] independently. Later on it was further investigated from the sequence space point of view and linked with summability theory by Et and Arsalanoğlu [1], Fridy [3], Fridy and Orhan [4], Nuray and Rhoades [5], Salat [6] and many others. In this study we introduce and examine the concepts of Wijsman lacunary statistical boundedness and give some relations between Wijsman lacunary statistical convergence and Wijsman lacunary statistical boundedness.

Keywords. Density, Statistical Convergence, Statistical Boundedness, Sequences of Sets, Wijsman convergence

REFERENCES

- [1] M. Et and M. Arslanoğlu, On f-lacunary statistical convergence of order α of sequences of sets, AIP Conference Proceedings 2037, 020009 (2018); doi: 10.1063/1.5078464.
- [2] H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951) 241-244.
- [3] J.A. Fridy, On statistical convergence, Analysis 5 (1985), no. 4, 301--313.
- [4] J.A. Fridy, C: Orhan, Statistical limit superior and limit inferior, Proc. Amer. Math. Soc. 125(12) (1997), 3625--3631.
- [5] F. Nuray and B.E. Rhoades, Statistical convergence of sequences of sets, Fasc. Math. No. 49 (2012), 87--99.
- [6] T. Salat, On statistically convergent sequences of real numbers, Math. Slovaca 30 (1980), 139-150.
- [7] I.J. Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- [8] H. Steinhaus, Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951), 73-74.

On λ -Statistical Boundedness of Sequences of Sets

Ayşe EREN¹ and Mikail ET²

¹Firat University, Elazığ, Turkey, gozukaraayse@gmail.com

²Firat University, Elazığ, Turkey, mikail68@gmail.com

Abstract The concept of statistical convergence was introduced by Steinhaus [8] and Fast [2], and later reintroduced by Schoenberg [7] independently. Later on it was further investigated from the sequence space point of view and linked with summability theory by Et and Arsalanoğlu [1], Fridy [3], Mursaleen [4], Nuray and Rhoades [5], Salat [6] and many others. In this study we introduce and examine the concepts of Wijsman λ -statistical boundedness and give some relations between Wijsman λ -statistical convergence and Wijsman λ -statistical boundedness.

Keywords. Density, Statistical Convergence, Statistical Boundedness, Sequences of Sets, Wijsman convergence

REFERENCES

- [1] M. Et and M. Arslanoğlu, On f -lacunary statistical convergence of order α of sequences of sets, AIP Conference Proceedings 2037, 020009 (2018); doi: 10.1063/1.5078464.
- [2] H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951) 241-244.
- [3] J.A. Fridy, On statistical convergence, Analysis 5 (1985), no. 4, 301--313.
- [4] Mursaleen M. λ -statistical convergence. Math. Slovaca. 2000;50(1):111-115..
- [5] F. Nuray and B.E. Rhoades, Statistical convergence of sequences of sets, Fasc. Math. No. 49 (2012), 87--99.
- [6] T. Salat, On statistically convergent sequences of real numbers, Math. Slovaca 30 (1980), 139-150.
- [7] I.J. Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- [8] H. Steinhaus, Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951), 73-74.

INEXTENSIBLE FLOWS OF CURVES WITH QUASI-FRAME IN GALILEAN SPACE G_3

Murat Turan¹, Hülya Gün Bozok² and Mahmut Ergüt³

^{1,2} Department of Mathematics, University of Osmaniye Korkut Ata, Osmaniye, Türkiye

³ Department of Mathematics, University of Tekirdağ Namık Kemal, Tekirdağ, Türkiye

muratturan@osmaniye.edu.tr, hulyagun@osmaniye.edu.tr and mergut@nku.edu.tr

Abstract

In this study we research inextensible flows of curve in 3-dimensional Galilean space G_3 with a new aspect. For this research we use a new adapted frame which called quasi-frame in 3-dimensional Galilean space G_3 . From this perspective, inextensible curve flows are examined with the help of this frame then important characterizations and results are obtained.

Keywords: Galilean space, inextensible flows of curves, quasi frame.

REFERENCES

1. Elsharkawy A., Tashkandy Y., Emam W., Cesarano C., Elsharkawy N., On Some Quasi-Curves in Galilean Three-Space, Axioms, 12, 823, 2023.
2. Elshenhab M.A., Moaaz O., Dassios I., Elsharkawy A., Motion along a Space Curve with a Quasi-Frame in Euclidean 3-Space: Acceleration and Jerk, Symmetry, 14, 1610, 2022.
3. Öğrenmiş A.O., Yeneroğlu M., Inextensible curves in the Galilean Space, International Journal of the Physical Sciences 5(9), 1424-1427, 2010.
4. Öztekin H. and Gün Bozok H., Inextensible flows of curves in 4-dimensional Galilean space G_4 , Math.Sci. Appl. E-Notes, 1(2), 28-34, 2013.

ON DIFFERENCES OF BOUNDED VARIATION FUZZY SEQUENCES

Murat Temizkan and Hıfı Altınok

Department of Mathematics, Firat University, Elazığ, Turkey

murattemizkanmat@gmail.com hifsialtinok@gmail.com

Abstract

Kızmaz [11] first introduced the concept of difference sequence in 1981. Afterwards, Et and Colak [8] generalized the difference sequences and examined some topological characteristics of the resultant sequence spaces. Matloka [13] introduced fuzzy number sequences and gave their basic properties. In this paper, we define the difference of a bounded variation fuzzy sequence by a lacunary sequence and examine some of their properties.

Keywords: Difference sequence; Lacunary sequence; Fuzzy sequence.

REFERENCES

1. Y. Altin, M. Mursaleen, H. Altınok, Statistical summability $(C,1)$ for sequences of fuzzy real numbers and a Tauberian theorem, *Journal of Intelligent & Fuzzy Systems*, 21(6) (2010), 379-384.
2. H. Altınok, Y. Altın and M. Et, Lacunary almost statistical convergence of fuzzy numbers, *Thai Jour. Math.* 2(2) (2004), 265-274.
3. H. Altınok, R. Çolak and M. Et, λ -Difference sequence spaces of fuzzy numbers, *Fuzzy Sets and Systems* 160(21) (2009), 3128-3139.
4. A.R. Freedman, I.J. Sember and M. Rapheal, Some Cesaro-type summability spaces, *Proc. London Math. Soc.* 37(3) (1973), 508-520.
5. J.A. Fridy and C. Orhan, Lacunary statistical convergence, *Pacific J. Math.* 160(1) (1993), 43-51.
6. H. Kızmaz, On certain sequence spaces, *Canad. Math. Bull.* 24(2) (1981), 169-176.
7. M. Matloka, Sequences of fuzzy numbers, *BUSEFAL* 28 (1986), 28-37.
8. B.C. Tripathy. and A.J. Dutta, Bounded variation double sequence space of fuzzy real numbers, *Comput. Math. Appl.* 59(2) (2010), 1031-1037.

ON LACUNARY STATISTICAL BOUNDEDNESS

Mithat KASAP¹ and Hıfı ALTINOK²

¹Department of Accounting, Sirnak University, Sirnak, Turkey

²Department of Mathematics, Firat University, Elazig, Turkey

fdd_mithat@hotmail.com hifsialtinok@gmail.com

Abstract

In the present paper, we introduce the concept of Δ^f -lacunary statistical boundedness of order β with respect to a modulus function f for sequences of fuzzy numbers and give some relations between Δ^f -lacunary statistical boundedness of order β and Δ^f -statistical boundedness with respect to a modulus function f .

Keywords: Lacunary sequence; Statistical boundedness; Modulus function; Fuzzy sequence.

REFERENCES

1. Aizpuru, A.; Listan-Garcia, MC and Rambla-Barreno, F. Density by moduli and statistical convergence. Quaest. Math. 37, (2014) 525-530.
2. Altinok, H. and Mursaleen, M. Δ -Statistical boundedness for sequences of fuzzy numbers, Taiwanese Journal of Mathematics, 15 (5), (2011), 2081-2093.
3. Altinok, H. Et, M. and Altin, Y. Lacunary statistical boundedness of order β for sequences of fuzzy numbers, J Intell Fuzzy Syst 35 (2018), 2383—2390
4. Aytar, S. and Pehlivan, S. Statistically monotonic and statistically bounded sequences of fuzzy numbers, Inform. Sci., 176(6) (2006), 734-744.
5. Bhardwaj, VK. and Dhawan, S. f -statistical convergence of order α and strong Cesaro summability of order α with respect to a modulus, J. Inequal. Appl. (2015) 2015:332 DOI 10.1186/s13660-015-0850-x.
6. Çolak, R. Statistical convergence of order α , Modern Methods in Analysis and Its Applications, New Delhi, India: Anamaya Pub, (2010) 121--129.
7. Freedman, A. R., Sember, J. J. and Raphael, M., Some Cesaro-type summability spaces, Proc. Lond. Math. Soc. 37(3) (1978), 508-520.
8. Matloka, M. Sequences of fuzzy numbers, BUSEFAL 28 (1986) 28-37.

ON A COEFFICIENT PROBLEM FOR FUNCTIONS BELONGS TO CERTAIN SUBCLASS OF UNIVALENT FUNCTIONS

Tuğba YAVUZ¹

¹Department of Mathematics, Istanbul Beykent University, Turkey

tugbayavuz@beykent.edu.tr,

Abstract

Let A denote the family of functions $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ analytic in the open unit disk $D = \{z: |z| < 1\}$ with the conditions $f(0) = f'(0) - 1 = 0$. If f does not take the same value twice, it is called univalent in D . For $-\infty < t < \infty$ and $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, the curve expressed by $w = w_0 \exp(-e^{-i\theta} t)$, where w_0 is a nonzero complex number, is called logarithmic θ -spiral. We know that 0-spirals are radial half line. For an analytic function, it is named as θ -spirallike if its range is θ -spirallike. Analytically, $f \in A$ belongs the class of S_θ if and only if $\operatorname{Re} \left(e^{-i\theta} \frac{zf'(z)}{f(z)} \right) > 0$.

In this study, the extreme points of certain subclasses of spirallike functions. By the applications of extreme point theory, we obtain sharp upper bounds for some nonlinear functionals defined in terms of functions in these classes.

Keywords: Spirallike Functions, Extreme Points, Subordination, Coefficient Estimates

REFERENCES

1. Suk Young Lee, Gae Sun Chung, "On extreme points of the family of spirallike functions", Commun. Korean Math. Soc., 8 (4), 603-615, 1993.
2. L. Spacek, Prispěvek k teorii funkcií prostých, Casopis Pěst. Mat. a Fys., 62 (1932), 12-19.
3. J. Zamorski, About the extremal spiral schlicht functions, Ann. Polon. Math., 9 (1962), 265-273.
4. Yavuz T, Altınkaya Ş. Notes on some classes of spirallike functions associated with the q-integral operator. Hacettepe Journal of Mathematics and Statistics, 53(1), 53-61, 2024.

New Wave Behaviors For Solutions Of The Truncated M-Fractional Variant Boussinesq System

İlkay Koçoğlu¹, Hasan Bulut²

¹ Department of Mathematics, Firat University, Elazig, Turkey,

²Department of Mathematics, Firat University, Elazig, Turkey,

* Corresponding author: ilkaykocoglu@icloud.com

Abstract

This manuscript focuses on New wave behaviors for solutions of the truncated M-fractional variant Boussinesq system which refers to a set of nonlinear partial differential equations used to model the behavior of waves in shallow water. This system is a modification or variant of the original Boussinesq equation. The goal of the study is to shed light on the model's underlying mathematical structures and contribute to a better understanding of the system for which is used to model waves in shallow water and electrical signals in telegraph lines based on tunnel diodes. The Sardar sub-equation method which is one of the powerful methods has been considered for finding exponential wave solutions of the system. Some 2D and 3D graphs were presented to explain the physical characteristics of variant Boussinesq system' solution. This algorithm yields new exponential function solutions to the system considered in this paper. Wolfram Mathematica 12 has been successfully used throughout the paper for mathematical calculations.

Keywords the variant Boussinesq system; Truncated M-fractional derivative; the Sardar sub-equation method; Soliton solutions.

References

- [1] G. Yel, H. Bulut, New wave approach to the conformable resonant nonlinear Schrödinger's equation with Kerr-law nonlinearity, *Opt. Quantum Electron.* 54 (2022) 1–13. <https://doi.org/10.1007/s11082-022-03655-2>.
- [2] M. Cinar, A. Secer, M. Ozisik, M. Bayram, Derivation of optical solitons of dimensionless Fokas-Lenells equation with perturbation term using Sardar sub-equation method, *Opt. Quantum Electron.* 54 (2022) 1–13. <https://doi.org/10.1007/s11082-022-03819-0>.
- [3] H.U. Rehman, I. Iqbal, S. Subhi Aiadi, N. Mlaiki, M.S. Saleem, Soliton Solutions of Klein–Fock–Gordon Equation Using Sardar Subequation Method, *Mathematics*. 10 (2022) 1–10. <https://doi.org/10.3390/math10183377>.
- [4] I. Onder, M. Cinar, A. Secer, A. Yusuf, M. Bayram, T.A. Sulaiman, Comparative Analysis for the Nonlinear Mathematical Equation with New Wave Structures, *Eur. Phys. J. Plus* 137 (2022) 1120. <https://doi.org/10.1140/epjp/s13360-022-03342>.
- [5] J.V.D.A.C. Sousa, E.C.D.E. Oliveira, A New Truncated M-Fractional Derivative Type Unifying Some Fractional Derivative Types with Classical Properties, *Int. J. Anal. Appl.* 16 (2018) 83–96. <https://doi.org/10.28924/2291-8639-16-2018-83>.

A Collocation Method for Numerical Solution of Linear Integro-Differential Equations by Stancu Polynomials

Neşe İşler Acar¹

¹Department of Mathematics, University of Mehmet Akif Ersoy, Burdur, Turkey

nisler@mehmetakif.edu.tr,

Abstract

In this paper, an alternative collocation method has been produced for numerical solution of linear Fredholm and Volterra type integro-differential equations. As the proposed method has been introduced, the Stancu polynomials that are generalization of the Bernstein polynomials and their algebraic properties have been used for the proposed method. Some examples of the Fredholm and Volterra integro-differential equations have been considered in order to indicate applicability of the method. Likewise, the numerical results of the proposed method have been presented as tables. Moreover, the numerical results of the proposed method have been compared with the numerical results of the Bernstein collocation method to show the how much the proposed method is efficient.

Keywords: Collocation Method; Stancu Polynomials; linear integro-differential equations; matrix equations.

REFERENCES

1. Stancu D.D., Approximation of functions by a new class of linear polynomial operators, Rev. Roumaine Math. Pure Appl., Vol:1968, No:13, 1173-1194, 1968.
2. Stancu, D.D., Approximation of functions by means of a new generalized Bernstein operatör, Calcolo, Vol:1983, No:20, 211-229, 1983.
3. Bernstein, S., Démonstration du théorème de Weierstrass Fondée sur le calcul des probabilités, Commun. Soc. Math. Kharkow, Vol:13, No:1, 1-2, 1912.
4. İşler Acar, N., Daşcıoğlu, A., A projection method for linear Fredholm-Volterra integro-differential equations, Journal of Taibah University for Science, Vol:13, No:1, 644-650, 2019. doi.org/10.1080/16583655.2019.1616962.

APPROXIMATION PROPERTIES OF KANTOROVICH TYPE SAMPLING SERIES IN WEIGHTED SPACES OF FUNCTIONS

Metin Turgay¹

¹Department of Mathematics, Selçuk University, Konya, Turkey

metinturgay@yahoo.com

Abstract

In this talk, we study Kantorovich type sampling series in weighted spaces of functions. We investigate the approximation properties of the newly introduced operators, presenting convergence results and providing a quantitative form of the convergence via weighted modulus of continuity.

Keywords: Sampling type series; weighted approximation, Voronovskaja type theorem.

**The author have been supported within TUBITAK (The Scientific and Technological Research Council of Turkey) 1001-Project 123F123.*

REFERENCES

- T. Acar, M. Turgay, Approximation by bivariate generalized sampling series in weighted spaces of functions, Dolomites Res. Notes Approx., 16 (2023), 11--22.
- T. Acar, O. Alagoz, A. Aral, D. Costarelli, M. Turgay, G. Vinti, Convergence of generalized sampling series in weighted spaces, Demonstr. Math., 55, (2022), 153-162.
- O. Alagoz, M. Turgay, T. Acar, M. Parlak, Approximation by Sampling Durrmeyer Operators in Weighted Space of Functions, Numerical Functional Analysis and Optimization, 43 (10), (2022), 1223-1239.
- T. Acar, O. Alagoz, A. Aral, D. Costarelli, M. Turgay, G. Vinti, Approximation by sampling Kantorovich series in weighted spaces of functions, Turk. J. Math., 46 (7) (2022), 2663-2676.
- T. Acar, D. Costarelli, G. Vinti. Linear prediction and simultaneous approximation by m-th order Kantorovich type sampling series, Banach J. Math. Anal., 14 (2020) 1481-1508.

Analytical Solution of Hirota Equation by Rational Sine-Gordon Method

Beyhan Kemalolu¹, Gülnur Yel², Hasan Bulut¹

²Department of Mathematics, Firat University, Elazig, Turkey

¹ Faculty of Education, Final International University, Kyrenia, Mersin 10, Turkey

beyhanozturk1980@gmail.com, gulnur.yel@final.edu.tr, hbulut@firat.edu.tr

Abstract

In this work, we consider the Hirota equation [1-3],

$$iu_t + u_{xx} + 2|u|^2 u + i\alpha u_{xxx} + 6i\alpha |u|^2 u_x = 0$$

The Hirota equation shows the pulses that are physically generated by the help of solitons per unit second in single special fibers. We define the complex form amplitude of the optical field over time with $u = u(x,t)$. Here α is a sufficiently small parameter. Also, and $u_t, u_{xx}, |u|^2 u, u_{xxx}$ and $|u|^2 u_x$ terms represent linear progression, For this equation, we used an analytical method which is the rational sine-Gordon expansion method [4]. By using the proposed method, some different wave solutions are achieved. We discussed the physical dynamics of all obtained solutions with respect to graphical simulations.

Keywords: The Rational sine-Gordon expansion method; Hirota equation; dark-bright solitary waves

References

1. Gao W., Rezazadeh H., PinarZ., Baskonus H. B., Sarwar S. and Yel G., Novel explicit solutions for the nonlinear Zoomeron equation by using newly extended direct algebraic technique. *Optical and Quantum Electronics*, 52:52, 2020
2. Eslami, M., Mirzazadeh, M.A. Neirameh, A., (2015) New exact wave solutions for Hirota equation, *Pramana J. of Physics*, 84(1), 3-8. c 2016 BSKA Bilis Abazari, R.: The solitary wave solutions of Zoomeron equation. *Appl. Math. Sci.* 5(59), 2943–2949, 2011
3. Jia TT, Chai YZ, Hao HQ. (2017) Multi-soliton solutions and breathers for the generalized coupled nonlinear Hirota equations via the Hirota method. *Superlattices Microstruct* ;105:172–82
4. Yel G., Bulut, H. New wave approach to the conformable resonant nonlinear Schödinger's equation with Kerr-law nonlinearity, *Optical and Quantum Electronics*, 54(4), (2022).

BLOCKCHAIN APPLICATIONS IN MEDULA SYSTEM

Burak Çevik¹ Muharrem Tuncay Gençoğlu²

¹ SGK Provincial Directorate, Elazığ, Türkiye

² Fırat University, Elazığ, Türkiye

kingleon23@hotmail.com,

Abstract

Medical Messenger System has a very important place in today's healthcare industry. The system includes reporting, e-prescription, HIS (Hospital Information Management System), e-medical, and e-pharmacy drug systems. All these systems are open to security and privacy violations; Ensuring the security of the data it contain is mandatory by patient privacy and social security institution payments. There are constant cyber attacks against the Medula System. Cyber attacks will cause e-signature, e-reporting, and e-pharmacy systems to fall into the hands of unauthorized persons. This poses both a danger to the patient's life and financial damage to the state, as it may cause patient data to be changed.

In this study, is planned to make the Medula System more reliable, the control architecture more unique, the patient interaction power faster, and the access to both medicine and medical infrastructure more economical by using Blockchain. It is anticipated that the Medula System will provide a more reliable, more economical, and more ergonomic infrastructure to the healthcare system by taking advantage of the superior features of Blockchain.

Keywords: Blockchain, Medula systems

REFERENCES

1. PanWang, BoTian, Kun Sun, Feng-Hua Qi, Thetitle of Paper, Applied Mathematics and Computation, Vol:2, No:1, 233–242, 2015.

Crank-Nicolson Finite Difference Treatment of Time Fractional Klein Gordon Equation

Muhammed Huzeyfe Uzunyol¹, Berat Karaagac², Alaattin Esen¹

¹ Department of Mathematics, University of Inonu, Malatya, Turkey

² Department of Mathematics Education, University of Adiyaman, Adiyaman, Turkey

mhuzeyfeuzunyol@hotmail.com, bkaraagac@inonu.edu.tr, alaattin.esen@inonu.edu.tr

Abstract

This paper explores treatment of Klein Gordon equation whose time derivative has Caputo meaning fractional derivative via Crank Nicolson finite difference method. The proposed technique relies on the obtaining an algebraic equation using L2 algorithm for time fractional derivative, Crank Nicolson and finite difference approach for spatial derivatives. The obtained results are compared with exact ones and presented for the different values of fractional order α ($1 \leq \alpha \leq 2$). Graphical findings depict the geometric behavior of the approximation solutions for different values of fraction order.

Keywords: Time fractional Klein Gordon equation, Finite difference method, Caputo derivative

REFERENCES

1. Hu Chen, Lü Shujuan, Chen Wenping, A fully discrete spectral method for the nonlinear time fractional Klein-Gordon equation, Taiwanese Journal of Mathematics, 21.1, 231-251, 2017.
2. S Kumbinarasaiah, A new approach for the numerical solution for nonlinear Klein–Gordon equation, SeMA Journal, 77.4, 435-456, 2020.
3. Joaquín Quintana Murillo, Santos Bravo Yuste, an explicit difference method for solving fractional diffusion and diffusion-wave equations in the Caputo form, Journal of computational and nonlinear Dynamics, 6.2, 021014, 2011.
4. Keith Oldham, Jerome Spanier. The fractional calculus theory and applications of differentiation and integration to arbitrary order, Elsevier, 1974.

The Relations of Soft Topological Hyperstructures

Gülay OĞUZ¹, Abdülkadir Olcay¹

¹Department of Mathematics, Faculty of Arts and Sciences, Harran University, Şanlıurfa, Turkey

gulay.oguz@harran.edu.tr

kadirolcay@hotmail.com

Abstract

Soft sets are a mathematical framework that extend classical set theory to handle uncertainty and vagueness in data and decision-making processes. Soft sets were introduced by Molodtsov in 1999 as a generalization of classical sets. Soft sets are particularly useful in situations where precise membership information is not available, and instead, elements have degrees of membership or uncertainty associated with them. Soft sets have found applications in various fields, including decision-making, data analysis, and information fusion, where uncertainty and imprecision are inherent. One of the application areas of soft sets is topological hyperstructures, which play a very important role in mathematics. In this work, some notions and results concerning soft topological hypergroupoids and soft topological hypergroups are presented. Also, the relations between soft topological hypergroupoids and soft topological hypergroups are analysed, and relevant important conclusions are obtained.

Keywords: Soft Groups Soft topological hypergroupoids; Soft topological hypergroups.

REFERENCES

1. Molodtsov, D. Soft Set Theory First Results. Comput. Math. Appl. Vol: 37, 19–31, 1999.
2. Oguz, G. A New View on Topological Polygroups. Turkish Journal of Science, Vol: 5, No:2, 110—117, 2020.
3. Maji, P. K., Biswas, R. and Roy, R. An Application of Soft Sets In A Decision Making Problem. Computers and Mathematics with Applications, Vol: 44, 1077-1083, 2002.
4. Oguz, G. On soft topological hypergroups. Journal of Hyperstructures, Vol:9, No:2, 81-95, 2020.
5. Marty, F. Sur une generalization de la notion de group, 8th Congres Math. Scandinaves, Stockholm, 45-49, 1934.
6. Oguz, G., Davvaz, B. Soft topological hyperstructure. Journal of Intelligent and Fuzzy Systems, Vol:40, No:5, 8755-8764, 2021.

Rough Approximation Operators on Algebraic Hyperstructures

Gülay OĞUZ¹, Ayhan YÜKSEL¹

¹Department of Mathematics, Faculty of Arts and Sciences, Harran University, Şanlıurfa, Turkey

gulay.oguz@harran.edu.tr

ayhan_680@hotmail.com

Abstract

Rough set theory is an important mathematical tool for modeling uncertainty. This theory has reached a wide potential for study in many fields, especially mathematics, engineering, medicine and computer science. On the other hand, new structures have been introduced by combining algebraic hyperstructures, which are generalizations of classical algebraic structures, with rough set theory. In this study, definitions and some properties of hypergroups, polygroups, rough sets and rough groups are presented. Then, the connections between rough hypergroups and rough polygroups are examined as using the notion of lower and upper approximations, and some important characterizations are studied.

Keywords: Hypergroup; Polygroup; Rough set; Rough hypergroup; Rough polygroup.

REFERENCES

1. Pawlak, Z. Rough sets. Int J Inf Comp Sci. Vol:11, 341–356, 1982.
2. Davvaz, B. Rough polygroups, Italian J. Pure Appl. Math. Vol: 12, 91-96, 2002.
3. Davvaz, B. Rough subpolygroups in a factor polygroup, Journal of Intelligent and Fuzzy Systems, Vol:17, No:6, 613-621, 2006.
4. Marty, F. Sur ungeneralisation de la notion degroup, 8th Congress of Scandinavian Mathematicians, Stockholm, 45-49, 1934.
5. Corsini, P., Leoreanu, V. Applications of hyperstructures theory, Kluwer Academic Publishers, 2003.
6. Oguz, G. Rough Approximation Operators in a Topological Ring. Erzincan University Journal of Science and Technology, 13(1), 144-151, 2020.
7. Fotea, V.L. The lower and upper approximations in a hypergroup. Information Sciences, Vol:178, No:18, 3605-3615, 2008.

Analytic Solutions for Third-Order Fractional Partial Differential Equation Using Modified Double Laplace Transform Method

Shorish Omer Abdulla¹, Mahmut MODANLI²

¹ Department of Mathematics, Garmian University, Kalar, Iraq

² Department of Mathematics, Harran University, Şanlıurfa, Turkey

shorish.omer@garmian.edu.krd¹, mmodanli@harran.edu.tr²

Abstract

This paper finds analytical solutions to third-order partial differential equations with fractional derivatives, formulated by the Atangana-Baleanu Caputo (ABC) formulation. Under certain initial and boundary conditions, the study finds analytical expressions using the double Laplace transform method that satisfy the fractional partial differential equation.

Keywords: Third-order partial differential equation; Atangana-Baleanu Caputo (ABC) fractional derivative; Double Laplace decomposition method.

REFERENCES

1. Sadaq Taha Abdulazeez, Mahmut Modanli, Analytic solutions of fractional order Pseudo-Hyperbolic Telegraph equation using modified double Laplace transform method, International Journal of Mathematics and Computer in Engineering, Vol:1, No:1, 105–114, 2023.
2. Mohamed Z. Mohamed, Tarig M. Elzaki, Mohamed S. Algom, Eltaib M. AbdElmohmono, Amjad E. Hamza, New Modified Variational Iteration Laplace Transform Method Compares Laplace Adomian Decomposition Method for Solution Time-Partial Fractional Differential Equations, Journal of Applied Mathematics, Vol: , 1-10, 2021.
3. Lokenath Debnath, The double Laplace transforms and their properties with applications to functional, integral and partial differential equations, International Journal Applied Computational Mathematics, Vol:2, 223-241, 2016.

Some New Results for Exponential-type Durrmeyer Sampling Series

Sadettin Kursun¹

¹Department of Basic Sciences, National Defence University, Turkish Military Academy,
Ankara, Türkiye

sadettinkursun@yahoo.com,

Abstract

In the present paper, we deal with approximation by exponential sampling Durrmeyer operators in logarithmic weighted spaces. We will present pointwise and uniform convergence of the family of operators for functions belonging to logarithmic weighted space of functions and give a rate of convergence via suitable modulus of continuity.

Keywords: Exponential sampling series; Durrmeyer operators; Pointwise/Uniform convergence; Rate of convergence.

REFERENCES

1. Acar T., Kursun S., Pointwise convergence of generalized Kantorovich exponential sampling series, Dolomites Research Notes on Approximation, Vol:16, No:1, 1-10, 2023.
2. Aral A., Acar T., Kursun S., Generalized Kantorovich forms of exponential sampling series, Analysis and Mathematical Physics, Vol:12, No:1, 2022.
3. Bardaro C., Faina L., Mantellini I., A generalization of the exponential sampling series and its approximation properties, Mathematica Slovaca, Vol:67, No:6, 1481-1496, 2017.
4. Bardaro C., Mantellini I., On a Durrmeyer-type modification of the exponential sampling series, Rendiconti del Circolo Matematico di Palermo Series 2, Vol:70, 1289-1304, 2021.

New soliton solutions with generalized exponential rational function method

Ugur BAYRAKCI¹, Seyma TULUCE DEMIRAY¹, Huseyin YILDIRIM²

¹ Department of Mathematics, University of Osmaniye Korkut Ata, Osmaniye, Turkey

² Department of Mathematics, University of Kahramanmaraş Sütçü İmam, Kahramanmaraş, Turkey

ubayrakci42@gmail.com, seymatuluce@gmail.com, hyildir.ksu.edu.tr

Abstract

In this study, the fractional generalized perturbed KdV equation is studied. The generalized exponential rational function method is applied to this equation. Thus, various soliton solutions of this equation are obtained. Various values are given to the obtained solutions and graphical drawings are made.

Keywords: Generalized exponential rational function method; Fractional generalized perturbed KdV equation; Soliton solutions.

REFERENCES

1. Marwan Alquran, Mohammed Ali, Kamel Al-Khaled, George Grossman, Simulations of fractional time-derivative against proportional time-delay for solving and investigating the generalized perturbed-KdV equation, *Nonlinear Engineering*, 12(1), 20220282, 2023.
2. Khan Shehzada, Aman Ullah, Sayed Saifullah, Ali Akgül, Fractional generalized perturbed KdV equation with a power Law kernel: A computational study, *Results in Control and Optimization*, 12, 100298, 2023.
3. Hajar Farhan Ismael, Hasan Bulut, Hacı Mehmet Baskonus, W-shaped surfaces to the nematic liquid crystals with three nonlinearity laws, *Soft Computing*, 25, 4513-4524, 2021.
4. Şeyma Tülüce Demiray, Uğur Bayrakçı, A study on the solutions of (1+1)-dimensional Mikhailov-Novikov-Wang equation, *Mathematical Modelling and Numerical Simulation with Applications*, 3(2), 101-110, 2023.

INTRODUCTION TO \mathcal{M} -STURM-LIOUVILLE PROBLEM FOR DIFFUSION OPERATOR

Merve KARAOGLAN¹ Erdal BAS¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

karaoglanmerve24@gmail.com, erdalbas@gmail.com

Abstract

In this article, we present the \mathcal{M} -Sturm Liouville problem for the diffusion operator depending on initial condition. The aim of the paper is to prove a more general version of the representation of the solution of the Sturm Liouville problem for diffusion operator in classical analysis. We achieve the representation of the solution of the \mathcal{M} -Sturm Liouville problem for diffusion operator through the \mathcal{M} -Laplace transform[1-4].

Keywords: \mathcal{M} -derivative; \mathcal{M} -Laplace transform; Diffusion operator.

REFERENCES

1. Acay, B., Bas, E., Abdeljawad, T., Non-local fractional calculus from different viewpoint generated by truncated M-derivative, Journal of Computational and Applied Mathematics, 366, 112410, 2020.
2. Bas, E., Acay, B., The direct spectral problem via local derivative including truncated Mittag-Leffler function. Applied Mathematics and Computation, 367, 124787, 2020.
3. Bas, E., The Inverse Nodal problem for the fractional diffusion equation. Acta Scientiarum. Technology, 37(2), 251-257, 2015.
4. Koyunbakan, H., Panakhov, E. S., Half-inverse problem for diffusion operators on the finite interval, Journal of Mathematical Analysis and Applications, 326(2), 1024-1030, 2007.

PROPERTIES OF ROUGH SUBGRUPOIDS

M. Mustafa BEYDAĞI¹, A. Fatih ÖZCAN², İlhan İÇEN³

¹Department of Mathematics, Faculty of Arts and Sciences, Inonu University, Malatya, Turkey

mbeydagi23@gmail.com

²Department of Mathematics, Faculty of Arts and Sciences, Inonu University, Malatya, Turkey

abdullah.ozcan@inonu.edu.tr

³Department of Mathematics, Faculty of Arts and Sciences, Inonu University, Malatya, Turkey

ilhan.icen@inonu.edu.tr

Abstract

In this paper, the definition of rough subgroupoid is given using the concepts of groupoid and rough subgroup. By defining the lower and upper approximations of a given set according to the normal subgroupoid, the rough set of this set is obtained. If the lower and upper approximations in the rough set are subgroupoids of the groupoid, then this rough set is a rough subgroupoid with respect to the lower and upper approximations of the groupoid. The properties of the rough subgroups also apply to the rough subgroupoids.

Keywords: Rough Set, Groupoid, Rough Group, Rough Subgroup, Rough Subgroupoid.

REFERENCES

1. Pawlak, Z. (1982). Rough sets, Int. J. Comput. Inform. Sci., 11, 341-356.
2. Biswas, R., Nanda, S. (1994). Rough groups and rough subgroups, Bull. Polish. Acad. Sci. Math., 42, 251-254.
3. Kuroki, N., Wang, P.P. (1996). The lower and upper approximations in a fuzzy group, Inform. Sci., 90, 203-220.
4. Erdogan, F. (2006). ~ The Relationship between fuzzy rough sets and rings, M.Sc. Thesis, Gaziosmanpaşa University, Tokat, Turkey.
5. Taşbozan, H. (2017). Lokal Rough Kümeler ve Rough Altgrupoidler, Ph. D. Thesis, İnönü University, Malatya, Turkey.

Solving Dynamic Complexity with Analytical Solution Techniques

Ulviye Demirbilek¹, Mehmet Şenol², Hasan Bulut³

¹ Department of Mathematics, University of Mersin, Mersin, Turkey

² Department of Mathematics, University of Hacı Bektaş Veli, Nevşehir, Turkey

³ Department of Mathematics, University of Firat, Elazığ, Turkey

udemirbilek@mersin.edu.tr, msenol@nevsehir.edu.tr, hbulut@firat.edu.tr,

Abstract

In this study, we introduce innovative exact solutions for a dynamic model applicable in various fields such as fiber optics, ferromagnetics, water engineering, and oceanography. Our solutions comprise diverse mathematical forms, including exponential, trigonometric, and rational functions. Utilizing advanced symbolic computational software, we visualize the physical behavior of these solutions comprehensively, incorporating 2D and 3D graphs. The various solution techniques presented herein provide valuable insights into dynamic model solutions across different application domains. Moreover, numerous solution techniques documented in the literature significantly contribute to these diverse application areas [1-3].

Keywords: Analytical techniques, dynamic model, exact solution.

REFERENCES

1. Şenol, M., Gençyiğit, M., Demirbilek, U., Akinyemi, L., and Rezazadeh, H. New analytical wave structures of the $(3+1)$ -dimensional extended modified Ito equation of seventh-order. *Journal of Applied Mathematics and Computing*, 1-17, 2024.
2. Kirci, Ö., Koç, D. A., and Bulut, H. Dynamics of traveling wave solutions of conformable time-fractional ISLW and DJKM equations via a new expansion method, 2024.
3. Yue, J., Zhao, Z., and Wazwaz, A. M. Solitons, nonlinear wave transitions and characteristics of quasi-periodic waves for a $(3+1)$ -dimensional generalized Calogero-Bogoyavlenskii-Schiff equation in fluid mechanics and plasma physics. *Chinese Journal of Physics*, 2024.

GRUPOID ATLASES

Semih GEÇEN¹, İlhan İÇEN², A. Fatih ÖZCAN³

¹Department of Topology, Graduate School of Nature and Applied Sciences, Inonu University, Malatya, semihgecen44@gmail.com

²Department of Topology, Graduate School of Nature and Applied Sciences, Inonu University, Malatya, ilhan.icen@inonu.edu.tr

³Department of Topology, Graduate School of Nature and Applied Sciences, Inonu University, Malatya, abdullah.ozcan@inonu.edu.tr

Abstract

The concept of global effect defined by Anthony Bak is a more complex algebraic structure that reinterprets known classical group effects within certain rules. Many mathematicians have worked on this concept and made various contributions to the literature.

A. Bak developed a combinatorial approach to higher K-theory, in which control is kept of the elementary operations involved, through paths and ‘paths of paths’ in what he called a global action. The homotopy theory of these was developed by G. Minian. R. Brown and T. Porter developed applications to identities among relations for groups, and also the extension to groupoid atlases.

Here, we focused on the concept of groupoid atlas, which provides the necessary transitions while transferring the global effect to the concept of groupoid.

Keywords : Global action, K-theory, groupoid atlases

REFERENCES

[1] A. Bak, ‘Global actions: The algebraic counterpart of a topological space.’ *Uspeki Mat. Nauk.*, English Translation: *Russian Math. Surveys* 525 (1997) 955 –996.

[2] H. Abels and S. Holz, ‘Higher generation by subgroups.’ *J. Alg* 160 (1993) 311– 341.

[3] A. Bak, ‘Topological methods in algebra.’ ‘Rings, Hopf Algebras and Brauer Groups,’ (eds S. Caenepeel and A. Verschoren) (M. Dekker, New York, 1998), no. 197 in *Lect. Notes in Pure and Applied Math.*

[4] A. Bak, R. Brown, G. Minian and T. Porter, ‘Global actions, groupoid atlases and related topics.’ University of Wales, Bangor, Mathematics Preprint 99.27.

Modified Kudryashov Method for Solving Van der Waals Gas System

Arif ÖZKUL¹, Tolga AKTÜRK², Hasan BULUT³

¹Department of Mathematics, Faculty of Science, Firat University, Turkey

²Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey

³Department of Mathematics, Faculty of Science, Firat University, Turkey

¹arifozkul@gmail.com, ²tolgaakturkk@gmail.com, ³hbulut@firat.edu.tr

Abstract

In the paper, we obtain some the exact solutions to the nonlinear Van der Waals Gas system which refers to a model for the behavior of gases. This system were analyzed by utilizing the modified Kudryashov method.

Keywords: Modified Kudryashov method; Van der Waals gas system; Exact solution.

REFERENCES

1. H. Bulut, T.A. Sulaiman, F. Erdogan, H.M. Baskonus, On the new hyperbolic and trigonometric structures to the simplified MCH and SRLW equations, Eur. Phys. J. Plus. 132 (2017) 1–12.
2. K.K. Ali, R. Yilmazer, A. Yokus, H. Bulut, Analytical solutions for the $(3+1)$ -dimensional nonlinear extended quantum Zakharov–Kuznetsov equation in plasma physics, Phys. A Stat. Mech. Its Appl. 548 (2020) 124327.
3. G. Yel, H.M. Baskonus, H. Bulut, Novel archetypes of new coupled Konno–Oono equation by using sine–Gordon expansion method, Opt. Quantum Electron. 49 (2017) 285.

A STUDY ON NUMERICAL SOLUTION OF THE REGULARIZED LONG WAVE EQUATION

Sibel Özer¹, Yusuf Uçar¹, Damla Özçelik¹

¹ Department of Mathematics, Inonu University, Malatya, Turkey

sibel.ozer@inonu.edu.tr, yusuf.ucar@inonu.edu.tr, 36223614919@ogr.inonu.edu.tr

Abstract

This study involves the numerical solution of the 1-Dimensional Regularized Long Wave (RLW) equation, which has an important place in fluid dynamics. After the RLW equation is discretized according to temporal and spatial variables using the three-step Adams-Bashforth and finite difference methods, respectively, numerical solutions are found with the help of the obtained schemes. The proposed schemes are applied to the single solitary wave problem and error norms and conservation constants are calculated. The calculated error norms and conservation constants are presented in tables and compared with the studies of the previous researchers. Additionally, the problem is presented visually by plotting the graphs of the numerical results obtained. From the tables and graphs, it is seen that the proposed schemes are compatible with those of the analytical solution of the single solitary wave problem and preserved the conservation constants.

Keywords: Regularized Long Wave Equation; Adams-Bashforth Method; Finite Difference Method .

REFERENCES

1. S. Kutluay, A. Esen, A finite difference solution of the regularized long-wave equation, Mathematical Problems in Engineering, Vol:2006, 1-14, 2006.
2. J. Lin, Z. Xie, J. Zhou, High-order compact difference scheme for the regularized long wave equation, Communications in Numerical Methods in Engineering, Vol:23, No:2, 135-156, 2007.
3. N.M. Yagmurlu, Y. Ucar, I. Celikkaya, Operator splitting for numerical solutions of the RLW equation, Journal of Applied Analysis and Computation, Vol:8, No:5, 1494-1510, 2018.
4. J.S. Hu, J.J. Li, X. Wang, New High-Order Conservative Difference Scheme for Regularized Long Wave Equation with Richardson Extrapolation, Thermal Science, Vol. 23, No: 3, S737-S745, 2019.

*This work has been supported by İnönü University Scientific Research Projects Unit under Grant No. FYL-2024-3459.

DOMAIN OF MERSENNE MATRIX OPERATOR IN THE SPACE OF CONVERGENT SEQUENCES

Sezer Erdem¹ and Serkan Demiriz²

¹Department of Basic Engineering Sciences, Malatya Turgut Özal University, Malatya, Turkey

²Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Turkey

sezererdem8344@gmail.com,

serkandemiriz@gmail.com

Abstract

In this study, we obtain a new sequence space as the domain of regular Mersenne matrix operator in the space of convergent sequences and we examine some fundamental properties of the aforementioned space.

Keywords: Mersenne matrix operator; Sequence space; Duals; Matrix transformations.

REFERENCES

1. Feyzi Başar, Summability Theory and its Applications, 2nd ed., CRC Press/Taylor & Francis Group, Boca Raton·London·New York, 2022.
2. Muhammet Cihat Dağlı, Matrix mappings and compact operators for Schröder sequence spaces. Turkish Journal of Mathematics, Vol:46, 2304-2320, 2022.
3. Serkan Demiriz, Sezer Erdem, Mersenne Matrix Operator and Its Application in p-Summable Sequence Space, Communications in Advanced Mathematical Sciences, Vol:7, No:1, 42-55, 2024.
4. Emrah Evren Kara, Some topological and geometrical properties of new Banach sequence spaces, J. Inequal. Appl., Vol:2013, No:38, 2013.
5. Taja Yaying, Bipan Hazarika, On sequence spaces defined by the domain of a regular Tribonacci matrix, Math. Slovaca, Vol:70, No:3, 697-706, 2020.
6. Taja Yaying, Merve İlkhana Kara, On sequence spaces defined by the domain of tribonacci matrix in c_0 and c , The Korean Journal of Mathematics, Vol:29, No:1, 25-40, 2021.

A NOTE ON ALMOST CONVERGENT MERSENNE SEQUENCE SPACE

Serkan Demiriz¹ and Sezer Erdem²

¹ Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Turkey

² Department of Basic Engineering Sciences, Malatya Turgut Özal University, Malatya Turkey

serkandemiriz@gmail.com

sezererdem8344@gmail.com

Abstract

This study aims to present a new sequence space obtained as the domain of regular Mersenne matrix in the space of almost convergent sequences and to determine its duals and some properties.

Keywords: Mersenne matrix; Almost convergence; Sequence space; Duals.

REFERENCES

1. Feyzi Başar, Summability Theory and its Applications, 2nd ed., CRC Press/Taylor & Francis Group, Boca Raton·London·New York, 2022.
2. Muhammet Cihat Dağlı, A new almost convergent sequence space defined by Schröder matrix, Linear and Multilinear Algebra, Vol:71, No:11, 1863-1874, 2023.
3. Serkan Demiriz, Merve İlkhani, Emrah Evren Kara, Almost convergence and Euler totient matrix, Ann. Funct. Anal., No:11, 604–616, 2020.
4. Serkan Demiriz, Sezer Erdem, Mersenne Matrix Operator and Its Application in p-Summable Sequence Space, Communications in Advanced Mathematical Sciences, Vol:7, No:1, 42-55, 2024.
5. Emrah Evren Kara, Some topological and geometrical properties of new Banach sequence spaces, J. Inequal. Appl., Vol:2013, No:38, 2013.
6. Taja Yaying, Bipan Hazarika, On sequence spaces defined by the domain of a regular Tribonacci matrix, Math. Slovaca, Vol:70, No:3, 697-706, 2020.
7. Taja Yaying, Merve İlkhani Kara, On sequence spaces defined by the domain of tribonacci matrix in c_0 and c , The Korean Journal of Mathematics, Vol:29, No:1, 25-40, 2021.

GRAVITY MODELLING AND EARTHQUAKE ANALYSIS FOR EAST ANATOLIAN FAULT ZONE AND SURROUNDING AREA

Bülent Oruç¹, Mustafa Berkay Doğan², Emir Balkan³, İlkin Özsöz², Sunay Mutlu³, Aybala Büşra Çalışkur¹

¹Kocaeli University, Engineering Faculty, Department of Geophysical Engineering, 41001 İzmit-Kocaeli, TÜRKİYE

²General Directorate of Mineral Research and Exploration, Marine Research Department Çukurambar District, 06800, Çankaya, Ankara, TÜRKİYE

³Eskişehir Technical University, Institute of Earth and Space Sciences, 26555, Eskişehir, Türkiye

bulent.oruc@kocaeli.edu.tr

Abstract

The 6 February 2023 Kahramanmaraş Earthquakes with moment magnitudes of 7.7 and 7.6 were the disaster of the century for Türkiye. The sequence consisted of two mainshocks demonstrated that East Anatolian Fault Zone (EAFZ) is active not only destructive earthquake potential, but also shallow depth of focus at upper crustal domain. Thus the strength of the upper crustal domain is the important factor in terms of behavior of the upper-lower crust system with earthquake activities. In this study, the crustal deformation and structures of the EAFZ and surrounding area have been investigated using gravity and earthquake analysis. We first calculate radially averaged logarithmic amplitude spectrum of WGM12 isostatic residual gravity anomalies to estimate the average depths of the deeper interfaces and critical cut-off wavenumbers for filtering operations. The average depths of the Conrad and basement have been calculated as 18 km and 6.6 km for the linear segments of the spectrum, respectively. The Conrad undulation is constructed with the gravity inversion based on Parker-Oldenburg's algorithm. Then linear inversion technique, namely the weighted and damped minimum norm inverse solution was then used to estimate density contrasts down to Conrad depth. This inversion technique indicates that deeper interfaces are more undulated in the northern part of the EAFZ region. In addition, sharpen gradient in upper crustal thicknesses in the northern part of the EAFZ, where significant lateral density contrast variations has been revealed. We point out that increasing earthquake activities are caused by intraplate deformation and weaker strength of the crust considering undulated interfaces and sharpen density contrast variation.

Keywords: East Anatolian Fault Zone; Gravity inversion; Upper crust; Seismicity

GENERALIZED FRACTIONAL THE VERTICAL MOTION OF A FALLING BODY PROBLEM

Erdal BAS¹ Ali SELCUK¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

erdalmat@yahoo.com , ali_selcuk@hotmail.com

Abstract

In this study, the vertical motion of falling body problem is considered by the newly given M -derivative and generalized fractional derivative. Analytical solutions of the modeling problem are obtained and supported by various graphs including different values comparatively with M -derivative and generalized fractional derivative. The Laplace transform is used as the method of choice. [1-4].

Keywords: M -derivative; Generalized fractional derivative; Mathematical Model; Laplace transform.

REFERENCES

1. Ozarslan,R.Ercan,A.,Bas,E.(2019).Novel fractional models compatible with real world problems. *Fractal and Fractional*,3(2),15.
2. Acay, B.,Bas, E.,Abdeljawad, T. (2020). Non-local fractional calculus from different viewpoint generated by truncated M -derivative. *Journal of Computational and Applied Mathematics*, 366,112410.
3. Jarad,F.,Abdeljawad,T.(2020).Generalized fractional derivatives and Laplace transform. *Discrete and Continuous Dynamical Systems -S*,2020,13(3):709-722.
4. Murray R. Spiegel - Applied Differential Equations-Prentice-Hall, Inc: 62-67 (1967).

SOME NEW RESULTS OF THE NONLINEAR CONFORMABLE MODEL ARISING IN PLASMA PHYSICS

Md. Nur Alam¹, Onur Alp İlhan^{2, a}, Md. Shahid Hasan³, Uzzal Saha⁴

And F. Berna Benli^{2, b}

^{1,3,4}Department of Mathematics, Pabna University of Science and Technology,
Pabna, 6600, Bangladesh.

²Faculty of Education, Erciyes University, 38039 Melikgazi, Kayseri, Turkey

E-mail address: ¹nuralam.pstu23@gmail.com, ^{2, a}oilhan@erciyes.edu.tr,
³shahid.math43@gmail.com, ⁴uzzalsaha671@gmail.com and
^{2, b}akpinarb@erciyes.edu.tr

Abstract

The nonlinear conformable model that arises in plasma physics is the 3D conformable Zakharov-Kuznetsov equation (CZKE) with power law nonlinearity (PLNL). The current study applies a modification of the (G'/G) -expansion (MG'/GE) approach to this model and obtains certain closed-form precise wave solutions. By going backwards into the 3D CZKE with PLNL, the obtained results are confirmed, and they are noted as being particularly advantageous over a number of current methods. For the other nonlinear conformable models in physics, mathematics, and engineering, the aforementioned approach could also be used to obtain closed-form wave solutions (CFWSSs).

Keywords: Conformable derivative The variation of (G'/G) -expansion method the 3D CZKE with power law nonlinearity.

REFERENCES

- [1] H. Y. Martineza, J. F. G. Aguilarb, D. Baleanu, Beta-derivative and sub-equation method applied to the optical solitons in medium with parabolic law nonlinearity and higher order dispersion, Optik 155 (2018) 357–365.
- [2] L. Alzaleq, D. Al-zaleq, S. Alkhushayni, Traveling Waves for the Generalized Sinh-Gordon Equation with Variable Coefficients, Mathematics 10 (2022), 822.
- [3] H. M. Baskonus, T. A. Sulaiman, H. Bulut, T. Aktürk, Investigations of dark, bright, combined dark-bright optical and other soliton solutions in the complex cubic nonlinear schrödinger equation with δ -potential, Superlattices and Microstructures 115 (2018) 19–29.

New Exact Solutions of the (1+1) dimensional nonlinear Ostrovsky equation

Sebahattin Ertas¹, Hasan Bulut¹, Yusuf Pandir²,

¹Department of Mathematics, University of Firat, Elazig, Turkey

²Department of Mathematics, Yozgat Bozok University, Yozgat, Turkey

hbulut@firat.edu.tr, yusuf.pandir@bozok.edu.tr

Abstract

In this study, we obtain new exact solutions to the (1+1) dimensional nonlinear Ostrovsky equation which models for weakly nonlinear surface and internal waves in a rotating ocean by using new version trial equation method. This method allows new exact solutions of the nonlinear partial differential equations. We can say that these new exact solutions which are not in the literature. In addition, two and three dimensional graphs were drawn to illustrate the physical behavior of these new exact solutions.

Keywords: New version trial equation method; (1+1) nonlinear Ostrovsky equation; Exact solutions.

REFERENCES

1. Liu Cheng Shi, Using trial equation method to solve the exact solutions for two kinds of KdV equations with variable coefficients. Acta Physica Sinica, Vol:54, No:10, 4506-4510, 2005.
2. Pandir Yusuf, Gurefe Yusuf, Misirli, Emine, Classification of exact solutions to the generalized Kadomtsev-Petviashvili equation. Physica Scripta, Vol:87, No:2, 025003, 2013.
3. Pandir Yusuf, Gurefe Yusuf, Misirli Emine, A multiple extended trial equation method for the fractional Sharma-Tasso-Olver equation, AIP Conference Proceedings, Vol:1558, No:1, 1927-1930, 2013.
4. Pandir Yusuf, Ekin Ali, Dynamics of combined soliton solutions of unstable nonlinear Schrodinger equation with new version of the trial equation method, Chinese Journal of Physics, Vol:67, 534-543, 2020.

GENERALIZED SYSTEMS OF LINEAR EQUATIONS WITH LOCAL DERIVATIVE

Enise KARTAL¹, Erdal BAS¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

kartalenise2323@gmail.com, erdalbas@gmail.com

Abstract

In this paper, we consider systems of linear differential equations with local derivative. The aim of this paper is to solve systems of linear equations with \mathcal{M} -derivative in a more general version. We solve \mathcal{M} -derivative systems of linear equations through the \mathcal{M} -Laplace transform. In addition, the Cramer method is used by adapting the local derivative and the necessary solutions are given by Mittag-Leffler functions [1-4].

Keywords: Linear equation systems; \mathcal{M} -derivative; \mathcal{M} -Laplace transform.

REFERENCES

1. Acay, B., Bas, E., Abdeljawad, T., Non-local fractional calculus from different viewpoint generated by truncated M-derivative, Journal of Computational and Applied Mathematics, 366, 112410, 2020.
2. Bas, E., Acay, B., The direct spectral problem via local derivative including truncated Mittag-Leffler function, Applied Mathematics and Computation, 367, 124787, 2020.
3. Sene, N., Mittag-Leffler input stability of fractional differential equations and its applications, Discrete and Continuous Dynamical Systems-S, 13(3), 867-880, 2020.
4. Sousa, J. Vanterler da C., Capelas de Oliveira, E., A new truncated M-fractional derivative type unifying some fractional derivative types with classical properties, International Journal of Analysis and Applications, 16.1, 83-96, 2018.

Model Order Reduction for Shigesada-Kawasaki-Teramoto Cross-Diffusion Systems

Glden Mlayim¹

¹Department of Mathematics, Adıyaman University, Adıyaman, Turkey

gmulayim@adiyaman.edu.tr,

Abstract

Shigesada-Kawasaki-Teramoto (SKT) is one of the most popular nonlinear cross-diffusion systems in population ecology. Full-order model (FOM) solutions of the SKT system are computed by applying the symmetric interior penalty discontinuous Galerkin method (SIPG) for space discretization and the semi-implicit Euler method for time discretization. Reduced solutions are calculated by using the proper orthogonal decomposition (POD) method. Also, the discrete empirical interpolation method (DEIM) is used in the computation of nonlinearities of the SKT system. Full-order model solutions and reduced order models (ROM) solutions are compared in numerical results which show the accuracy and efficiency of POD and POD-DEIM methods for the SKT system.

Keywords: SKT systems; full order model solutions; reduced order model solutions; proper orthogonal decomposition method (POD); discrete empirical interpolation method (DEIM).

REFERENCES

1. N. Shigesada, K. Kawasaki, E. Teramoto, Spatial segregation of interacting species, J. Theor. Biol., Vol:79, No:1, 83-99, 1979.
2. K. Kunisch, S. Volkwein, Galerkin proper orthogonal decomposition methods for parabolic problems, Numer. Math., Vol:90, No:1, 117-148, 2001.
3. S. Chaturantabut, D. C. Sorenson, A state estimation for POD-DEIM nonlinear model reduction, SIAM J. Numer. Anal., Vol:50, No:1, 46-43, 2012.

AN INVERSE NODAL PROBLEM OF A CONFORMABLE STURM-LIOUVILLE PROBLEM BY RETARDED CONSTANT

Auwalu Sa'idu¹, Hikmet Kemaloğlu (Koyunbakan)²,

Department of Mathematics, Firat University, Elazig, Turkey

¹asaaidu@yumsuk.edu.ng, ²hkoyunbakan@gmail.com

Abstract

In this paper, we presented a new technique to the literature: a conformable derivative for the inverse problem of a Sturm-Liouville problem with constant delay. We treated the characteristic equation in four different cases and calculated the asymptotic formulas for the eigenvalues and eigenfunctions for each case and demonstrated the existence of the solution. Additionally, we identified the nodal points, from which we generated the problem's potential functions. Next, we used the Lipschitz stability approach and demonstrated the problem's stability.

Keywords: Local derivative, Conformable Sturm-Liouville problem, Spectrum, Constant delay

References

1. Abdeljawad T. On conformable fractional calculus. *Journal of computational and Applied Mathematics* 2015; 279:57-66.
2. Adalar I, Ozkan AS. Inverse problems for a Fractional Sturm-Liouville operators. *Journal of Inverse and Ill-posed Problem* 2020; 28 (6): 777-782.
3. Allahverdiev BP, Tuna H, Yalcinkaya Y. Conformable Fractional Sturm Liouville equation. *Mathematical Methods in the Applied Sciences* 2019; 42 (10):3508-3526.
4. Freiling G, Yurko, VA. Inverse problems for Sturm Liouville differential operators with a constant delay. *Applied Mathematics Letter* 2012; 25 (11), 1999- 2004.
5. Mortazaasl H, Jodayree Akbarfam A. Trace formula and inverse nodal problem for a conformable fractional Sturm-Liouville problem. *Inverse Problems in Science and Engineering* 2020; 28 (4):524-55. <https://doi.org/10.1080/17415977.2019.1615909>.
6. Sa'idu A, Koyunbakan H. Inverse fractional Sturm-Liouville problem with eigenparameter in the boundary conditions. *Mathematical Methods in the Applied Sciences* 2022; 1-10. <https://doi.org/10.1002/mma.8433>
7. Sa'idu A, Koyunbakan H. Transmutation of conformable Sturm-Liouville operator with exactly solvable potential. *Filomat*. 2023;37(11):3383-90.
8. Saidu A, Koyunbakan H. A conformable inverse problem with constant delay. *J. Adv. App. Comput. Math.* 2023; 10: 26-38. DOI: <https://doi.org/10.15377/2409-5761.2023.10.3>

FUNDAMENTAL ALGEBRAIC AND TOPOLOGICAL CONCEPTS IN GEOMETRIC ANALYSIS

Emre Civgin¹, Numan Yalcin^{1*}

¹ Department of Electronics and Automation, Vocational School of Gumushane, University of Gumushane, Gumushane, Turkey

numan@gumushane.edu.tr,

emrec57@hotmail.com

Abstract

In this study, the fundamental definitions and theorems given for some algebraic and topological concepts known from classical analysis are given in geometric analysis.

Keywords: Geometric Calculus ; Topological concept; Algebraic concept.

REFERENCES

- [1] Grossman, Michael, and Robert Katz. Non-Newtonian Calculus: A Self-contained, Elementary Exposition of the Authors' Investigations... Non-Newtonian Calculus, 1972.
- [2] Çakmak, Ahmet Faruk, and Feyzi Başar. "Some new results on sequence spaces with respect to non-Newtonian calculus." Journal of Inequalities and Applications 2012 (2012).
- [3] Çakmak, Ahmet Faruk, and Feyzi Başar. "Certain spaces of functions over the field of non-Newtonian complex numbers." Abstract and Applied Analysis. Vol. 2014. Hindawi, 2014.
- [4] Çakmak, Ahmet Faruk, and Feyzi Basar. "Some sequence spaces and matrix transformations in multiplicative sense." TWMS Journal of Pure and Applied Mathematics 6.1 (2015): 27-37.
- [5] Tekin, Sebiha, and Feyzi Başar. "Certain sequence spaces over the non-Newtonian complex field." Abstract and Applied Analysis. Vol. 2013. Hindawi, 2013.
- [6] Türkmen, Cengiz, and Feyzi Başar. "Some basic results on the sets of sequences with geometric calculus." AIP conference proceedings. Vol. 1470. No. 1., 2012.
- [7] Türkmen, C., and F. Basar. "Some basic results on the geometric calculus, Commun." Fac. Sci. Univ. Ank. Series A 1: 61.

Blow up at finite time for sixth-order evolution equations with time dependent coefficient

Ayşe Fidan¹, Erhan Pişkin²

¹Dicle University, Institute of Natural and Applied Sciences, Department of Mathematics,
Diyarbakır, Turkey

afidanmat@gmail.com,

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

episkin@dicle.edu.tr

Abstract

In this presentation, we consider sixth-order evolution equations with time-dependent coefficients. We establish both lower and upper bounds for the blow-up time using a differential inequality argument to determine when blow-up occurs.

Keywords: Blow-up, Parabolic-type equations, Variable Coefficients.

REFERENCES

1. H. Di, Y. Shang, Blow-up phenomena for a class of metaparabolic equations with time dependent coefficient, AIMS Ser. Appl. Math. 2(4) (2017), 647-657.
2. E. Pişkin, B. Okutmuşur, An Introduction to Sobolev Spaces, Bentham Science, 2021.
3. E. Pişkin, A. Fidan, Finite time blow up of solutions for the m-Laplacian equation with variable coefficients, Al-Qadisiyah Journal of Pure Science, 28(1)(2023).
4. E. Pişkin, A. Fidan, Nonexistence of global solutions for the strongly damped wave equation with variable coefficients, Universal Journal of Mathematics and Applications, 5 (2) (2022), 51-56.

A COMPARATIVE STUDY OF FINITE ELEMENT METHODS WITH CUBIC AND QUINTIC BASIS FUNCTIONS FOR THE SMCH EQUATION

Hatice KARABENLİ¹, Yusuf UÇAR², E. Nesligül AKSAN², Alaattin ESEN²

¹Nuray Tuncay Kara Science and Art Center, Ministry of Education, Gaziantep, Turkey

²Department of Mathematics, University of Inonu, Malatya, Turkey

haticekarabenli@gmail.com, yusuf.ucar@inonu.edu.tr, naksan@inonu.edu.tr,
alaattin.esen@inonu.edu.tr

Abstract

In this research, the numerical approaches for solving simplified modified Camassa-Holm(SMCH) equation are going to be examined. For this purpose, the studied model problem have been transformed into a linear form using Rubin-Graves linearization method. Then, the finite element method is applied to this linear equation using cubic and quintic B-Spline basis functions. The newly found numerical results are compared according to the degrees of the basis functions at different final time values. As per the obtained results, the error norms L_2 and L_∞ of the numerical results attained using quintic basis functions are smaller than the error norms of the numerical results attained using cubic basis functions. The finite element study based on different basis functions indicate that as the degree of the basis functions increases, the numerical solutions improve in accuracy and convergence. Moreover, higher accuracy is observed with not only larger of elements but also smaller time steps.

Keywords: Finite Element Method, Cubic B-Spline, Quintic B-Spline, Simplified Modified Camassa-Holm Equation.

REFERENCES

1. A.M. Wazwaz, A class of nonlinear fourth order variant of a generalized Camassa-Holm equation with compact and noncompact solutions, Applied Mathematics and Computation, Vol:165, No:2, 485-501, 2005.
2. P. M. Prenter, Splines and variational methods, Wiles, New York, 1975.
3. R. Camassa and D. Holm, An integrable shallow water equation with peaked solitons, Phys. Rev. Lett., Vol:71, No:11, 1661-1664, 1993.
4. S.G. Rubin and R.A. Graves, A Cubic Spline Approximation for Problems in Fluid Mechanics, NASA, Washington, District of Columbia, October, 1975.
5. S. Kutluay, Y. Ucar, and N. M. Yagmurlu, Numerical Solutions of the Modified Burgers Equation by a Cubic B-spline Collocation Method. Bull. Malays. Math. Sci. Soc. Vol:39, No:4, 39, 1603-1614, 2016.

A NEW GENERALIZED METHOD FOR THE FRACTIONAL NONLINEAR EQUATION

Dilara ALTAN KOÇ¹ Yusuf PANDIR² Hasan BULUT³

¹Department Mathematics, Faculty of Science Mugla Sıtkı Kocman University, Turkey,

²Department Mathematics, Faculty of Science, Bozok University, Yozgat, Turkey,

³Department of Mathematics, Faculty of Science, Firat University, Elazığ, Turkey,

dilaraaltan@mu.edu.tr, yusuf.pandir@bozok.edu.tr, hbulut@firat.edu.tr,

Abstract

In this paper, we propose a new generalized method to obtain some new exact trigonometric, hyperbolic, and rational function solutions of the special nonlinear equation. First, we introduce the new version of the generalized method, and then we give the exact solutions of the given nonlinear fractional differential equation. The obtained results are examined under different conditions. The three-dimensional figures for the reported solutions are drawn.

Keywords: Exact solution; Traveling wave solution; Fractional calculus.

REFERENCES

1. Anıqa Anıqa, Jamshad Ahmad, Soliton Solution of Fractional Sharma-Tasso-Olver Equation via an Efficient $(\frac{G'}{G})$ -expansion method. Ain Shams Engineering Journal, 13, 101528, 1-23, 2022.
2. Abdon Atangana, Dumitru Baleanu, New Fractional Derivatives with Nonlocal and Non-Singular Kernel: Theory and Application to Heat Transfer Model. Thermal Science, 20(2), 763-769, 2016.
3. Sagar B, Saha Ray S., An Efficient Meshfree Numerical Technique to Solve Fractional Schamel-KdV Equation for Ion-Acoustic Solitary Waves in Dusty Plasma, Ieee Transactions on Plasma Science, vol. 51, no. 7, July 2023.
4. Hasan Bulut, Hajar F. Ismael, Exploring new features for the perturbed Chen-Lee-Liu model via $(m + \frac{1}{G'})$ -expansion method. Proc. Inst. Math. Mech. Natl. Acad. Sci. Azerb. 48(1), 164-173, 2022.

SOME PROPERTIES OF THE EIGENFREQUENCIES OF THE PLATE ON THE DOMAIN

Natavan Allahverdiyeva¹, Yusif Gasimov²

¹Sumgayit State University, Sumgayit Azerbaijan

²Azerbaijan University, Baku, Azerbaijan

natavan.sdu@gmail.com

yusif.gasimov@au.edu.az

It is known that the plates are elements included in the composition of various mechanical constructions, and these constructions are widely used in industry, technology, and the construction sector covering the most diverse fields [2,]. Physical properties of different types of plates (free, fixed and clamped) are usually investigated in works dedicated to the study of these issues. But in many cases, these physical characteristics depend not only the material of which the board is made, the environment, etc. but also on its geometric properties. In this sense, the shape and the geometrical structure of the area of the plate are of great importance. Since it is possible to control the geometric parameters of the board in many cases, optimization of one or another physical characteristics required in practice can be achieved through these parameters. Such characteristics can be taken as mass, volume, critical power or eigenfrequency [1, 4, 6].

Here we consider the convex dependence of the eigenfrequency of the clamped plate on its shape.

Let the domain of the plate be D with boundary $S_D \in C^2$. It is known that [6,] the function $w(x_1, x_2, t)$ characterizing the across vibrations of the clamped plate satisfies the following equation

$$\omega_{x_1 x_1 x_1 x_1} + 2\omega_{x_1 x_1 x_2 x_2} + \omega_{x_2 x_2 x_2 x_2} + \omega_{tt} = 0. \quad (1)$$

As we consider the clamped plate the boundary condition should be given as below

$$u = 0, \quad \frac{\partial u}{\partial n} = 0, \quad x \in S_D. \quad (3)$$

Basing on the former results obtained in this direction [3, 5] the following theorems are proved.

Theorem 1. Let the support function of the area of the clamped plate satisfies the condition $P'_{D(t)}(x) > 0$ (or $P'_{D(t)}(x) < 0$) for each $t \in [t_0, t_1]$, where t is some parameter. Then the functional $\lambda_j(t)$ is quazi-convex.

Theorem 2. Let the eigenfrequency $\lambda(t)$ of the clamped plate is twice differentiable with respect to the parameter t and the derivative of the support function of the area of the plate keeps its sign in the interval $[t_0, t_1]$. Then there exists a number $k_0 \in R$ such that for any $k \geq k_0$ the functional $\Lambda(t) = e^{k\lambda(t)}$ is convex.

REFERENCES

1. Banichuk N.B. (1986). Introduction to Optimization of the Constructions. Moscow, Nauka.
2. Bucur D., Buttazzo G. (2005). Variational Methods in Shape Optimization Problems. Series: Progress in Nonlinear Differential Equations and Their Applications, 65, 2005, VIII, 216 p.
3. Gasimov Y.S. (2003). On some properties of eigenvalues when the domain varies. *Mathematical Physics, Analysis, Geometry*, 10(2), 249-255.
4. Gasimov, Y.S. (2008). Some shape optimization problems for eigenvalues. *Journal of Physics A: Mathematical and Theoretical*, 41(5), 055202.
5. Gasimov, Y.S. (2013). On a shape design problem for one spectral functional. *Journal of Inverse and Ill-Posed Problems*, 21(5), 629-637.
6. Gould S. (1970). Variational Methods in Eigenvalue Problems. Moscow, Mir.

EXPLOSIVE SOLUTIONS FOR A FOURTH-ORDER REACTION-DIFFUSION EQUATION IN VARIABLE EXPONENT SOBOLEV SPACES

Gülistan Butakın¹

Dicle University Institute of Natural and Applied Sciences Diyarbakır, Turkey

gulistanbutakin@gmail.com

Erhan Pişkin²

Dicle University Department of Mathematics Diyarbakır, Turkey

episkin@dicle.edu.tr

Abstract

In this work focuses on a fourth-order reaction-diffusion equation featuring variable exponents. Firstly, we explore the phenomenon of finite-time blow-up in solutions with positive initial energy. Later, we establish an upper limit on the blow-up time utilizing a technique involving differential inequalities.

Keywords: Explosive Solutions; Reaction-diffusion equation; Variable exponent.

REFERENCES

1. R. Abita, Blow-up phenomenon for a semilinear pseudo-parabolic equation involving variable source, *Applied Analysis*, 102(1) (2023), 88-103.
2. G. Butakın, E. Pişkin, Existence and Blow up of Solutions of a Viscoelastic $m(x)$ -Biharmonic Equation with Logarithmic Source Term, *Miskolc Mathematical Notes*, (in press).
3. G. Butakın, E. Pişkin, Existence and Blow up of Solutions for $m(x)$ -Biharmonic equation with Variable Exponent Sources, *Filomat*, (in press).

Decay of Solutions for a Nonlinear Hyperbolic-type Equations with Variable Exponents

Nebi Yılmaz¹, Erhan Pişkin²

¹Dicle University, Institute of Natural and Applied Sciences, Department of Mathematics, Diyarbakır, Turkey

nebiyilmaz1981@gmail.com,

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

episkin@dicle.edu.tr

Abstract

In this study, we investigate the decay of solutions to a nonlinear hyperbolic-type equation with variable exponents, focusing on the case without a source term. By employing Komornik's lemma, we establish the decay of the solutions under appropriate conditions on the variable exponents.

Keywords: Decay, hyperbolic-type equation, variable exponents.

REFERENCES

1. E. Pişkin, B. Okutmuşur, An Introduction to Sobolev Spaces, Bentham Science, 2021.
2. J. Ferreira, W. S. Panni, S. A. Messaoudi, E. Pişkin, M. Shahrouzi, Existence and Asymptotic Behavior of Beam-Equation Solutions with Strong Damping and $p(x)$ -Biharmonic Operator. Journal of Mathematical Physics, Analysis, Geometry, 18(4) (2022)488-513.
3. V. Komornik, Exact Controllability and Stabilization. The Multiplier Method, Masson-John Wiley, Paris, 1994.
4. N. Yılmaz, E. Pişkin, E. Çelik, Well-Posedness and Blow-up of Solutions for a Variable Exponent Nonlinear Petrovsky Equation, Advances in Mathematical Physics, Volume 2023, Article ID 8866861, 1-13.

FORCED VIBRATION ANALYSIS OF FUNCTIONALLY GRADED RODS BY PSEUDOSPECTRAL CHEBYSHEV METHOD

Özlem Cerit¹, Durmuş Yarımpaç¹

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

ozlemcerit08@gmail.com, durmusyarpabuc@osmaniye.edu.tr

Abstract

Forced vibration analysis of functionally graded rod with variable cross-sectional area in axial direction is considered. It is assumed that material properties such as elastic modulus and density are graded in the axial direction by the Voigt homogenization model. These conditions result in a partial differential equation with variable coefficients that is difficult to solve with conventional analytical methods. Under the Laplace transform, the partial differential equation is transformed into a time-independent boundary value problem in the axial direction and solved by the pseudospectral Chebyshev Method. With the modified Durbin method, displacements in physical space are obtained by taking the inverse transformation to the time domain. The results obtained are compared with the literature. The effects of randomly selected material mixture on displacement distributions are discussed.

Keywords: Forced vibration; Functionally graded materials; pseudospectral Chebyshev Method; Laplace transform, modified Durbin Method.

REFERENCES

1. Li, Q. S., Exact solutions for free longitudinal vibration of non-uniform rods. J. Sound Vib. 234(1):1–19, 2000.
2. Yardimoglu, B., Aydin, L., Exact longitudinal vibration characteristics of rods with variable crosssections. Shock and Vibration 18: 555 -562, 2011
3. Celebi, K., Keles, I., Tutuncu, N., Exact solutions for forced vibration of non-uniform rods by Laplace transformation. Gazi University Journal of Science 24(2): 347- 353, 2011
4. Akgoz, B., Civalek, Ö., Longitudinal vibration analysis of strain gradient bars made of functionally graded materials (FGM). Comp.: Part B: Eng. 55: 263- 268, 2013
5. Celebi, K., Yarımpaç, D., Baran, T., Forced vibration analysis of inhomogeneous rods with non-uniform cross-section, Journal of Engineering Research, 6(3), 189-202, 2018.

COMBINATORIAL INVARIANTS OF SATURATED NUMERICAL SEMIGROUPS

Meral Süer¹

¹Department of Mathematics, Batman University, Batman, Turkey

meral.suer@batman.edu.tr,

Abstract

In this study, our attention will be directed towards characterizing some families of saturated numerical semigroups. We will discuss about concepts factorizations of an element and the distances between factorizations of elements in terms of numerical semigroups. With the help of these concepts, we will also present results regarding the catenary degree of these numerical semigroup families, which is a combinatory concept and measures the spread of the distance between the factorizations of a given element.

Keywords: Saturated numerical semigroups; Catenary degree; Factorization.

REFERENCES

1. PanWang, BoTian, Kun Sun, Feng-Hua Qi, Thetitle of Paper, Applied Mathematics and Computation, Vol:2, No:1, 233–242, 2015.
2. F. Aguilo-Gost and P.A. Garc'ia-S'anchez, Factorization and catenary degree in 3-generated numerical semigroups, Electronic Notes in Discrete Mathematics, Vol:34,157–161, 2009.
3. A. Assi, P.A. Garc'ia-S'anchez, Numerical semigroups and applications, Springer Cham, RSME Springer Series, Switzerland, 2016.
4. V. Barucci, D.E. Dobbs, and M.Fontana, Maximality properties in numerical semigroups and applications to one-dimensional analytically irreducible local domains, Memoirs of the American Mathematical Society, Vol:598, 1–77, 1997.
5. S.T. Chapman, P.A. Garc'ia-S'anchez, and D. Llena, The catenary and tame degree of numerical monoids, Forum Mathematicum, Vol:21, No: 1, 117-129, 2009.
6. S.T. Chapman, P.A. Garc'ia-S'anchez, D. Llena, V. Ponomarenko, and J.C. Rosales, The catenary and tame degree in finitely generated commutative cancellative monoids, Nova Science Publishers, New York, 1999.
7. J.C. Rosales, P.A. Garc'ia-S'anchez, J.I. Garc'ia-Garc'ia, and M.B. Branco, Saturated numerical semigroups, The Houston Journal of Mathematics, Vol: 30, 321–330, 2004.

THERMAL ANALYSIS OF FUNCTIONALLY GRADED 2D PLATE

Oğuzhan Demirel¹, Durmuş Yarımpaç¹

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

demirelemu@hotmail.com
durmusyarpabuc@osmaniye.edu.tr,

Abstract

A closed form solution of the heat transfer problem of a two-dimensional heterogeneous plate in steady state condition has been obtained under the most general boundary conditions. It is assumed that the thermal conductivity of the material varies exponentially in two directions. An analytical solution to the partial differential equation obtained under these conditions is obtained by using the method of separation of variables. The results are discussed on graphs for specifically selected material properties.

Keywords: Heterogeneous plate; Functionally graded materials; Separation of variables

REFERENCES

1. Adineh, M. and Kadkhodayan, M., Three-Dimensional Thermo-Elastic Analysis and Dynamic Response of a Multi-Directional Functionally Graded Skew Plate on Elastic Foundation, Composites Part B: Engineering, vol. 125, pp. 227–240, 2017.
2. Alibeigloo, A., Exact Solution for Thermo-Elastic Response of Functionally Graded Rectangular Plates, Composite structures, vol. 92, no. 1, pp. 113–121, 2010.
3. Apalak, M. and Demirbas, M., Thermal Stress Analysis of In-Plane Two-Directional Functionally Graded Plates Subjected to In-Plane Edge Heat Fluxes, Proceedings of the Institution of Mechanical Engineers, Part L: Journal of Materials: Design and Applications, vol. 232, no. 8, pp. 693–716, 2018.
4. Yarımpaç D., Cihan, E., Kerimcan, C., and Eker, M., Heat Conduction Analysis of Two-Dimensional Anisotropic Plate, Çukurova University Journal of the Faculty of Engineering and Architecture, vol. 25, no. 1, pp. 139–147, 2020.

EXISTENCE AND UNIQUENESS RESULTS FOR SINGULAR FRACTIONAL DIFFERENTIAL EQUATIONS WITH P-LAPLACIAN OPERATOR

Nuket Aykut Hamal¹, Furkan Erkan¹

¹Department of Mathematics, University of Ege, Izmir, Turkey

nuket.aykut@ege.edu.tr , 91210000387@ogrenci.ege.edu.tr

Abstract

In this study, we investigate the existence and uniqueness of solutions for a singular fractional boundary value problem involving the p-Laplacian operator. Our analysis is based on a fixed point theorem. Examples are given for illustrating our main results.

Keywords: Fractional derivative, the p-Laplacian operator, existence and uniqueness, singular.

Acknowledgements: This study was supported by the Scientific and Technological Research Council of Turkey (TUBITAK) under the Grant Number 123F242. The authors thank to TUBITAK for their supports.

REFERENCES

1. Ahmed Alsaedi, Madeaha Alghanmi, Bashir Ahmad Boshra Alharbi, Uniqueness of solutions for a Ψ -Hilfer fractional integral boundary value problem with the p-Laplacian operator, Demonstratio Mathematica, Vol:56, No:1, 233–242, 2023.
2. Hanan A Wahash, Mohammed S Abdo, Abdulkafi M Saeed, Satish K Panchal, Singular fractional differential equations with Ψ -Caputo operator and modified Picard's iterative method, Appl. Math. E-Notes, Vol:20, 215-229, 2020.

THE SPECIFIC ENERGY AND SPECIFIC ANGULAR MOMENTUM ON ROTATIONAL SURFACES IN PSEUDO EUCLIDEAN 4-SPACE WITH INDEX 2

Fatma Almaz¹

¹ Department of Mathematics, University of Batman, Batman, TÜRKİYE

fatma.almaz@batman.edu.tr

Abstract

In taking into consideration the mathematical problem of the geodesics on a surface, there is an enormous advantage in conceptual comprehending that results from taking the point of view of a physicist by explaining parametrized geodesics as the paths traced out in time by the motion of a point on the surface, this combination of the constants of the motion is of course also constant along a geodesic. The existence of this constant is a conclusion of the one-parameter rotational group of symmetries of the rotational surfaces, like this constant of the movement introduces a new thing when the surface is invariant under any one-parameter group of symmetries, which is seen in the variation approximate to the geodesic equations easily. Mathematically, this quantity is a constant obtained by Clairaut for geodesic movement on surface defined in a coordinate system adapted to this one-parameter group of symmetries. In this paper, the results show that the specific energy and the specific angular momentum on the surfaces of rotation can be expressed in E_2^4 using some certain results describing the geodesics on the rotational surface are given. As a first instance, using the conditions of being geodesic, in which the curves can be chosen to be time-like curves, which allows us to constitute the specific energy and specific angular momentum.

Keywords: The specific energy; specific angular momentum; geodesic curves

REFERENCES

1. D. Lerner, Lie derivatives, izometries, and Killing Vectors, Depar. Math. Univ. of Cansas, Lawrence, Kansas, 66043-7594. 12, 2010.
2. F. Almaz, M.A. Kulahcı, The notes on rotational surfaces in Galilean space, International Journal of Geometric Methods in Modern Physics, 18(2), 2021.
3. F. Almaz, M.A. Kulahcı, The research on rotational surfaces in pseudo Euclidean 4-space with index 2, Acta Mathematica Universitatis Comenianae 93(3), 263-297, 2023.
4. I.M.Yaglom, A simple non-Euclidean geometry and its physical basis, Springer, New York, 1979
5. J.D. Walecka, Topics in Modern Physics: Theoretical Foundations, World Scientific, 1979.
6. J.D. Walecka, Introduction to General Relativity, World Scientific, Singapore, 2007.

Expansion Theorem for Sturm-Liouville Problem including Local Derivative

Mahmut Ozusan , Hikmet Kemaloğlu

Department of Mathematics, Firat University, Elazig, Turkey
ozusanmahmut@gmail.com, hkoyunbakan@gmail.com

Abstract

In this study, we give an expansion theorem for the Sturm-Liouville problem with M -derivative. For the classical Sturm-Liouville problem it was given by Levitan and Sargjan in [3] We will give this result for the following problem;

$$-D_{\alpha,M}^x(D_{\alpha,M}^x)y + q(x)y = \lambda y \quad (1.1)$$

$$D_{\alpha,M}^x y(0) - hy(0) = 0 \quad (2.2)$$

$$D_{\alpha,M}^x y(0)y(0) + HD_{\alpha,M}^x y(0) = 0 \quad (1.3)$$

where h and H are constants and $D_{\alpha,M}^x$ define the M -derivative for $0 < \alpha \leq 1$, $q(x)$ is real integrable and refers to potential. [1,2,4]

Keywords: Sturm-liouville problem, Potential functon, Spectrum, Conformable M-derivative

References

1. Abdeljawad, T. On Conformable Fractional Calculus, *Journal of computational and Applied Mathematics*, 2015, vol. 279, p. 57-66.
2. Rivero, M., Trujillo, J., & Velasco, M. A fractional approach to the Sturm-Liouville problem. *Open Physics*, 2013, 11(10), 1246-1254.
3. Levitan, B.M. and Sargsjan, I.S. (1975) Introduction to Spectral Theory: Self Adjoint Ordinary Differential Operators. American Mathematical Society, Providence. Mortazaasl, H., & Jodayree Akbarfam, A. Trace formula and inverse nodal problem for a conformable fractional Sturm-Liouville problem. *Inverse Problems in Science and Engineering*, 2020, 28(4), 524-555.

FOCAL CURVES ACCORDING TO THE ALTERNATIVE FRAME

Gülşah Belhan¹, Vedat ASİL¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

gulsahbelhan23@gmail.com, vasil@firat.edu.tr

Abstract

Curves in Euclidean space have been examined in many studies according to different frame works. While examining curves, researchers studied frameworks such as the Frenet frame, Bishop frame and Alternative frame. These frame help us with characterizations of curves.

The aim of this paper is to investigate focal curves according to alternative frame work in 3-dimensional Euclidean space. For this purpose, an alternative frame was first defined. Then, focal curves were given depending on this frame and various characterizations were obtained.

Keywords: Frenet frame, Focal curves, Alternative frame, Euclidean space

REFERENCES

1. Uribe-Vargas, R. On Vertices, Focal Curvatures And Differential Geometry Of Space Curves, Bull Braz Math Soc, 36(3), 285-307, 2005.
2. Alegre, P., Arslan, K., Carriazo, A., Murathan, C. and Öztürk, G. Some special types of developable ruled surface, Hacettepe Journal of Mathematics and Statistics, 39(3), 319-325, 2010.
3. Körpınar, T., Baş, S. and Asil, V. New Characterization of D-Focal Curves in Minkowski 3-space, Bol. Soc. Paran. Math., 38(2), 115-123, 2020.
4. Uzunoğlu, B., Gök İ. and Yaylı, Y. A new approach on curves of constant precession. *Applied Mathematics and Computation*, 275, 317-323, 2016.
5. Chen, B.Y. Constant ratio Hypersurface. *Soochow J.Math.*, 27(4), 353-362, 2001.

CUBIC HERMITE COLLOCATION METHOD FOR THE EQUAL WIDTH WAVE EQUATION

Nuri Murat Yağmurlu¹, Selçuk Kutluay¹, Ali Sercan Karakaş¹

¹Department of Mathematics, University of İnönü, Malatya, Turkey

murat.yagmurlu@inonu.edu.tr, selcuk.kutluay@inonu.edu.tr, ali_sercan_44@hotmail.com

Abstract

The fundamental aim of the present article is to numerically solve the non-linear Equal Width-Wave (EW) equation. For this purpose, the nonlinear term appearing in the equation is firstly linearized by Rubin-Graves type approach. After that, to reduce the equation into a solvable discretized linear algebraic equation system which is the essential part of this study, the Crank-Nicolson type approximation and cubic Hermite collocation method are respectively applied to obtain the integration in the temporal and spatial domain directions. To demonstrate how good the offered method generates approximate numerical results, six experimental problems exhibiting different wave profiles known as the motion of single, interacting two and three, the Maxwellian, undular bore and colliding soliton waves given with different initial and boundary conditions of the equation will be taken into consideration. Since only the first model problem has an exact solution, to measure error magnitudes widely used mean squared and maximum norms between exact and approximate solutions are calculated and also compared with those from other existing works in the literature. Furthermore, the three conservation constants known as mass, moment and energy quantities are also calculated and presented throughout the wave simulations with increasing time. In addition, a tabular comparison of the newly computed norms and conservation constants show that the current scheme produces better and compatible solutions than those of the most of the previous works with the same parameters. Apart from those, the stability analysis for this present scheme has been illustrated using the von Neumann method.

Keywords: Equal width-wave equation; cubic Hermite collocation method; solitary waves; stability analysis; Crank-Nicolson type approximation; Rubin-Graves type linearization.

REFERENCES

1. Kumar H. (2024) Numerical Study of Differential Equations Via Cubic Hermite Collocation Technique, NATURALISTA CAMPANO, 28(1)
2. S. Kutluay, N. M. Yağmurlu and A. S. Karakaş, An Effective Numerical Approach Based on Cubic Hermite B-spline Collocation Method for Solving the 1D Heat Conduction Equation, New Trends in Mathematical Sciences, 10, No. 4, 20-31 (2022) doi: <http://dx.doi.org/10.20852/ntmsci.2022.485>

This study is supported by Inonu University Scientific Research Project with project number FDK-2023-3402.

ON THE APPROXIMATION BY NONLINEAR OPERATORS OF MAX-PRODUCT KIND

Sevilay KIRCI SERENBAY¹, Ecem ACAR^{2*}

¹Department of Mathematics, Harran University, Şanlıurfa, Turkey

²Department of Mathematics and Science Education, Harran University, Şanlıurfa, Turkey

sevilaykirci@gmail.com, karakusecem@harran.edu.tr

Abstract

Max-product type operators which are a type of nonlinear positive operators, use the maximum and product operations to approximate functions. The max-product type operators have better order of approximation compared to linear operator sequences. In this paper, we introduce the nonlinear bivariate Bernstein max-product type operators by using the GBS operators (generalized Boolean sum) and we investigate their approximation properties by obtaining their rates of convergence.

Keywords: Max-product type operators; Bivariate operators; Order of approximation.

REFERENCES

1. Bede, B., Coroianu, L., Gal, S. G., Approximation by max-product type operators, Heidelberg: Springer, 2016.
2. Butzer, P.L., On two-dimensional Bernstein polynomials, Can. J. Math. 5, 107-113 , 1953.
3. Bögel K., Mehrdimensionale Differentiation von Funktionen mehrerer Veränderlicher, J Reine Angew Math., 170, 197-217, 1934.
4. E. Acar and A. İzgi, The Approximation of Bivariate Generalized Bernstein-Durrmeyer Type GBS Operators, International Journal of Maps in Mathematics, Vol:5, No:1, 2–20, 2022.

Examination Of Mhd Effect and Fractional Derivative Model Between Porous Medium Parallel Plates In Time-Dependent Flow

Kübra Heredağ¹ Fatma Ayaz²

¹Department of Mathematics, Gazi University , Ankara, Turkey

²Department of Mathematics, Gazi University , Ankara, Turkey

kubra.heredag@gazi.edu.tr, fayaz@gazi.edu.tr,

Abstract

This study investigates the effects of heat and mass transfer on time-dependent flow between porous parallel plates under the influence of a magnetic field. The time-dependent term in the governing equation is approximated by a fractional-order derivative. The partial differential equation system is solved using the finite difference method, with the Grünwald-Letnikov approach for the time derivative and the Crank-Nicolson method for other terms. The Thomas Algorithm is utilized for solving the equation system. Variations in the parameters of the equations are examined, and the resulting differences are presented graphically. It is observed that an increase in Gr and Gm numbers leads to an increase in the velocity profile, while a decrease in M results in an increase in velocity. A decrease in K corresponds to a decrease in temperature, and a decrease in Pr leads to a decrease in temperature. An increase in Sc causes an increase in concentration. Lastly, the change in the fractional derivative order α is investigated. A similar profile is observed for α values of 1, 0.9, and 0.8. As α decreases, the velocity, temperature, and concentration increase.

Keywords: Fractional Derivative; Grünwald Letnikov Approach,; Heat and Mass Transfer.

REFERENCES

1. Baytaş, A. C., Gözenekli ortamlarda taşınım olayı, İTÜDERGİSİ/c, 4(1), 2011.
2. Nishimoto, K., An Essence of Nishimoto's Fractional Calculus, Descarter Press Co, 1991.
3. Weilbeer, M., Efficient Numerical Methods for Fractional Differential Equations and their Analytical Background, PhD Thesis, Von der Carl-Friedrich-GaubFakultät für Mathematik und Informatik der Technischen Universität Braunschweig, Germany, 2005.

ON THE SEMI-ANALYTICAL AND HYBRID METHODS FOR THE DRINFELD-SOKOLOV-WILSON SYSTEM MODELLING DISPERSIVE WATER WAVES

Emre Aydın¹ İnci Çilingir Süngü²

¹ Department of Mathematics, University of Ondokuz Mayıs, Samsun, Turkey

emre_aydn_55@outlook.com,

² Department of Mathematics Education, University of Ondokuz Mayıs, Samsun, Turkey

incicilingir@gmail.com

Abstract

In this study, MVIM, MVILTM and MVISTM are used to solve the Drinfeld-Sokolov-Wilson (DSW) system. Semi-analytical solutions have been obtained for the DSW system. The exact solutions and semi-analytical solutions of the DSW system are compared with each other. Maximum errors of semi-analytical solutions of the DSW system for various iteration values are given by tables. Comparison of relative errors for various iteration values and effect of change of wave constant is visualized by figures. Also, it commented on the effectiveness and usefulness of the methods when applied to the DSW system.

Keywords: Modified variational iteration method (MVIM); Modified variational iteration Laplace transform method (MVILTM); Modified variational iteration Sumudu transform method (MVISTM); The Drinfeld-Sokolov-Wilson (DSW) system.

REFERENCES

1. Vladimir Gershonovich Drinfeld, Vladimir Sokolov, Equations of Korteweg-de Vries type and simple Lie algebras, Doklady Akademii Nauk. Russian Academy of Sciences, Vol:258, No:1, 11-16, 1981.
2. Vladimir Gershonovich Drinfeld, Vladimir Sokolov, Lie algebras and equations of Korteweg-de Vries type, Journal of Soviet Mathematics, Vol:30, 1975-2036, 1985.
3. Wei Gao, Pundikala Veerasha, D.G. Prakasha, Hacı Mehmet Baskonus, Gulnur Yel, A powerful approach for fractional Drinfeld-Sokolov-Wilson equation with Mittag-Leffler law, Alexandria Engineering Journal, Vol:58, No:4, 1301-1311, 2019.

THE COMPUTATION OF \mathcal{H}_∞ -NORM OF TRANSFER FUNCTIONS OF LINEAR DAEs VIA TWO-STEP METHOD

Hasan Gunduz¹, Mesut Karabacak², Ercan Celik³

¹Department of Mathematics, University of Bingol, Bingol, Turkey

²Department of Mathematics, University of Ataturk, Erzurum, Turkey

³Department of Applied Mathematics and Informatics, University of Kyrgyz- Turkish Manas, Byshkek, Kyrgyzstan

hgunduz@bingol.edu.tr,

Abstract

In this paper, we evaluate \mathcal{H}_∞ -norm of a transfer function of a linear DAEs system for special case $D = 0$, using two-step method, which is based on to associate poles of transfer function $G(s)$ to related Hamiltonian matrix. Since the method needs only one iteration then it is extremely fast when compared to other similar methods, that are used to compute \mathcal{H}_∞ -norm.

Keywords: DAEs Systems; \mathcal{H}_∞ -Norm; Two-Step Method; Hamiltonian Matrix; Singular Values.

REFERENCES

1. Kuster, G. E., \mathcal{H} Infinity-Norm Calculation via a State Space Formulation (Master of Science in Mathematics), Faculty of Polytechnic Institute, Blacksburg, Virginia, 2012.
2. Bruinsma, N. A., Steinbuch, M., A Fast Algorithm to Compute the \mathcal{H}_∞ Norm of a Transfer Function Matrix, Systems & Control Letters, 14(4):287–293, 1990.
3. Doyle, J., Glover, K., Khargonekar P., Francis, B., State-Space Solutions to Standard \mathcal{H}_2 and \mathcal{H}_∞ Control Problems, IEEE Trans. Automatic Control 34, 831- 847, 1989.
4. Enns, D. F., Model Reduction with Balanced Realizations: An Error Bound and a Frequency Weighted Generalization, Proceedings IEEE Conference on Decision and Control (Las Vegas, NV) 127-132, 1984.
5. Francis, B. A., A Course in \mathcal{H}_∞ Control Theory,(Springer-Verlag, Berlin-New York),1987.
6. Leibfritz, F., Lipinski, W., Description of Benchmark Examples in COMPl_eib 1.0 Tech. Report, 2003.
7. Gunduz, H., Celik, E., \mathcal{H}_∞ -Norm Evaluation for a Transfer Matrix via Bisection Method, Thermal Science, Vol 26, Special Issue 2:745-51, 2022.

Investigation of extended type a NLS equation using the extended direct algebraic method

Ahmed Abuhatim¹ and Ebru Cavlak Aslan¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

ebrucavlak@hotmail.com,

Abstract

In this study, the soliton solutions of the extended type nonlinear Schrödinger equation are investigated using an extended direct algebraic method. The solutions are found in the form of hyperbolic, trigonometric, and rational functions. Various types of well-known optical solitons, including dark, bright and combo optical soliton have been extracted here.

Keywords: Extended direct algebraic method; the extended type NLSE; Optical soliton; Soliton.

REFERENCES

1. Y. Ren, H. Zhang, New generalized hyperbolic functions and auto- Bäcklund transformation to find new exact solutions of the (2+1)- dimensional NNV equation, Physics Letters A, Vol: 357, No: 6, 438-448, 2006.
2. H. Yopez-Martinez, H. Rezazadeh, A. Souleymanou, S.P.T. Mukam, V.K. Kuetche, A. Bekir, The extended modified method applied to optical solitons solutions in birefringent fibers with weak nonlocal nonlinearity and four wave mixing, Chinese Journal of Physics, Vol: 58, 137-150, 2019.
3. W.B. Rabie, H.H. Hussein, H. M. Ahmed, M. Alnahhass, W. Alexan, Abundant solitons for highly dispersive nonlinear Schrödinger equation with sextic-power law refractive index using modified extended direct algebraic method, Alexandria Engineering Journal, Vol: 86, 680-689, 2024.

GROWTH OF SOLUTION FOR REACTION DIFFUSION EQUATION WITH KIRCHHOFF TERM AND MULTIPLE NONLINEARITIES

Muhteşem DEMİR¹

Erhan PİŞKİN²

¹ Ministry of National Education, Diyarbakır, Turkey,

² Department of Mathematics, Dicle University, Diyarbakır, Turkey

demirmuhtesem@gmail.com, episkin@dicle.edu.tr

Abstract

In this presentation, we consider a reaction diffusion equation with Kirchhoff term and multiple nonlinearities. We prove the growth of solution with negative initial energy under suitable conditions.

Keywords: Growth; Reaction diffusion equation; Kirchhoff equation.

REFERENCES

1. Pişkin E. ve Ekinçi F. 2020, Blow up and growth of solutions for a parabolic type Kirchhoff equation with multiple nonlinearities, *Konuralp Journal of Mathematics*, 8(1),216-222.
2. Pişkin E. ve Ekinçi F. 2019, Nonexistence and growth of solutions for a parabolic p-Laplacian system, *Sigma Journal of Engineering and Natural Sciences*, 10(3),301- 307.

CHARACTERIZATION OF PARAMETRIC SURFACES IN LIE GROUPS USING ALTERNATIVE FRAME

Betül OĞRAŞ İKİZ, Zühal KÜÇÜKARSLAN YÜZBAŞI

Department of Mathematics, Fırat University, Elazığ, Turkey

betulogras@hotmail.com

zuhal2387@yahoo.com.tr

Abstract

This study focused on describing a surface family by utilizing the alternative moving frame in the 3-dimensional Lie group. We determine the necessary and sufficient conditions for the geodesic curve, asymptotic curve, and line of curvature to serve as an isoparametric curve, respectively. Finally, we provided and visualized some examples based on the obtained results.

Keywords: Lie group; Frenet Frame; Alternative Frame.

REFERENCES

- [1] Küçükarslan Yüzbaşı Z., Bektaş M., On the construction of a surface family with common geodesic in Galilean space G_3 , *Open Phys.*, 14 (2016) 360–363.
- [2] Küçükarslan Yüzbaşı Z., Yoon D. W., On constructions of surfaces using a geodesic in Lie group, *J. Geo.*, 110(2) (2019) 1-10.
- [3] Yoon, D. W., Y Küçükarslan Yüzbaşı Z., Bektaş, M. (2017). An approach for surfaces using an asymptotic curve in Lie group. *Journal of Advanced Physics*, 6(4), 586-590.

IMPLEMENTATION OF BATTLE ROYALE OPTIMIZATION ALGORITHM FOR 0-1 KNAPSACK PROBLEM USING S-SHAPED AND V-SHAPED TRANSFER FUNCTIONS

Gülşen Orucova Büyüköz¹

¹ Department of Mathematics and Computer Science, Necmettin Erbakan University, Konya, Turkey

gorucova@erbakan.edu.tr,

Hüseyin Hakli²

² Department of Computer Engineering, Necmettin Erbakan University, Konya, Turkey

hhakli@erbakan.edu.tr

Abstract

In this study, the Battle Royale Optimization Algorithm is applied to 0-1 Knapsack Problems (0-1 KP), which are in the NP-hard problem class. The Battle Royale optimization algorithm proposed for continuous problems cannot be directly applied to 0-1 KP problems with binary structure. Therefore, S-shaped and V-shaped transfer functions are used to adapt the continuous search space to the binary search space. The proposed algorithm is used to solve 0-1 KP problems and the results obtained are compared to determine which transfer function gives more effective results. The proposed algorithm is also compared with different algorithms in the literature.

Keywords: Battle Royale Optimization Algorithm, 0-1 Knapsack Problems, Transfer Functions, Binary Optimization.

REFERENCES

1. Dantzig, G.B., 1957. Discrete-Variable Extremum Problems, Source: Operations Research.
2. Roederkerk, R.P. and van Heerde, H.J., 2016. Robust optimization of the 0-1 knapsack problem: Balancing risk and return in assortment optimization. European Journal of Operational Research, 250(3), pp. 842-854.
3. Rahkar Farshi, T. (2021). Battle royale optimization algorithm. Neural Computing and Applications, 33(4), 1139-1157.

HOUSING PRICE DETERMINANTS: A BIG SPATIAL DATA ANALYSIS

Bahadır Yüzbaşı¹

¹Department of Econometrics, University of Inonu, Malatya, Turkey

b.yzb@hotmail.com,

Abstract

The purpose of this paper is to use a new approach to predict the underlying prices of properties in a city. To evaluate the efficacy of this technique, a thorough investigation of the spatial regression strategy is undertaken through the utilization of geostatistical-based Monte Carlo simulations and the analysis of real data collected from Airbnb.

Keywords: Spatial regression; Big spatial data; Machine Learning.

REFERENCES

1. Ahmed, S. E., Ahmed, F., & Yüzbaşı, B. (2023). Post-shrinkage strategies in statistical and machine learning for high dimensional data. Chapman and Hall/CRC.
2. Fotheringham, A. S., Brunsdon, C., & Charlton, M. E. (2009). Geographically weighted regression. The Sage handbook of spatial analysis, 1, 243-254.
3. Harris, P. (2019). A simulation study on specifying a regression model for spatial data: choosing between autocorrelation and heterogeneity effects. Geographical Analysis, 51(2), 151-181.

PSEUDOSPECTRAL CHEBYSHEV APPROACH FOR NONLINEAR TEMPERATURE DISTRIBUTIONS IN FUNCTIONALLY GRADED DISKS

Enver Temo¹, Mehmet Eker², Durmuş Yarımpanbuç^{1*}

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

²Department of Mechanical Engineering, Tarsus University, Mersin, Turkey

envertemmo@gmail.com, mehmeteker@tarsus.edu.tr, durmusyarimpabuc@osmaniye.edu.tr,

Abstract

In this study, the nonlinear heat conduction of a functionally graded disk under various temperature loads is discussed using pseudospectral Chebyshev method. Among the thermal material properties of disk are assumed to be graded in radial direction with Voigt homogenization scheme. Beside, due to its temperature sensitivity, the heat conduction coefficient is taken to depend on the temperature. While the inner surface of the disk is subjected to constant base temperature load, the outer surface is assumed to be under both Dirichlet and Neumann boundary conditions. Also, the effect of the convection coefficient is examined to model different operating conditions. These conditions lead to nonlinear differential equations that conventional methods may not solve. The nonlinear temperature distributions of the functionally graded disk under the thermo-mechanical loads are determined and illustrated graphically.

Keywords: Nonlinear temperature distribution; pseudospectral Chebyshev method; Voigt homogenization scheme; Functionally graded materials.

REFERENCES

1. Surja Deka, Ashis Mallick, Pratyush P. Behera, Prakash Thamburaja, Thermal stresses in a functionally graded rotating disk: An approximate closed form solution. Journal of Thermal Stresses, Vol:44, No: 1, 20-50, 2020.
2. Rajiv Ranjana, Ashis Mallicka, and Prasun, Thermoelastic study of a functionally graded annular fin with variable thermal parameters using semiexact solution. Journal of Thermal stresses, Vol:42, No: 10, 1272-1297, 2019.
3. David Gönczi, Istvan Ecsedi, Thermoelastic analysis of functionally graded hollow circular disc. Archive of Mechanical Engineering, Vol:5, No:18, 2015.

New Optical Soliton Solutions of the NLS Equation with Jacobi Elliptic Function Expansion Method

Derya DENİZ¹ and Ebru CAVLAK ASLAN¹

¹Department of Mathematics, University of Firat, Elazığ, Turkey

deryadeniz485@yandex.com , ebrucavlak@hotmail.com

Abstract

In this presentation optical soliton solutions are extracted for the proposed equation using the Jacobi elliptic function expansion technique. To better comprehend the dynamic characteristics of the retrieved solutions their graphical visualization are provided.

Keywords: Jacobi elliptic function expansion method; Soliton; Optical Soliton; NLS equation.

REFERENCES

1. T. A. Khalil, N. Badra, H. M. Ahmed, W. B. Rabie, Optical solitons and other solutions for coupled system of nonlinear Biswas–Milovic equation with Kudryashov’s law of refractive index by Jacobi elliptic function expansion method, Optik, Vo: 253, No: 168540, 2022.
2. E. Cavlak Aslan, L. Gürgöze, Soliton and Other Function Solutions of the Potential KdV Equation with Jacobi Elliptic Function Method, International Journal of Innovative Engineering Applications, Vol: 6, No: 2, 2022.
3. W. B. Rabie and H. M. Ahmed, “Cubic-quartic solitons perturbation with couplers in optical metamaterials having triple-power law nonlinearity using extended F-expansion method,” Optik (Stuttg.), Vol. 262, No. 169255, 2022.
4. T. Yazgan, G. Yel, E. Celik and H. Bulut, ”On Survey of the Some Wave Solutions of the Non-Linear Schrödinger Equation (NLSE) in Infinite Water Depth,” GU J Sci, Vol. 36, No. 2, 819-843, 2023.

MOTION OF THE FILAMENT IN MINKOWSKI SPACE

Zühal KÜÇÜKARSLAN YÜZBAŞI¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

zuhal2387@yahoo.com.tr,

Abstract

This research paper provides an in-depth geometric explanation for nonlinear partial differential equations (NPDEs). It achieves this by utilizing three different spacelike curves that demonstrate geometric flow motion, respectively.

Keywords: Curve Flow; Minkowski Space; Spacelike Curve.

REFERENCES

1. Yuzbasi, Z.K., Yoon, D.W., Myrzakulov, R. Mapping between space-like curve flows and soliton equations in Minkowski space. Int. J. Geom. Methods Mod. Phys., (2023), doi.org/10.1142/S0219887824500804.
2. Sun, X., Wang, Y. (2011). KdV geometric flows on Kähler manifolds. Int. J. Math., 22(10), 1439-1500.
3. Ding, Q., Wang, Y.D. Geometric KdV flows, motions of curves and the third-order system of the AKNS hierarchy. Int. J. Math., 22 (2011), 1013-1029.

ON THE UNIFORMLY PARIKH-FRIENDLY WORDS

Mehmet Uçar ¹, Aynur Yalçın ²

^{1,2}Department of Mathematics, University of Selcuk, Konya, Türkiye

¹ 218214001001@lisansustu.selcuk.edu.tr, ² ayalciner@selcuk.edu.tr

Abstract

Mateescu and others [1] introduced Parikh matrix mapping as a sharpening of the Parikh mapping, where somewhat more information is preserved than in the original Parikh mapping. The Parikh matrix of a word is an upper triangular matrix and the classical Parikh vector appear in such a matrix as the second diagonal. All other entries above the main diagonal contain information on the number of occurrences of certain subwords of that word. Şerbanuta [3] defined the Parikh matrix mapping in a way that expands it to the Parikh matrix mapping created according to a word u , instead of defining it according to an $\Sigma_k = \{a_1 < a_2 < \dots < a_k\}$ ordered alphabet. The extending Parikh matrix coincides with Parikh matrix if it is defined with respect to the word $u = a_1 a_2 \dots a_k$. Salomaa [2] introduced the notion of Parikh-friendly permutation and posed the characterization of Parikh-friendly permutations as open problem. Teh [4] showed that every permutation is Parikh-friendly. Also, he defined the uniformly Parikh-friendly words and proved the existence of the uniformly Parikh-friendly words for every alphabet Σ .

In this talk, we will give a generalization of uniformly Parikh-friendly words.

Keywords: Parikh matrix mapping; Parikh-friendly permutation; uniformly Parikh-friendly word.

REFERENCES

1. Alexandru Mateescu, Arto Salomaa, Kai Salomaa, Sheng Yu, A sharpening of the Parikh mapping, *Theoretical Informatics and Applications*, Vol:35, 551–564, 2001.
2. Arto Salomaa, Parikh matrices: Subword indicators and degrees of ambiguity, In *Adventures Between Lower Bounds and Higher Altitudes*, Lecture Notes in Computer Science Springer, Cham, Vol:11011, 100–112, 2018.
3. Traian-Florin Şerbănuță, Extending Parikh matrices, *Theoretical Computer Science*, Vol: 310, 233–246, 2004.
4. Wen Chean Teh, Parikh-friendly permutations and uniformly Parikh-friendly words, *Australasian Journal of Combinatorics*, Vol:76, No:1, 208–219, 2020.

A MESHFREE METHOD FOR NUMERICAL SOLUTIONS OF SOME REACTION-DIFFUSION TYPE EQUATIONS

Ömer Oruç¹

¹Department of Mathematics, Dicle University, Diyarbakır, Turkey

omer.oruc@dicle.edu.tr,

Abstract

This study proposes a numerical approach to obtain approximate solutions of some nonlinear partial differential equations (PDEs) of reaction-diffusion type. To this end, for discretization of space variable a meshfree method based on radial basis function differential quadrature method is used. For discretization of time variable of considered PDE, a finite difference method is used. In this way, a full discrete system is obtained whose solution is used for construction of approximate solution. Some test problems are solved via proposed method. Acquired approximate solutions are compared with exact solutions and with some methods available in literature. From the comparisons, accuracy and feasibility of the proposed method are confirmed

Keywords: Radial basis function differential quadrature; Reaction-diffusion PDE; Numerical method.

REFERENCES

1. J. Nagumo, S. Arimoto, S. Yoshizawa, An active pulse transmission line simulating nerve axon, Proceedings of the IRE, 50, 2061–2070, 1962.
2. R. Fitzhugh, Impulse and physiological states in models of nerve membrane. Biophys Journal, 1(6), 445–66, 1961.

Optimal culling strategy for the fractional-order brucellosis transmission model

Dilara Yapişkan¹ and Beyza Billur İskender Eroğlu^{1,*}

¹Department of Mathematics, Balıkesir University, Balıkesir, Turkey

* biskender@balikesir.edu.tr

Abstract

We suggest and solve an optimal culling problem within the fractional-order brucellosis model between interspecies. We aim to reduce exposed and infected sheep with the culling strategy while also minimizing the health and economic burden associated with brucellosis. We characterize the fractional optimal control with the Pontryagin Maximum Principle approach, considering the necessary optimality conditions proposed for the Atangana-Baleanu derivative [1,2], and solve the problem numerically by implementing the Adams-type predictor-corrector algorithm combined with the forward-backward sweep method [3].

Keywords: Adams-type predictor-corrector algorithm; Atangana- Baleanu; brucellosis; culling; optimal control;

REFERENCES

1. Mohamed Gaber Bahaa, Abdon Atangana, Necessary and sufficient optimality conditions for fractional problems involving atangana-baleanu's derivatives, in Fractional Derivatives with Mittag-Leffler Kernel: Trends and Applications in Science and Engineering edited by J. F. Gomez, L. Torres, and R. F. Escobar, 194 Springer, Cham, 13–33, 2019.
2. Lev Pontryagin, Mathematical theory of optimal processes. CRC Press, London, 1987.
3. Ismail Gad Ameen, Dumitru Baleanu, Hegagi Mohamed Ali, An efficient algorithm for solving the fractional optimal control of SIRV epidemic model with a combination of vaccination and treatment, Chaos Solit. Fractals., Vol:137, No:2020, 109892, 2020.

POINTWISE HEMI-SLANT SUBMERSIONS FROM COSYMPLECTIC MANIFOLDS

Meltem Karaismailoğlu¹, Sezin Aykurt Sepet², Mahmut Ergüt³

^{1,2} Kırşehir Ahi Evran University, Art and Science Faculty, Department of Mathematics,
Kırşehir, Turkey

³ Tekirdağ Namık Kemal University, Art and Science Faculty, Department of Mathematics,
Tekirdağ, Turkey

mltm.33.27@hotmail.com, saykurt@ahievran.edu.tr, mergut@nku.edu.tr

Abstract

In this paper, we study pointwise hemi-slant submersions as a generalization of pointwise slant submersions and hemi-slant submersions from cosymplectic manifolds onto Riemannian manifolds. We investigate the integrability of distributions and the geometry of totally geodesic foliations which arise from the definition of such submersions. Moreover, we study the φ -pluriharmonicity of such maps.

Keywords: Riemannian submersion; pointwise hemi-slant submersion; cosymplectic manifold; pluriharmonicity.

REFERENCES

1. M.A. Akyol, Y. Gündüzalp, Hemi-slant submersions from almost product Riemannian manifolds, Gulf Journal of Mathematics, Vol. 4, 15-27, Issue 3(2016).
2. A. Gray, Pseudo-Riemannian almost product manifolds and submersions, J. Math. Mech., 715-737, 16(1967).
3. Y. Ohnita, On pluriharmonicity of stable harmonic maps, J. Lond. Math. Soc., 563-568, 2(1987), <https://dx.doi.org/10.1112/jlms/s2-35.3.563>.
4. B. O'Neill, The fundamental equations of a submersion, Michigan Math. J., 458-469, 13(1966).
5. B. Şahin, Slant submersions from almost Hermitian manifolds, Bull Math. Soc. Sci. Math. Roumanie, 93-105, 54(2011).
6. H.M. Taştan, B. Şahin, Ş. Yanan, Hemi-slant submersions, Mediterr J. Math., DOI:10.1007/s00009-015-0602-7.

MULTIPLICATIVE RECTIFYING CURVE IN MULTIPLICATIVE EUCLIDEAN SPACE

Muhittin Evren Aydın¹ Aykut Has^{2*}, Beyhan Yılmaz³

¹Department of Mathematics, Firat University, Elazig, Turkey

^{2*}Department of Mathematics, Kahramanmaraş Sutcu Imam University, Turkey

³Department of Mathematics, Kahramanmaraş Sutcu Imam University, Turkey

meaydin@firat.edu.tr, ahas@ksu.edu.tr, beyhanyilmaz@ksu.edu.tr

Abstract

In this talk the rectifying curves are studied in terms of the tools from multiplicative differential geometry. By several characterizations, we classify such curves in the multiplicative Euclidean space by means of the curves on a multiplicative sphere. Several examples are also given by figures.

Keywords: Rectifying curve; spherical curve, multiplicative calculus; multiplicative Euclidean space.

Acknowledgements: This study was supported by Scientific and Technological Research Council of Turkey (TUBITAK) under the Grant Number 123F055. The authors thank to TUBITAK for their supports.

REFERENCES

1. Georgiev S.G., Multiplicative Differential Geometry (1st ed.), Chapman and Hall/CRC., New York, 2022.
2. Chen B.-Y., When Does the Position Vector of a Space Curve Always Lie in Its Rectifying Plane?, The American Mathematical Monthly, 110(2) (2003), 147-152.
3. Deshmukh, S., Chen, B.-Y., Alshammari, S. H., On rectifying curves in Euclidean 3-space, turk. J. Math. 42 (2018), 609-620.

Solution of Fractional Order Partial Differential Equations with Hosoya Neural Network

Merve Zeynep KAYA^{1,*}, Ercan ÇELİK², Mesut KARABACAK³

¹Distance Education Center, Agri Ibrahim Cecen University, Agri, Turkey

²Department of Applied Mathematics and Informatics, Faculty of Science, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan

³ Department of Mathematics, Faculty of Science, Ataturk University, Erzurum, Turkey

mzgecmen@agri.edu.tr, ercan.celik@manas.edu.kg, mkarabacak@atauni.edu.tr

Abstract

In this article, we propose an innovative method that leverages the Hosoya neural network to solve fractional-order reaction-diffusion equations. The architecture of the Hosoya neural network comprises input, hidden, and output layers, each consisting of perceptrons. Specifically, we employ various degrees of the Hosoya polynomial as training functions for the hidden layer. Subsequently, the fractional-order diffusion equation is transformed into an optimization problem. In the subsequent step, we obtain approximate and exact solution graphics using the Python programming language within a specific algorithm. Notably, the results obtained from this study demonstrate greater practicality and efficiency compared to conventional solutions.

Keywords: Partial Differential Equations; Neural Network; Hosoya Polynomial; Fractional Calculus; Reaction Diffusion Equation

REFERENCES

1. Podlubny I., Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications. Elsevier, New York, 1998.
2. Dwivedi, K. D. and Rajeev, Numerical solution of fractional order advection reaction diffusion equation with Fibonacci neural network, Neural Processing Letters 53: 2687-2699.
3. Bishop, M. J., History and philosophy of neural networks, 2015.

A NEW FACIAL EXPRESSION RECOGNITION METHODS BASED ON HYBRID FEATURE

Muhammed Kerem Turkes¹ Yıldız Aydın¹

¹ Department of Computer Engineering, Erzincan Binali Yildirim University,

Erzincan 24000, Turkey

mktrks1@gmail.com, yciltas@erzincan.edu.tr

Abstract

Within the scope of this study, an application that can predict emotions from people's facial expressions was developed. This developed application consists of two steps: feature extraction and classification. It is observed that there is a significant increase in classifier performance by obtaining robust features, which is an essential factor when extracting features. For this reason, in recent years, hybrid features have been received by combining features instead of classical feature extraction, and more robust features have been used to get more successful results in problems. In this study, support vector machine (SVM), XGBoost (XGB), logistic regression (LR) and random decision trees (RF) classification methods were used to compare the performance of classical features such as SIFT, SURF, KAZE and the HessianSIFT hybrid feature obtained by building the SIFT descriptor on the key points detected with the Hessian detector. The SIFT feature achieved the highest success rate of 95.41% accuracy with the SVM classifier, the SURF feature achieved the highest success rate of 86.75% accuracy with the SVM classifier, and the KAZE feature achieved the highest success rate of 93.07% accuracy with the RF classifier. CK+48 data set was used to obtain the study's performance data. As a result, by classifying the hybrid feature created using the Hessian detector and SIFT descriptive using the SVM method, the classification performance was significantly improved and a success rate of 98.67% was achieved.

Keywords: Robust Features; HessianSIFT; HOG+LBP; SURF; Classification; SVM.

REFERENCES

1. Aydın Yıldız, Automated identification of copy-move forgery using Hessian and patch feature extraction techniques, Journal of Forensic Sciences, Vol:69, No:1, 131-138, 2024.
2. Berretti Stefano, Ben Amor Boulbaba, Daoudi Mohamed, Del Bimbo Alberto, 3D facial expression recognition using SIFT descriptors of automatically detected keypoints. The Visual Computer, Vol:27, 1021-1036, 2011.
3. Turan Cigdem, Lam Kin-Man, Histogram-based local descriptors for facial expression recognition (FER): A comprehensive study. Journal of visual communication and image representation, Vol:55, 331-341, 2018.

f –STATISTICAL CONVERGENCE OF DOUBLE SEQUENCES IN TOPOLOGICAL GROUPS

Suleyman Sarikaya and Yavuz Altin

Department of Mathematics, Firat University, Elazig, Turkey

ssarikaya@firat.edu.tr, yaltin23@yahoo.com

Abstract

In 2003, Mursaleen and Edeley [5] first introduced the idea of statistical convergence of double sequences. Later, Cakallı and Savaş [3] defined this concept in topological groups. In this study, we gave the concept of f –statistical convergence for double sequences in topological groups by using the concept of unbounded modulus function. We also gave some inclusion theorems.

Keywords: Statistical convergence; Double sequence; Modulus function; Topological groups.

REFERENCES

1. Aizpuru, A., Listán-García M.C., Rambla-Barreno F. Density by moduli and statistical convergence, Quaestiones Mathematicae, (2014), 37 (4), 525-530.
2. Cakalli, H. Lacunary statistical convergence in topological groups, Indian Journal of Pure and Applied Mathematics, (1995), 26 (2), 113-119.
3. Çakalli, H., Savaş E. Statistical convergence of double sequences in topological groups, Journal of Computational Analysis and Applications (2010), 12 (2), 421-426.
4. Fast, H. Sur la convergence statistique, Colloquium Mathematicum, (1951), 2, 241-244.
5. Fridy, J.A. On statistical convergence, Analysis (1985), 5, 301-313.
6. Mursaleen, Osama H. H. Edely. Statistical convergence of double sequences, Journal of Mathematical Analysis and Applications (2003), 288, 223-231.

AN INVESTIGATION OF A FUZZY BOUNDARY VALUE PROBLEM

Hülya Gültekin Çitil¹, Fatma Gizem Özmen²

¹Department of Mathematics, Faculty of Arts and Sciences, Giresun University, Giresun, Turkey

² Institute of Science, Giresun University, Giresun, Turkey

hulya.citil@giresun.edu.tr, gzzmsyn@gmail.com

Abstract

In this paper, we investigate a fuzzy boundary value problem. We obtain fundamental results for solutions of the problem. Also, we prove when the solutions are valid fuzzy functions.

Keywords: Fuzzy boundary value problem; Fuzzy function; Fuzzy number.

REFERENCES

1. S. Tudu, S. P. Mondal, A. Ahmadian, A. K. Mahmood, S. Salahshour, M. Ferrara, Solution of generalised type – 2 Fuzzy boundary value problem, Alexandria Engineering Journal, 60(2), 2725-2739, 2021.
2. H. Gültekin Çitil, On a Boundary Value Problem with Fuzzy Forcing Function and Fuzzy Boundary Values, International Journal of Mathematical Combinatorics, 2, 1-16, 2021.
3. M. Chehlabi, T. Allahviranloo, Positive or negative solutions to first-order fully fuzzy linear differential equations under generalized differentiability, Applied Soft Computing, 70, 359-370, 2018.

EXPLAINING OF DECISION MAKING PROCESSES WITH THE HELP OF INTUITIONISTIC FUZZY SETS

Feride Tuğrul¹

¹ Department of Computer Engineering, Faculty of Engineering,

Munzur University, Tunceli, Türkiye

feridetugrul@munzur.edu.tr

Abstract

Mathematics and logic are deeply interconnected disciplines, with logic providing the framework for mathematical reasoning and mathematics contributing to the development of logical systems. Fuzzy logic provides a mathematical framework for dealing with vague and uncertain information. Intuitionistic fuzzy logic extends the concepts of fuzzy logic by incorporating hesitation and non-membership degrees alongside membership degrees, providing a richer framework for modeling uncertainty and ambiguity. Intuitionistic fuzzy logic is particularly well-suited for decision making problems where decisions need to be made based on multiple conflicting criteria. In such scenarios, intuitionistic fuzzy sets can represent the varying degrees of satisfaction or preference for different criteria, and decision making algorithms can be designed to take into account both the degrees of membership and non-membership in the decision process. In this study, a study was conducted on the applications of intuitionistic fuzzy logic and decision making methods. The methods themselves, which play an important role in the decision making processes, are interpreted by taking into account factors such as weight functions and the importance of decision makers, and their real life results through an original application.

Keywords: Intuitionistic fuzzy sets; Decision making.

REFERENCES

1. Krassimir Atanassov,. Intuitionistic Fuzzy Sets, VII ITKR Session, Sofia, 20-23, (Deposed in Centr. Sci.-Techn. Library of the Bulg. Acad. of Sci., 1697/84) (in Bulgarian), (Reprinted: Int. J. Bioautomation 20(S1) 2016), June, 1983.

A NEW PARANORMED SEQUENCE SPACE GIVEN BY JORDAN TOTIENT FUNCTION

Pınar Zengin Alp¹

¹Department of Mathematics, University of Duzce, Düzce, Turkey

pinarzenginalp@gmail.com,

Abstract

In this paper, we give a new paranormed sequence space by using the regular matrix given by Jordan Totient function and we prove that this space is linearly isomorphic to $\ell(p)$. Also we compute α -, β -, γ - duals and the Schauder basis of this space.

Keywords: Paranormed sequence space, α -, β -, γ - duals, Jordan Totient function.

REFERENCES

1. İlkan, Merve, Necip Şimşek, and Emrah Evren Kara, A new regular infinite matrix defined by Jordan totient function and its matrix domain in ℓ_p , Mathematical Methods in the Applied Sciences Vol:44,No:9, 7622-7633, 2021.
2. İlkan, Merve, Serkan Demiriz, and Emrah Evren Kara, A new paranormed sequence space defined by Euler totient matrix, Karaelmas Fen ve Mühendislik Dergisi Vol:9, No:2, 277-282, 2019.
3. Maddox, I.J, Spaces of strongly summable sequences, The Quarterly Journal of Mathematics Vol:18, No:1, 345-355, 1967.

Detecting Android Malware Using LightGBM: A Study on the TUANDROMD Dataset

Muhammed Veysi Güler^{1*}, Muhammed Emre Çolak¹

¹ Department of Artificial Intelligence and Data Engineering, Firat University, Elazig, Türkiye

(mvguler@firat.edu.tr, memrecolak@firat.edu.tr)

Abstract

This study emphasizes the importance of security measures for users and organizations facing an increasing number of malicious software threats. The rapid evolution of these threats and the inadequacy of traditional detection methods necessitate the development of new and effective detection techniques. Therefore, this study aims to strengthen both user and organizational digital security by utilizing lightweight and powerful machine learning algorithms such as LightGBM to detect malicious software on the Android platform. Within this context, a model has been developed for detecting Android malware and goodware using the Dataset TUANDROMD. The TUANDROMD dataset comprises 241 features from 4465 Android applications, and techniques such as smoothing and synthetic data generation were employed to address dataset imbalances. The model was trained using LightGBM, a lightweight gradient boosting framework. It achieved an accuracy of 99.44% on the test set, with precision, recall, and F1-score values of 99% for both classes. Additionally, specificity values of 99.42% for class 0 and 99.45% for class 1 were observed. According to the confusion matrix results, only four false positive/negative predictions were made for both classes. These results demonstrate the high performance potential of LightGBM for Android malware detection and the effectiveness of enhancing the dataset through synthetic data generation.

Keywords: Android malware detection, Classification performance, LightGBM, Machine learning, Synthetic data generation, TUANDROMD

REFERENCES

- [1] P. Borah, D. Bhattacharyya, ve J. Kalita, “Malware Dataset Generation and Evaluation”, içinde *2020 IEEE 4th Conference on Information & Communication Technology (CICT)*, Ara. 2020, ss. 1-6. doi: 10.1109/CICT51604.2020.9312053.
- [2] B. Wu *vd.*, “Why an Android App is Classified as Malware? Towards Malware Classification Interpretation”. arXiv, 04 Eylül 2020. doi: 10.48550/arXiv.2004.11516.
- [3] D. Arp, M. Spreitzenbarth, M. Hübner, H. Gascon, ve K. Rieck, “Drebin: Effective and Explainable Detection of Android Malware in Your Pocket”, içinde *Proceedings 2014 Network and Distributed System Security Symposium*, San Diego, CA: Internet Society, 2014. doi: 10.14722/ndss.2014.23247.
- [4] N. G. Ambekar, N. N. Devi, S. Thokchom, ve Yogita, “TabLSTMNet: enhancing android malware classification through integrated attention and explainable AI”, *Microsyst. Technol.*, Mar. 2024, doi: 10.1007/s00542-024-05615-0.

ON THE CONSTRUCTION OF A TOPOLOGY ON A ROUGH SEMIGROUP

Nurettin BAĞIRMAZ¹

¹Department of Mathematics, Mardin Artuklu University, Mardin, Turkey

nurettinbagirmaz@artuklu.edu.tr,

Abstract

The aim of this talk is to define the topology of a rough semigroup on an approximation space. Moreover, some basic properties and examples are presented.

Keywords: Rough sets, rough semigroup, topological semigroup, topological rough semigroup.

REFERENCES

1. Z. Pawlak, Rough sets, Int. J. Comput. Inform. Sci. 11 (1982) 341.356.
2. N. Bağirmaz, A. F. Özcan, Rough semigroups on approximation spaces, International Journal of Algebra, Vol. 9, 2015, no. 7, 339-350.
3. R. Biswas, S. Nanda, Rough groups and rough subgroups, Bull. Polish Acad. Sci. Math 42 (1994) 251-254.
4. N. Kuroki, Rough ideals in semigroups, Inform. Sci. 100 (1997) 139.163.
6. J. H. Carruth, J. A. Hildebrandt, R. J. Koch, The theory of topological semigroups, Marcel Dekker Inc, 1983.

VOTING CLASSIFIER BASED EXPLAINABLE ARTIFICIAL INTELLIGENCE METHOD FOR DETECTING GLIOMA GRADING USING CLINICAL AND MUTATION FEATURES

Ömer Miraç Kökçam^{1*}, Muhammed Emre Çolak², Özal Yıldırım²

¹ Department of Software Engineering, Fırat University, Elazığ, Türkiye

² Department of Artificial Intelligence and Data Engineering, Fırat University, Elazığ, Türkiye

(omkokcam@firat.edu.tr, memrecolak@firat.edu.tr, ozalyildirim@firat.edu.tr)

(0000-0003-1099-7513, 0000-0003-2317-2385, 0000-0001-5375-3012)

Abstract

In this study, a voting classifier based explainable approach is proposed for glioma grading detection. Logistic Regression, K Nearest Neighbor, Support Vector Machine and AdaBoost are used in this system, which gives output with a voting system in line with the decision outputs of each classifier on the input data in a structure where different classifier structures are used together. Local Interpretable Model-agnostic Explanations method was used to provide an explainable structure of which features the classifiers focus on when making decisions on the input data.

In the experimental studies, the performance of Logistic Regression, K Nearest Neighbour, Support Vector Machine and AdaBoost algorithms were 93.45%, 92.86%, 94.05% and 90.48% respectively. In this study, where classifiers were used together in various combinations, the accuracy rate reached to 92.86% when all classifiers were used with the voting classifier. The obtained results will play an effective role in creating reliable models in the field of health by determining that each classifier can make different decisions on the same input data and according to which criteria they make these decisions. This study has made an important contribution in terms of its explainable aspect and creating a system where different classifiers will make a common decision.

Keywords: *Biomedical Engineering, Explainable Artificial Intelligence, Glioma Grading, Machine Learning, Voting Classifier,*

REFERENCES

[1] A. S. Alhasan, “Clinical Applications of Artificial Intelligence, Machine Learning, and Deep Learning in the Imaging of Gliomas: A Systematic Review,” *Cureus*, Nov. 2021, doi: 10.7759/cureus.19580.

A COMPARATIVE ANALYSIS OF TREE-INSPIRED FRACTAL BRANCHINGS DENDRIFORM STRUCTURES, FROM THE BC TO THE L-SYSTEM BASED STRUCTURES.

Zeynep Gülcan Kaya^{1*}, Murat Şahin¹, Ayça Gülten¹

¹ Department of Architecture, University of Firat, Elazığ, Turkey

(zeynepgulcankaya@gmail.com, msahin@firat.edu.tr, aaytac@firat.edu.tr)

(0009-0005-9427-0996, 0000-0001-6733-1136, 0000-0001-9837-8674)

Abstract

Prior to the advent of fractal geometry, the principles of natural geometry were enigmatic and could not be elucidated through conventional methodologies. In the present era, it is possible to employ quantitative techniques to describe the majority of geometrical configurations. This study examined the compatibility of geometric compositions when transferring from Euclidean geometry-based hand-drawing environments to digital drawing and computational CAD tools in architectural products with complex parametric designs. The study considered historical structures such as Saint Chapelle, Gloucester Cathedral, La Sagrada Familia, and Stuttgart Airport Terminal, as well as contemporary examples generated by parametric design tools. In this study, both past and current examples of parametric design were utilized as benchmarks. The plans and column head views of buildings were analyzed using the Fractal Analysis Method with software called FracLac, which acts as a plug-in within ImageJ. Sophisticated column geometry was generated as dynamic geometry using L-System rules and iteration principles. A solid substance was then constructed using Dynamo-PythonScript, which served as an interface command within Autodesk Revit. It thus appears that, contrary to popular belief, the geometry of dendriform structures is not particularly complex, yet exhibits unexpected behavior.

Keywords: Computational architecture, Dendriform, Fractal geometry, Iterative generation method Parametric design.

REFERENCES

- [1] L. Nouri, K. T. Azari, and M. Alaghmandan, “Development of Algorithmic Applications in Architecture :A Review and Analysis of L-Systems: Bagh-e Nazar,” *Bagh-e Nazar*, vol. 19, no. 116, pp. 5–24, Feb. 2023, doi: 10.22034/BAGH.2022.327468.5119.
- [2] L. Moltedo, G. Mortelliti, O. Salvetti, and D. Vitulano, “Computer aided analysis of the buildings,” *Journal of Cultural Heritage*, vol. 1, no. 1, pp. 59–67, Jan. 2000, doi: 10.1016/S1296-2074(99)00115-6.

Multi-category classification of inappropriate content on social media using Natural Language Processing techniques and Transformer Models

Mohammed TALEB¹, Nouredine En-Nahnahi² and Nisrine Dad³

¹ LISAC Laboratory, Faculty of Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

mohammed.taleb1@usmba.ac.ma,

² LISAC Laboratory, Faculty of Sciences Dhar El Mahraz, Sidi Mohamed Ben Abdellah University, Fez, Morocco

nouredine.en-nahnahi@usmba.ac.ma,

³ SmartiLab Laboratory, Moroccan School of Engineering Sciences (EMSI), Rabat, Morocco

dad.nisrine@gmail.com,

Abstract

The expansion of social media utilization has led to a surge in user-generated content, presenting critical challenges in online content moderation task. Identifying and categorizing inappropriate content, such as offensive, toxic, sarcastic, and sexist text, has become imperative. In this paper, we present a novel approach to multi-category classification of inappropriate content on social media platforms utilizing advanced Natural Language Processing (NLP) techniques and Transformer-based Models[1] . Particularly, we explore the use of fine-tuned versions of BERT[2] and RoBERTa[3] pre-trained Transformer-based models for our classification task. Our strategy includes preprocessing the input content and training these models on a curated dataset sourced from more than ten datasets dealing with inappropriate content. We illustrate the viability of our approach by accomplishing accuracy scores of 0.98 and 0.97 for improper text detection and classification utilizing the fine-tuned BERT and RoBERTa models, respectively. Our findings emphasize the potential of leveraging state-of-the-art NLP techniques for strong detection of inappropriate content in social media.

Keywords: Natural Language Processing; BERT; RoBERTa; Offensive language; sexism detection ; toxic; sarcasm.

REFERENCES

- [1] A. Vaswani *et al.*, “Attention Is All You Need,” Jun. 2017.
- [2] J. Devlin, M.-W. Chang, K. Lee, and K. Toutanova, “BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding,” Oct. 2018.
- [3] Y. Liu *et al.*, “RoBERTa: A Robustly Optimized BERT Pretraining Approach,” Jul. 2019.

INVERSE SCATTERING PROBLEM FOR DISCONTINUOUS STURM-LIOUVILLE OPERATOR

Özge Akçay¹

¹ Department of Computer Engineering, Munzur University, Tunceli, Turkey

ozgeakcay@munzur.edu.tr,

Abstract

In this paper, the inverse scattering problem for Sturm-Liouville operator with discontinuous coefficient and transmission conditions at some point on the positive half plane is examined. The modified Marchenko equation of the inverse scattering problem is obtained and the reconstruction algorithm of the potential function from scattering data of this problem is given.

Keywords: Inverse scattering problem; Sturm-Liouville equation; main equation.

REFERENCES

1. O. Akçay, On the investigation of a discontinuous Sturm-Liouville operator of scattering theory, Math. Commun., Vol:27, 33–45, 2022.
2. Ö. Akçay, Inverse scattering problem for Sturm-Liouville operator with discontinuity conditions on the positive half line, Int. J. Pure Appl. Sci. Vol:7, 401–409, 2021.
3. Kh. R. Mamedov, On an inverse scattering problem for a discontinuous Sturm-Liouville equation with a spectral parameter in the boundary condition, Bound. Value Probl. 171967, DOI 10.1155/2010/171967.
4. V. A. Marchenko, Sturm-Liouville Operators and Applications, AMS Chelsea Publishing, Providence, Rhode Island, 2011.

A Probabilistic Chaotic Image Encryption Scheme

M. Ghebleh¹ A. Kanso¹ M. B. Khuzam²

¹Department of Mathematics, Kuwait University, Kuwait

² Department of Mathematics and Natural Sciences, American University of Iraq, Iraq

mohammad.ghebleh@ku.edu.kw

ali.kanso@ku.edu.kw

mazen.boukhuzam@auis.edu.krd

Abstract

We propose a probabilistic image encryption scheme that improves on existing deterministic schemes by using a chaining mode of chaotic maps in a permutation-masking process. Despite its simplicity, the permutation phase destroys any correlation between adjacent pixel values in a meaningful image. The masking phase modifies the pixel values of the image at hand using pseudorandom numbers with some other initiated random numbers so that any slight change in the plain image spreads throughout the corresponding cipher image. These random numbers ensure the generation of distinct cipher images for the same plain image encryption, even if it is encrypted multiple times with the same key, thereby adding some security features. Simulations show that the proposed scheme is robust to common statistical and security threats. Furthermore, the scheme is shown to be competitive with existing image encryption schemes.

Keywords: Image encryption; chaotic systems; pseudorandom number generators; permutation; diffusion

REFERENCES

*Abstract Submission should be prepared only **1 page**.

Exploring Divergence Measures: Concepts, Applications, and Advances across Disciplines

Servet Akbas¹ and Bilgi Yilmaz²

^{1,2} Department of Mathematics, Kahramanmaraş Sutcu Imam University, Kahramanmaraş, Turkey

[1akbass794@gmail.com](mailto:akbass794@gmail.com) , [2bilgiyilmaz@ksu.edu.tr](mailto:bilgiyilmaz@ksu.edu.tr)

Abstract

Divergence measures serve as crucial statistical instruments for quantifying dissimilarities between probability distributions. This study delves into the realm of divergence measurements, elucidating fundamental principles, diverse applications, and recent advancements. Offering a holistic comprehension of these measures, it underscores their pragmatic utilities across multiple domains. Embracing well-known metrics such as Kullback-Leibler, Jensen-Shannon, and Hellinger divergences alongside their extensions, the research explores their applications within machine learning, information theory, signal processing, and image analysis.

Keywords: Divergence measures; Probability distributions; Information theory; Machine learning

1. REFERENCES

2. Günder, Maurice; PIATKOWSKI, Nico; BAUCKHAGE, Christian. Full Kullback-Leibler-Divergence Loss for Hyperparameter-free Label Distribution Learning. arXiv preprint arXiv:2209.02055, 2022.
3. Pranesh Kumar and S Chhina. A symmetric information divergence measure of the csiszár's f-divergence class and its bounds. Computers & Mathematics with Applications, Vol: 49, NO: 4,575-588, 2005.
4. Syed Mumtaz Ali and Samuel D Silvey. A general class of coefficients of divergence of one distribution from another. Journal of the Royal Statistical Society. Series B (Methodological), 131-142, 1966.
5. Taneje, Inder Jeet. On symmetric and nonsymmetric divergence measures and their generalizations. Advances in Imaging and Electron Physics, 138: 177-250, 2005.

Fixed Points of multiplicative Zamfirescu Mapping in Multiplicative Metric Spaces

Fatih Avşar¹

¹Institute of Science, University of Atatürk, Erzurum, Turkey

fatihavsar7575@gmail.com,

Abstract

In this study, we will examine some fixed point theorems and their applications to Zamfirescu mappings in multiplicative metric spaces. First of all, we will give some basic concepts in the theorems that we will use in the study. Later, we will give preliminary information about some mappings in metric spaces and multiplicative metric spaces. Finally, based on the theorems, we will find the Zamfirescu mappings has a fixed point in the metric spaces and multiplicative metric spaces.

Keywords: Multiplicative metric space, fixed point, metric space, Zamfirescu mappings.

REFERENCES

1. Özavşar, M. ve Çevikel, A. C., 2017. Fixed points of multiplicative contraction mappings on multiplicative metric spaces. Journal of Engineering Tecnology and Applied Sciens 2(2), 65-79
2. He, X., Song, M. ve Chen, D., 2014. Common fixed points for weak commutative mappings on a multiplicative metric spaces. Fixed Point Theory Appl. 2014 (2014), doi:10.1186/1687-1812-2014-48, 9 pages.
3. T. Došenović, M. Postolache, S. Radenović, On multiplicative metric spaces: Survey, Fixed Point Theory Appl., 2016:92, (2016).
4. Abbas, M, Ali, B, Suleiman, Y: Common fixed points of locally contractive mappings in multiplicative metric spaces with applications. Int. J. Math. Math. Sci. 2015, Article ID 218683 (2015)
5. Sarwar, M, Badshah-e-Rome: Some unique fixed point theorems in multiplicative metric space (2014). arXiv:1410.3384v2 [math.GM]

q – Bell Statistical Convergence

Koray İbrahim Atabey¹, Muhammed Recai Türkmen², Mikail Et³, Muhammed Çınar⁴

¹ Muş Nizamülmülk Girl Anatolia İmam Hatip High School, Muş, Turkey

²Department of Mathematics Education, University of Afyon Kocatepe, Afyonkarahisar, Turkey

³Department of Mathematics, University of Firat, Elazig, Turkey

⁴ Department of Mathematics, University of Muş Alparslan, Turkey

korayatabey7@gmail.com,

mrtmath@gmail.com,

mikailet68@gmail.com,

muhammedcinar23@gmail.com

Abstract

In this paper, we use to the q – Bell matrix $\hat{B}_q = \hat{B}_{nk}(q)$ to introduce the notions of q – Bell summability, q – Bell statistical convergence, strongly $\hat{B}_q[p]$ – Cesaro summability and give some inclusion relations about these concepts.

Keywords: q – Bell numbers; q – analog; Statistical Convergences.

REFERENCES

1. Hüseyin Aktuğlu, Şerife Bekar, On q -Cesaro matrix and q -statistical convergence, J. Comput. Appl. Math., 235(16), 4717–4723, 2011.
2. George Eyre Andrews, q -Catalan identities, In The legacy of Alladi Ramakrishnan in the mathematical sciences. New York, NY: Springer New York, 183-190, 2010.
3. Koray İbrahim Atabey, Muhammed Çınar, Mikail Et, q -Fibonacci sequence spaces and related matrix transformations. Journal of Applied Mathematics and Computing, 69, 2135–2154, 2023.
4. Zsófia R. Kereskényiné Balogh, Michael J. Schlosser, q -Stirling numbers of the second kind and q -Bell numbers for graphs. Electronic notes in discrete mathematics, 54, 361-366, 2016.

Mathematical Analysis and Modeling of Biofouling in Urban Water Filtration Systems

Deniz Öztürk¹

¹Department of Mathematics, Sinop University, Sinop, Türkiye

denizozturk@sinop.edu.tr

Abstract

This study aims to introduce mathematical analysis and mathematical modeling strategies to increase the performance of microfiltration and ultrafiltration systems. Initially, we will discuss the biological and engineering background. Next, we address the mathematical analysis and formulations. Finally, the results will be analyzed and evaluated using some computer simulations.

Keywords: Microfiltration, Ultrafiltration; Biofouling; Mathematical Modeling

REFERENCES

1. Iritani, E., Katagiri, N., Takenaka, T., & Yamashita, Y. (2015). Membrane pore blocking during cake formation in constant pressure and constant flux dead-end microfiltration of very dilute colloids. *Chemical Engineering Science*, 122, 465-473.
2. Cogan N., Li J., Badireddy A. R., Chellam S. (2016), Optimal backwashing in dead-end bacterial microfiltration with irreversible attachment mediated by extracellular polymeric substances production, *Journal of Membrane Science*, vol. 520, pp. 337-344
3. Cogan N., Ozturk D., Ishida K., Safarik J. , Chellam S. (2022), Membrane aging effects on water recovery during full-scale potable reuse: Mathematical optimization of backwashing frequency for constant-flux microfiltration, *Separation and Purification Technology*, vol. 286, 120294
4. Chheang, M., Hongprasith, N., Ratanatawanate, C., and Lohwacharin, J. (2022). Effects of Chemical Cleaning on the Ageing of Polyvinylidene Fluoride Microfiltration and Ultrafiltration Membranes Fouled with Organic and Inorganic Matter. *Membranes*, 12(3), 280.

Effective Method for Analyzing Nonlinear Mathematical Model Behavior

Tolga Aktürk¹

¹Department of Mathematics and Science Education, University of Ordu, Ordu, Turkey

tolgaakturk@gmail.com,

Abstract

The exponential function method (MEFM) has been utilized to investigate the behaviors of the Joseph-Egri (TRLW) equation, which is one of the nonlinear mathematical models. This method is known to be an effective technique for obtaining solution functions that support such nonlinear mathematical models. In this study, solution functions satisfying the Joseph-Egri (TRLW) equation have been obtained using the MEFM. To further examine the behaviors of the resulting solutions in detail, computer codes suitable for programming have been written, and two- and three-dimensional as well as sensitivity and contour plots have been obtained.

Keywords: Modified Exponential Function Method (MEFM); Joseph-Egri (TRLW) equation; Behavior of traveling wave solutions.

REFERENCES

1. Taghizadeha, N., Mirzazadeha, M., & Paghaleh, A. S. (2012). Exact travelling wave solutions of Joseph-Egri (TRLW) equation by the extended homogeneous balance method. *Int J Appl Math Comput*, 4(1), 96-104.
2. He, Y., Li, S., & Long, Y. (2012). Exact solutions of the Klein-Gordon equation by modified Exp-function method. In *Int. Math. Forum* (Vol. 7, No. 4, pp. 175-182).
3. Al-Harbi, M., Al-Hamdan, W., & Wazzan, L. (2023). Exact Traveling Wave Solutions to Phi-4 Equation and Joseph-Egri (TRLW) Equation and Calogero-Degasperis (CD) Equation by Modified (G'/G²)-Expansion Method. *Journal of Applied Mathematics and Physics*, 11(7), 2103-2120.

Ostrowski type inequalities via fractional integrals and related results

Funda Türk¹, Samet Erden²

^{1,2}Department of Mathematics, University of Bartın, Turkey

fundaturk44@gmail.com, serden@bartin.edu.tr

Abstract

In this work, we introduced the Ostrowski inequality based on M-fractional integrals. we initially established an identity concerning this inequality to prove the Ostrowski inequality based on M-fractional integrals. We derive some results for the Ostrowski inequality utilizing this identity, different convex of classes of functions, and well-known inequalities.

Keywords: Ostrowski; Convex function; Fractional inequality.

REFERENCES

1. Muhammad, AdilKhan, Sumbel, Begum, Yousaf, Khurshid Yu-Ming, Chu, Ostrowski type inequalities involving conformable fractional integrals, JournalofInequalities and Applications, Vol:70, 1-14, 2018.
2. Alomari, M., Darus, M., Dragomir, S.S., Cerone, P., Ostrowski type inequalities for functions whose derivatives are s-convex in the second sense, Appl. Math. Letter, Vol:23, No:9, 1071–1076, 2010.
3. S. S. Dragomir. A functional generalization of Ostrowski inequality via Montgomery identity. Acta Math. Univ. Comenian (N.S.), Vol:84, No:1, 63–78, 2015.
4. G. A. Anastassiou, M-fractional integral inequalities, J. Comput. Anal. Appl, Vol:29, No:6, 1153-1158, 2021.
5. G. A. Anastassiou, Fractional Left Local General M-Derivative. Intelligent Analysis: Fractional Inequalities and Approximations Expanded, 01-510, 2020
6. G. A. Anastassiou, Fractional Right Local General M-Derivative. Intelligent Analysis: Fractional Inequalities and Approximations Expanded, 511-520, 2020.
7. G. A. Anastassiou, Ostrowski type inequalities, Proc. AMS, Vol:123, 3775-3781, 1995.
8. Chu, Y.-M., Zhang, X.-M., Wang, G.-D.: The Schur geometrical convexity of the extended mean values. J. Convex Anal. Vol:15, No:4, 707–718, 2008.

PERSONALITY ANALYSIS USING ARTIFICIAL INTELLIGENCE ACCORDING TO THE EYE DESCRIPTIONS IN MARIFETNÂME

Semra Çelebi¹, İbrahim Türkoğlu²

¹ Firat University, Faculty of Technology, Software Engineering Department, Elazig, Turkey

² Firat University, Faculty of Technology, Software Engineering Department, Elazig, Turkey

scelebi@firat.edu.tr,

iturkoglu@firat.edu.tr

Abstract

Body language, facial structure and general appearance can give many clues about a person's character at first impression. Therefore, people often form their prejudices about a person by basing their first impressions on these clues. The relationship between certain physical characteristics and character and intelligence has been remarkable historically as well as in modern science. In Old Turkish Literature, there are works aimed at character analysis from human physical characteristics. One of these works is Marifetnâme, written by the famous scientist and clergyman İbrahim Hakkı of Erzurum in 1756. In the section of Marifetname called "mirror of bodies", a connection is established between the human body and facial structures and the spiritual aspect. In this study, this connection written by İbrahim Hakkı was transformed into practice with today's technologies and artificial intelligence was used to carry out the character analysis of İbrahim Hakkı. For the decision process, training was performed with three different deep learning-based CNN models (AlexNet, ResNet and YOLOv8) and the results were compared. The proposed application process is based on three steps: In the first step, from each image in the dataset, first the face regions and then the areas where the eyes are located are cropped with a facial landmark detection algorithm. In the second step, an artificial intelligence model was determined to predict the eye structure in each frame. The third step aims to detect in real time the character trait associated with the eye structure determined using this model in Marifetname. As a result of this study, character analysis matched with 96% accuracy according to the developed artificial intelligence-based application process.

Keywords: İbrahim Hakkı; Marifetnâme; Artificial intelligence; Character analysis; Personality.

Exponential Inequalities Involving Riemann-Liouville Fractional Integral

Funda Türk¹, Samet Erden², Burçin Gökkurt Özdemir³

^{1,2}Department of Mathematics, University of Bartın, Turkey

³Department of Mathematics and Science Education, Faculty of Education, University of Bartın, Turkey

fundaturk44@gmail.com, erdensmt@gmail.com, gokkurtburcin@gmail.com

Abstract

New inequalities related to Riemann-Liouville Fractional integrals for exponential functions are provided. Also, some special cases of these results are examined.

Keywords: Riemann-Liouville Fractional Integral, Ostrowski Inequality, Exponential

REFERENCES

1. Cerone, P.; Dragomir, S.S., & Roumeliotis, Some Ostrowski type inequalities for n-time differentiable mappings and applications, *Demonstratio Math*, Vol:32, No:4, 697-712.
2. Sarikaya, M. Z., On the Ostrowski type integral inequality, *Acta Math. Univ. Comenianae*, Vol: LXXIX, No:1, 129-134, 2010.
3. Erden, S., Budak, H., Sarikaya, M. Z., Iftikhar, S., & Kumam, P., Fractional Ostrowski type inequalities for bounded functions, *Journal of Inequalities and Applications*, Vol:123, 1-11, 2020.
4. Budak, H. Sarikaya, M. Z., & Erden, S., New weighted Ostrowski type inequalities for mappings whose nth derivatives are of bounded variation, *International J. of Analysis and App*, Vol:12, No:1, 71-79, 2016.
5. Cerone, P.; Dragomir, S.S., & Roumeliotis, J., An inequality of Ostrowski type for mappings whose second derivatives are bounded and applications, *RGMIA Research Report Collection*, Vol:1, No:1, 35-42, 1998.
6. Dragomir, S. S., Cerone, P., & Roumeliotis, J., A new generalization of Ostrowski's integral inequality for mappings whose derivatives are bounded and applications in numerical integration and for special means, *RGMIA Research Report Collection*, Vol:2, No:1, 105-111, 1999.
7. Wang M. & Zhao, X., Ostrowski type inequalities for higher-order derivatives, *Journal of Inequalities and Applications*, Article ID 162689, 1-8, 2009.

CONFORMABLE STURM-LIOUVILLE PROBLEM WITH TWO PARAMETER

Aslı Öner¹, Sertac Goktas², Büşra Barut³

^{1,2,3} Department of Mathematics, Mersin University, Mersin, Turkey

aslioner98@gmail.com, srtcgoktas@gmail.com, bbarut0633@gmail.com

Abstract

In this study, we used conformable derivatives to define the Sturm-Liouville problem with two parameters and examined various spectral properties associated with them. First of these properties, the Sturm-Liouville problem with two parameters was reduced to the simpler one-parameter problem. This reduction provides an important perspective for better understanding the fundamental structures of the problems. Additionally, we focused on the orthogonality properties of eigenfunctions. These properties play a critical role in understanding the behavior and relationships of solutions. Investigating the reality of solutions is important to understand the physical relevance and practical usability of the considered eigenvalue problem. This analysis provides a deeper understanding of the system's behavior and potential applications. Finally, we examined integral relations, which are used to explain important connections and relationships between different aspects of the system.

References [1-5] form the basis of our study, and quality and effective results have been carried out from these studies.

Keywords: Conformable Derivative, Eigenvalue Problems, Sturm-Liouville Problems.

REFERENCES

1. Douglas R. Anderson and Darin J. Ulness, Newly Defined Conformable Derivatives. *Advances in Dynamical Systems and Application*, No: 10, Vol:2, 109-137, 2015.
2. Felix M. Arscott, Two-parameter Eigenvalue Problems in Differential Equation, *Proceedings of the London Mathematical Society*, Vol: 3, No: 3, 459-470, 1964.
3. Frederick V. Atkinson, Angelo B. Mingarelli, Multiparameter Eigenvalue Problems. Vol: 1. New York: Academic Press, 1972.
4. Roshdi Khalil, Mohammed Al Horani, A. Yousef and Mohammed Sababheh, A New Definition of Fractional Derivative, *Journal of Computational and Applied Mathematics*, Vol: 264, 65-70, 2014.
5. Thabet Abdeljawad, On Conformable Fractional Calculus, *Journal of Computational and Applied Mathematics*, Vol: 279, 57-66, 2015.

q –Pell Sequence Spaces

Koray İbrahim Atabey¹, Murat Karakaş²

¹ Muş Nizamülmülk Girl Anatolia İmam Hatip High School, Muş, Turkey

²Department of Mathematics, University of Bitlis Eren, Bitlis, Turkey

korayatabey7@gmail.com,

mkrks33@gmail.com,

Abstract

In this study, we build q -analogue of the q -Pell matrix $\tilde{P}_q = (\tilde{P}_{nk}(q))$ defined by

$$\tilde{P}_q = (\tilde{P}_{nk}(q)) = \begin{cases} \frac{q^k P_{k-1}(q)}{P_n(q) - P_{n-1}(q)}, & n-1 \leq k \leq n \\ 0, & \text{otherwise} \end{cases} \quad n, k = \{2, 3, \dots\}.$$

After, we use this analogue to define the sequence spaces $c(\tilde{P}_q)$, $c_0(\tilde{P}_q)$, $\ell_\infty(\tilde{P}_q)$, $\ell_p(\tilde{P}_q)$ ($1 \leq p < \infty$). Then, we provide some inclusion relations for these spaces and examine a few topological characteristics. Furthermore, we construct a basis for the space $\ell_p(\tilde{P}_q)$, calculate α -, β -, γ -duals of the same space, describe certain matrix classes and look at some geometric properties.

Keywords: Pell numbers; q –Pell numbers; q –analogue; Dual Spaces; Matrix Transformations; Banach-Saks Property.

REFERENCES

1. Hüseyin Aktuğlu, Şerife Bekar, On q -Cesaro matrix and q -statistical convergence, J. Comput. Appl. Math., 235(16), 4717–4723, 2011.
2. George Eyre Andrews, q -Catalan identities, In The legacy of Alladi Ramakrishnan in the mathematical sciences. New York, NY: Springer New York, 183-190, 2010.
3. Koray İbrahim Atabey, Muhammed Çınar, Mikail Et, q -Fibonacci sequence spaces and related matrix transformations. Journal of Applied Mathematics and Computing, 69, 2135–2154, 2023.
4. Toufik Mansour, Mark Shattuck, Restricted partitions and q –Pell numbers. Open Mathematics, 9(2), 346-355, 2011.

ON ρ -STATISTICAL CONVERGENCE DEFINED BY MODULAR SEQUENCE SPACES OF ORDER α

Gülcan Atıcı Turan¹

¹ Vocational School of Tunceli, Munzur University, Tunceli, Turkey

gatici23@hotmail.com,

Abstract

In this paper, we introduce Wijsman ρ -statistical convergence of order α and Wijsman strongly ρ -convergence of order α and define $[W_\rho^\alpha, \mathcal{M}, v, p]$ by using a sequence of Orlicz functions. Also, some inclusion theorems are presented.

Keywords: Wijsman convergence; statistical convergence; Orlicz function.

REFERENCES

1. Fatih Nuray and Billy Eugene Rhoades, Statistical convergence of sequences of sets, Fasc. Math., Vol: 49, 87–99, 2012.
2. Uğur Ulusu and Erdinç Dünder, I-Lacunary Statistical Convergence of Sequences Of Sets, Filomat, Vol:28, No:8, 1567–1574, 2014.
3. Uğur Ulusu and Fatih Nuray, Lacunary statistical convergence of sequence of sets, Prog. Appl. Math., Vol:4, No:2, 99–109, 2012.
4. Isaac Jacob Schoenberg, The Integrability of Certain Functions and Related Summability Methods, Amer. Math. Monthly, Vol:66, No:5, 361–375, 1959.
5. Hugo Steinhaus, Sur La Convergence Ordinaire Et La Convergence Asymptotique, Colloq. Math., Vol:2, 73–74, 1951.
6. Joram Lindenstrauss and Lior Tzafriri, On Orlicz sequence spaces, Israel Journal of Mathematics, Vol:10, 379–390, 1971.
7. Karl Lindberg, On Subspaces of Orlicz Sequence Spaces, Studia Math., Vol:45, No:2, 119–146, 1973.
8. Nazlım Deniz Aral, Hacer Şengül Kandemir, Mikail Et, On P-Statistical Convergence of Order α of Sequences of Sets, Miskolc Mathematical Notes, Vol:24, No:2, 569–578, 2023.

Developing High-Efficiency Organic Solar Cells through Molecular Design Analysis of Novel D-A-Di-A-D Conjugated Compounds

Imane EL Mhamedi¹, Zakaria El Malki¹

¹Moulay Ismail University, High School of Technology, (ESTM), Modeling, Materials and Systems Control (MMSC), Computer Engineering and Intelligent Electrical Systems (2ISEI).) BP: 3103, Toulal, Meknes, Morocco.

i.elmhamedi@edu.umi.ac.ma, zelmalki@yahoo.fr

Abstract

In this investigation, we employed the DFT/TD-DFT/B3LYP/6-31G(d,p) method to analyze the structural, optoelectronic, and optical properties of a series of conjugated compounds characterized by a modular D-A-Di-A-D architecture. These compounds feature a D donor unit (carbazole), an A acceptor unit (benzothiadiazole), and various Di donor units. Utilizing AMPS-1D, we scrutinized the photovoltaic properties of these compounds when combined with the PCBM acceptor, dividing our analysis into three distinct phases. Initially, we assessed the energy conversion efficiency of the compounds, observing performances ranging from 7.11% to 11.70%. Introducing a PEDOT layer between the active layer and the anode in the second phase resulted in a significant enhancement in photovoltaic performance, achieving energy conversion efficiencies of up to 15.31%, the highest recorded in our study. Lastly, incorporating ZnO as an intermediate interface layer in the third phase notably improved photovoltaic performance across all compounds, with energy conversion efficiency values reaching 17.13%, 17.20%, and 18%. These findings underscore the effectiveness of ZnO addition in enhancing photovoltaic conversion, highlighting its potential for organic solar cell applications. Overall, our results suggest that these compounds hold promise as viable candidates for bulk heterojunction organic solar cell applications.

Keywords: Organic compounds; BHJ; Efficiency; PEDOT; ZnO

A NEW FRACTIONAL MODELLING OF RC ELECTRIC CIRCUIT

Enes Ata¹, İ. Onur Kıymaz², Hacı Mehmet Başkonuş³

^{1,2} Department of Mathematics, Kirsehir Ahi Evran University, Kirsehir, Turkey

³ Department of Mathematics and Science Education, Harran University, Sanliurfa, Turkey

enesata.tr@gmail.com, iokiymaz@ahievran.edu.tr, hmbaskonus@gmail.com

Abstract

In this study, we introduce a new fractional RC electric circuit model with the generalized Caputo fractional derivative and obtain analytical solutions via the generalized Laplace and inverse Laplace transforms. We also compare the approximate behavior of the solution of the new fractional RC electric circuit model with the approximate behavior of the solutions of RC electric circuit models defined by the Caputo, Caputo-Fabrizio fractional derivatives and conformable operator, which can be found in the literature.

Keywords: Fractional derivatives; Electric circuits; Mathematical modellings.

REFERENCES

1. Lokenath Debnath, Dambaru Bhatta, Integral Transforms and Their Applications, CRC Press, 2015.
2. Anatoli Aleksandrovich Kilbas, Hari Mohan Srivastava, Juan Jose Trujillo, Theory and Applications of Fractional Differential Equations, Elsevier, 2006.
3. Igor Podlubny, Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications, Elsevier, 1998.
4. Bo Zhang, Xujian Shu, Fractional-Order Electrical Circuit Theory, Springer, 2022.
5. John Bird, Electrical Circuit Theory and Technology, Routledge, 2017.

ON WAVE STRUCTURES OF TIME CONFORMABLE ZAKHAROV–KUZNETSOV EQUATION

Meltem Uzun¹

¹Department of Aeronautical Engineering, Istanbul Gelisim University, Istanbul, Turkey

meuzun@gelisim.edu.tr,

Abstract

This study reveals the soliton wave solutions of the time conformable Zakharov–Kuznetsov equation, which models complex wave phenomena in different contexts, such as surface water waves, ion-acoustic waves in plasma, and some optical models. A robust solution technique, Sardar sub-equation method is utilised to obtain solutions of the equation with dual power law nonlinearity. The physical implications of the obtained results are illustrated by symbolic computation tools.

Keywords: Zakharov–Kuznetsov equation; Sardar sub-equation method; Dual-power law nonlinearity.

REFERENCES

1. R. Khalil, M. Al Horani, A. Yousef , M. Sababheh, A New Definition of Fractional Derivative, Journal of Computational and Applied Mathematics, Vol:264, , 65–70, 2014.
2. Y. Saliou, S. Abbagari, A. Houwe, W-Shape Bright And Several Other Solutions to the (3+1)-Dimensional Nonlinear Evolution Equations, Modern Physics Letters B, 2150468 (23 pages), 2021.
3. B. T. Matebese, A. R. Adem, C. M. Khalique, A. Biswas, Solutions of Zakharov–Kuznetsov Equation with Power Law Nonlinearity in (1+3) Dimensions, Physics of Wave Phenomena, Vol. 19, No. 2, pp. 148–154, 2011.

ON USING A NEW APPROACH TO DETERMINE THE ROOT OF NONLINEAR EQUATIONS

Yasemin Bakır^{1*}, Oya Mert², Gülay Karahanlı³

^{1*}Department of Management Information System, University of Fenerbahçe, İstanbul, Türkiye

yasemin.bakir@fbu.edu.tr,

²Department of Mathematics, University of Tekirdağ Namık Kemal, Tekirdağ, Türkiye

oyamert@nku.edu.tr,

³Department of Information Technology, University of Tekirdağ Namık Kemal, Tekirdağ, Türkiye

gulaykarahanli@nku.edu.tr,

Abstract

This article presents a recently developed method for solving nonlinear equations. Various iterative methods can be employed to solve nonlinear equations, including the Bisection method, the Newton-Raphson method, the Secant method, the Regula Falsi method, and others. The objective is to identify the roots of nonlinear equations by utilizing polynomial curves obtained through Lagrange interpolation under specified initial conditions, without resorting to any derivative processes. The efficacy of the proposed method has been demonstrated through the examination of well-known examples from the literature.

Keywords: Lagrange approximation; Nonlinear equation; Root finding

REFERENCES

1. Cleonice F. Bracciali, Michael Carley, Quasi-analytical root-finding for non-polynomial functions, Numerical Algorithms, Vol: 76, No: 3, 639–653, 2017.
2. Stephen J. Garrett, Newton-Raphson Method, Introduction to Actuarial and Financial Mathematical Methods, 2015.
3. Amir Naseem, M. A. Rehman, Thabet Abdeljawad, A Novel Root-Finding Algorithm With Engineering Applications and Its Dynamics via Computer Technology, IEEE Access, 10, 19677-19684, 2022.

Modeling Epidemics Using Ising Model and Voronoi Tessellation: A Novel Study and Epidemiological Applications

Şeyma Firdevs Korkmaz¹, Hasan Bulut¹, Gülnur Yel²

¹Department of Mathematics, Firat University, Elazığ, Turkey

² Faculty of Educational Sciences, Final International University, Kyrenia, Mersin 10, Turkey

sfkorkmaz@firat.edu.tr

hbulut@firat.edu.tr

gulnuryel33@gmail.com

Abstract

This article aims to investigate the spread of diseases by drawing inspiration from the Ising model in physics. The Ising model, known for simulating magnetic properties, is introduced into the field of disease epidemiology as a novel application area. While Voronoi tessellation provides location data representing individuals' health statuses, the Ising model is employed to model interactions among neighbors. A modified Ising model will be developed to analyze the speed and impact of disease spread, to be examined through Monte Carlo simulations and the Metropolis algorithm. The goal of this article is to assess the potential use of the Ising model in disease epidemiology, contributing to a better understanding of disease spread processes and the development of effective control strategies.

Keywords: Voronoi Tessellation; Ising Model; Monte Carlo Simulation; Metropolis Algorithm; Disease Spread.

REFERENCES

1. Mata, A.S., Dourado, S. M. P., (2021). Mathematical modeling applied to epidemics: an overview, São Paulo Journal of Mathematical Sciences, 15, 1025-1044. DOI 10.1007/s40863-021-00268-7.
2. Anderson R. M., May R. M. (1991). Infectious diseases of humans. Oxford: Oxford University Press.
3. N. Walker, K.M. Tam, M. Jarrell, Deep learning on the 2-dimensional Ising model to extract the crossover region with a variational autoencoder, Sci. Rep. 10 (2020) 1–12.
4. H. Imai, M. Iri, K. Murota, Voronoi Diagram in the Laguerre Geometry and Its Applications., SIAM J. Comput. 14 (1985) 93–105. <https://doi.org/10.1137/0214006>.

Designing a novel radial basis process for the nonlinear prey-predator system

Zulqurnain Sabir^{1,2}, Ayse Nur Akkilic³, Hasan Bulut^{3,4}

¹ Department of Computer Science and Mathematics, Lebanese American University, Beirut, Lebanon

² Department of Mathematics and Statistics, Hazara University, Mansehra, Pakistan

³ Department of Mathematics, Firat University, Elazığ, Turkey

⁴ Azerbaijan University, Baku, Azerbaijan

Email: zulqurnain_maths@hu.edu.pk, akkilicaysenur@gmail.com, hbulut@firat.edu.tr

Abstract: In this research, a novel deep neural network process (DNNP) is proposed for solving one of the dynamical prey-predator nonlinear systems (PPNS). The mathematical PPNS is based on two categories, one is prey and other is a predator. The proposed DNNP contains three hidden layers with 15, 30 and 45 numbers of neurons, while the log-sigmoid activation function is used in each hidden layer in order to solve the dynamical PPNS. The optimization is performed through the scale conjugate gradient procedure for solving the dynamical PPNS. An Adam method is used to get the dataset, which is used to train the process through scale conjugate gradient by dividing the data into training as 74%, while 13% for both validation and testing. The correctness of the DNNP using the optimization of scale conjugate gradient is observed by using the overlapping of the proposed and reference solutions. The accuracy of the scheme is also observed through the small absolute error values and correlation coefficient values that are obtained as 1 for solving the model. Moreover, the values of the state transitions, error histogram and fitness function performances approve the accuracy of the proposed DNNP using the optimization of scale conjugate gradient for solving the dynamical PPNS.

Keywords: Dynamical prey-predator, Deep neural network; Scale conjugate gradient; Log-sigmoid; Hidden layers.

References

- [1] Sabir, Z., Botmart, T., Raja, M.A.Z. and Weera, W., 2022. An advanced computing scheme for the numerical investigations of an infection-based fractional-order nonlinear prey-predator system. Plos one, 17(3), p.e0265064.
- [2] Alkaabi, H., Alkarbi, N., Almemari, N., Said, S.B. and Sabir, Z., 2024. Gudermannian neural network procedure for the nonlinear prey-predator dynamical system. Heliyon.
- [3] Umar, M., Sabir, Z., Raja, M.A.Z., Amin, F., Saeed, T. and Sanchez, Y.G., 2023. Design of intelligent computing solver with Morlet wavelet neural networks for nonlinear predator-prey model. Applied Soft Computing, 134, p.109975.
- [4] Ruttanaprommarin, N., Sabir, Z., Núñez, R.A.S., Az-Zobi, E., Weera, W., Botmart, T. and Zamart, C., 2023. A stochastic framework for solving the prey-predator delay differential model of holling type-III. CMC-Comput. Mater. Cont., 74(3), pp.5915-5930.

LIE ALGEBRA AND SOME GEODESIC PROPERTIES

Mevlüt Ağar¹, Mustafa Yeneroğlu¹

¹Department of Mathematics, University of Firat, Elazığ, Turkey

sha.mvlt07@gmail.com, mustafayeneroglu@gmail.com

Abstract

Geodesy is a word of Greek origin. Geo=earth, world; dezi=means to divide, measure. Geodesy is the science of measuring and projection of the earth's surface. we tried to define Lie groups and Lie algebra based on some definitions about groups and algebra in real numbers. Then, we talked about some Geodesic features and tried to analyze this Geodesy mathematically first. Later, we tried to work on their adaptation to space geometry. For example; we related this to Euler's formula by applying it to the rotational ellipsoid. On the other hand, with the help of Frenet trihedron, we defined tangent and normal unit vectors on the surfaces and obtained curvature and curvature radii accordingly. The reason for all this work was to obtain the Geodesy curve and equation. Therefore; with the help of these studies, we tried to obtain the curvature of a curve and its equation on any surface in space. As a matter of fact, when we adapted this to the earth ellipsoid, we obtained Geodesic equations under the name of Geodesic curvature. Then, using this Geodesic curvature, we obtained the Geodesic torsion and its equation. In conclusion; we have concluded that the shortest curve between any two points on a surface in space is the Geodetic curve.

In this study, we took into consideration the previous studies and tried to find definitions and equations related to different Lie groups and Lie algebras and obtain Geodesic curvature, Geodesic equation and Geodesic torsion.

Keywords: Lie Group, Lie Algebra, Geodesic Curvature, Geodesic Equation, Geodesic Torsion.

REFERENCES

1. Akkaş, S., Hacısalihoğlu, H., H., Özel, Z., Sabuncuoğlu, A., Soyut Matematik, Ankara, 1998.
2. Takushiro, O., Tsunero, T. "TheGroup of Isometries a LeftInvaryantRiemanMetricon a Lie Group", Osaka University, Toyonaka, Japan, 1976.
3. Warner, W., F., "Foundations of DifferentiableManifoldsand Lie Groups", Universty of Pennsylvania, London, 1971.
4. Iqbal, M.A., Ali, A., Alshammari, I., and Ozel, C., Construction of new Lie group and its geometric properties. AIMS Mathematics, 9(3), 2023.
5. Prada, E., Miková, L., Virgala, I., Kelemen, C., Sincák, P.J., Mykhailyshyn, R., Mathematical Modeling of Robotic Locomotion Systems. Symmetry, 16, 2024.

CURVATURES COMPUTATION FOR CURVES IN AFFINE SPACE USING FRACTIONAL CALCULUS

Meltem Ogrenmis¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

mogrenmis@firat.edu.tr,

Abstract

This paper proposes a method to compute the curvatures of equiaffine curves in affine space using local fractional derivatives. It introduces the concepts of α -equiaffine arc length and α -equiaffine curvatures through a generalized local approach involving conformable, V-derivative, and similar methods. Equiaffine Frenet formulas and curvatures are reestablished within fractional calculus.

Keywords: Affine space; Fractional Calculus; Curvatures.

REFERENCES

1. M.E. Aydin, A. Mihai, A. Yokus, Applications of Fractional Calculus in Equiaffine Geometry: Plane Curves With Fractional Order, *Mathematical Methods in the Applied Sciences*, 44 (17), 13659-13669, 2020.
2. M.E. Aydin, M. Bektas, A.O. Ogrenmis, A. Yokus, Differential Geometry of Curves in Euclidean 3-Space With Fractional Order, *International Electronic Journal of Geometry*, 14(1), 132-144, 2021.
3. K. Lazopoulos, A. K. Lazopoulos, Fractional Differential Geometry of Curves and Surfaces, *Progress in Fractional Differentiation and Applications*, 2(3), 169-186, 2016.
4. V. E. Tarasov, On Chain Rule for Fractional Derivatives, *Communications in Nonlinear Science and Numerical Simulation*, 30(1), 1-4, 2016.
5. J. Vanterler da C. Sousa, E. Capelas de Oliveira, A New Truncated M-Fractional Derivative Type Unifying Some Fractional Derivative Types With Classical Properties, *International Journal of Analysis and Applications*, 16(1), 83-96, 2018.
6. J. Vanterler da C. Sousa, E. Capelas de Oliveira, Mittag-Leffler Functions and the Truncated V-Fractional Derivative, *Mediterranean Journal of Mathematics*, 244(14), 2017.
7. D. Davis, Generic Affine Differential Geometry of Curves in R_n , *Proceedings of the Royal Society of Edinburgh Section A*, 136, 1195-1205, 2006.

EXPANDING FRACTIONAL EQUIAFFINE CURVATURES OF PLANE CURVES

Meltem Ogrenmis¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

mogrenmis@firat.edu.tr,

Abstract

This study focuses on equiaffine plane curves, employing a generalized local fractional derivative. It introduces the concepts of α -equiaffine arc length and curvature. Equiaffine Frenet formulas and an equivalent of the fundamental theorem are reintroduced within fractional calculus. Additionally, the study describes plane curves with constant fractional equiaffine curvature.

Keywords: Affine space; Fractional Calculus; Curvatures.

REFERENCES

1. S.G. Samko, A.A. Kilbas, OI Marichev, Fractional Integrals and Derivatives. Theory and Applications, Gordon and Breach Science Publishers, London, UK, 1993.
2. M.E. Aydin, Effect of local fractional derivatives on Riemann curvature tensor. Exam. and Counter. 5, 100134, 2024.
3. M.E. Aydin, A. Mihai, A. Yokus, Applications of Fractional Calculus in Equiaffine Geometry: Plane Curves With Fractional Order, Mathematical Methods in the Applied Sciences, 44 (17), 13659-13669, 2020.
4. T. Yajima, Nagahama H, Differential geometry of viscoelastic models with fractional order derivatives, Journal of Physics A: Mathematical and Theoretical, 43(38),385207, 2010.
5. V.E. Tarasov, On Chain Rule for Fractional Derivatives, Communications in Nonlinear Science and Numerical Simulation, 30(1), 1-4, 2016.
6. J. Vanterler da C. Sousa, E. Capelas de Oliveira, Mittag-Leffler Functions and the Truncated V-Fractional Derivative, Mediterranean Journal of Mathematics, 244(14), 2017.
7. H.W. Guggenheimer, Differential Geometry, McGraw-Hill, New York, 1963.

Several Integral Representations of the ${}_pH_k$ Srivastava's triple hypergeometric functions

Ali Olgun¹, Zekiye Rana Lüsna¹ and Oğuz Yağcı¹

¹Department of Mathematics, University of Kırıkkale, Kırıkkale, Turkey

aliolgun71@gmail.com.tr, zekiyelusna@gmail.com.tr,

oguzyagci26@gmail.com.tr,

Abstract

This paper introduces newly ${}_pH_{A,k}$, ${}_pH_{B,k}$, and ${}_pH_{C,k}$ Srivastava's triple hypergeometric function using ${}_pH_k$ Pochhammer symbol. We also present relationship between ${}_pH_k$ Srivastava's triple hypergeometric functions and classical Srivastava's triple hypergeometric functions. Then, we obtain some properties of the ${}_pH_k$ Srivastava's triple hypergeometric functions such as integral representations and recurrence formulas.

Keywords: Srivastava hypergeometric function; Integral representations; Derivative formula; Recurrence relations

REFERENCES

1. Kuldeep Singh Goglot, Two parameter gamma function and its properties, *arXiv preprint arXiv:1701.01052*, 2017.
2. Shadid Mubeen,, Sana Iqbal,, Gauhar Rahman, Contiguous function relations and an integral representation for Appell k-series $F_{1,k}$, *International Journal of Mathematical Research*, Vol: 4, No:2, 53-63, 2015.
3. Shadid Mubeen, Habibullah, An integral representation of some k-hypergeometric functions, *Int. Math. Forum*, Vol:7, No:4, 203-207, 2012.
4. İsmail Onur Kıymaz, Ayşegül Çetinkaya, Praveen Agarwal, A study on the k-generalizations of some known functions and fractional operators, *Journal of Inequalities and Special Functions*, Vol:8, No:4, 31-41, 2017.

Fractional Solutions of the General Class of Non-Fuchsian Differential Equations

Mehmet Aydin¹, Resat Yilmazer¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

maydin2185@gmail.com, ryilmazer@firat.edu.tr,

Abstract

In this article, we consider the non-Fuchsian differential equation, which is a general class of second-order differential equations with singular points, such as Fukuhara, Tricomi, Whittaker, Bessel, whose solutions are obtained by different methods in the literature. Then, we obtain new fractional solutions with the help of the fractional operator. This operator is applied to homogeneous and non-homogeneous linear ordinary differential equations. Thus, we obtain new solutions in fractional forms with a newly developed method.

Keywords: Fractional calculus; Non-Fuchsian equations; Differential equations.

REFERENCES

1. Podlubny, I. Fractional Differential Equations; Academic Press: New York, NY, USA, 1999.
2. Lin, S.D.; Tu, S.T.; Srivastava, H.M. A Unified Presentation of Certain Families of Non-Fuchsian Differential Equations via Fractional Calculus Operators. Comput. Math. Appl. 45, 1861–1870, 2003.
3. Yilmazer, R. N-fractional calculus operator N_m method to a modified hydrogen atom equation. Math. Commun. 15, 489–501, 2010.
4. Yilmazer, R.; Inc, M.; Bayram, M. On Discrete Fractional Solutions of Non-Fuchsian Differential Equations, Mathematics, 6 (12), 308, 2018.

Weighted Statistical Convergence in Probability

Kübra Elif AKBAŞ¹ and Mahmut IŞIK²

¹Faculty of Medicine, Fırat University, Elazığ, Turkey; kelifakbas@gmail.com

²Faculty of Education, Harran University, Şanlıurfa, Turkey; misik63@yahoo.com

Abstract. In this study, we introduce the concept of weighted statistical convergence in probability and also give some inclusion relations about weighted statistical convergence.

Keywords: Density, Statistical convergence, Sequence of random variables, Weighted statistical convergence.

REFERENCES

- [1] Akbaş, KE, Işık, M. On asymptotically λ -statistical equivalent sequences of order α in probability, Filomat 34(13) (2020), 4359--4365.172.
- [2] H. Fast, Sur la convergence statistique, Colloq. Math. 2 (1951) 241-244.
- [3] JA. Fridy, On statistical convergence, Analysis 5 (1985), no. 4, 301--313.
- [4] Das, P. ; Ghosal, S. and Som, S. Statistical convergence of order α in probability, Arab J. Math. Sci. 21(2) (2015), 253—265.
- [5] Ghosal, S. Weighted statistical convergence of order α and its applications, J. Egyptian Math. Soc. 24(1) (2016), 60--67.
- [6] Ghosal, S. λ -convergence of a sequence of random variables, J. Egyptian Math. Soc. 23(1) (2015), 85--89.
- [7] T. Salat, On statistically convergent sequences of real numbers, Math. Slovaca 30 (1980), 139-150.
- [8] IJ. Schoenberg, The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- [9] H. Steinhaus, Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951), 73-74.

THE NEW GOMPERTZ DISTRIBUTION

Ayşe METİN KARAKAŞ¹ Sinan ÇALIK¹

¹Department of Statistics, University of Bitlis Eren, Bitlis, Turkey

²Department of Statistics, University of Firat, Elazığ, Turkey

aysekarakas5767@gmail.com,

Abstract

This research introduces a new four parameter Marshall Olkin Power Gompertz distribution model. The novel model has many sub-models that are very useful in modeling real-life data such as extended Gompertz distribution. The proposed distribution was compared with its sub-models and other existing models

Keywords: Gompertz Distribution, Marshall Olkin Distribution.

REFERENCES

1. Krishna, E., Jose, K. K., Alice, T., & Ristić, M. M. (2013). The marshall-olkin Fréchet distribution. Communications in Statistics-Theory and Methods, 42(22), 4091-4107.

ANALYSIS OF MATHEMATICAL MODEL WAVE SOLUTIONS WITH THE EXPONENTIAL FUNCTION METHOD

Elif Nur YILDIRIM¹, Münevver TUZ²

^{1,2} Department of Mathematics, Faculty of Science, Firat University, Elazig, Turkey

elfyldrm335@gmail.com, mtuz@firat.edu.tr

Abstract

In this study, the analysis of the exponential function method, which is used to obtain traveling wave solutions of partial differential equations and has many applications in the literature, will be included. In addition, the method was used to find the exact solution of the Gibbon equation and several solutions depending on arbitrary parameters were obtained. It has gained validity as an easy, simple and effective method for nonlinear equations that arise in mathematical physics.

Keywords: Exponential function method; exact solution; Gibbon equation; nonlinear wave equations

REFERENCES

- [1] He, J.H., Wu, X. H. (2006). Exp-function method for nonlinear wave equations, Chaos, Solitons and Fractals, 30, 700-708.
- [2] Zhao, M. M., Li, C. (2008). The-expansion method applied to nonlinear evolution equations, Science paper Online, 21789.
- [3] Hafez, M. G., Akbar, M. A. (2015). An exponential expansion method and its application to the strain wave equation in microstructured solids, Ain Shams Engineering Journal, 6(2), 683-690.
- [4] Islam, R., Alam, M. N., Hossain, A.K.M.K.S., Roshid, H. O. Akbar, M. A. (2013). Traveling wave solutions of nonlinear evolution equations via $\text{Exp}(-\Phi(\eta))$ -expansion method". Global Journal of Science Frontier Research, 13(11), 63-71.

PREDICTING STUDENT PERFORMANCE USING STATISTICAL LEARNING TECHNIQUES

Gökhan GÖKDERE¹, Melih TAŞ¹

¹Department of Statistics, University of Firat, Elazig, Turkey

ggokdere@firat.edu.tr

Abstract

Universities accumulate large amounts of student data electronically. Filtering a data according to certain criteria, based on the information stored in the database, becomes difficult when executed manually. Therefore, it is very important to implement tools that analyze data in statistical, descriptive or computational ways. The objectives of this study are listed below:

- To examine and identify the variables used to analyze student performance.
- To examine existing prediction methods for predicting student performance.
- To select a student dataset from the Kaggle website dataset repository ("https://www.kaggle.com/datasets"). Apply machine learning algorithms for classification and prediction on the dataset. Analyze and compare the performance of these algorithms.

Keywords: Machine learning algorithms; Student performance; Kaggle.

REFERENCES

1. Hou, R., Huang, H., Zeng, D., Xia, G., Ghany, K. K. A., & Zawbaa, H. M. (Eds.). (2023). *Big Data Technologies and Applications: 11th and 12th EAI International Conference, BDTA 2021 and BDTA 2022, Virtual Event, December 2021 and 2022, Proceedings* (Vol. 480). Springer Nature.
2. Romero, C., & Ventura, S. (2020). Educational data mining and learning analytics: An updated survey. *Wiley interdisciplinary reviews: Data mining and knowledge discovery*, 10(3), e1355.
3. Khan, A., & Ghosh, S. K. (2021). Student performance analysis and prediction in classroom learning: A review of educational data mining studies. *Education and information technologies*, 26, 205-240.
4. De Giorgi, M. G., Donateo, T., Ficarella, A., Menga, N., Chiodo, L. S., & Strafella, L. (2024). Coupling Principal Component Analysis-Based Sensor Data Reduction Techniques and Multi-net systems for Simultaneous Prediction of Multi-Component Degradation Levels in Hybrid Electric Rotorcraft Engines. *Measurement*, 114212.

ON THE INVERSE PROBLEM FOR A SECOND-ORDER DIFFERENTIAL OPERATOR WITH A MATRIX POTENTIAL

Keziban Taş¹

¹ Vocational School of Pertek Sakine Genc, University of Munzur, Tunceli, Turkey

kezibantas@munzur.edu.tr,

Abstract

In this paper, we considered an inverse problem for a second-order differential operator with a matrix potential. In the present study, using Mizutani's method, it was obtained the inequality concerning the structure of the potentials difference.

Keywords: Inverse problem; Matrix potential; Wellposedness

REFERENCES

1. Levitan, B.M., Sargsian, I.S., Introduction to spectral theory: selfadjoint ordinary differential operators, American Mathematical Soc., 1975.
2. Mizutani, A.. On the inverse Sturm-Liouville problem. J. Fac. Sci. Univ. Tokyo Sect. IA Math, 31(2), 1984.
3. Uskova, N.B., On Spectral Properties of Sturm–Liouville Operator with Matrix Potential, Ufa Math. J., vol. 7, No: 3, 88–99, 2015.
4. Uskova, N.B., On the Spectral Properties of a Second-Order Differential Operator with a Matrix Potential, Differential Equations, Vol. 52, No:5, 557–567, 2016.

COMPARISON OF DETERMINISTIC AND STOCHASTIC DYNAMICS OF SIR MODEL

Tuğçem Partal¹ and Melike Kakşı

¹Department of Mathematics, Recep Tayyip Erdogan University, Rize, Turkey

tugcem.partial@erdogan.edu.tr,

Abstract

In this paper we analyse the deterministic and stochastic SIR (Susceptible, Infected, Recovered) models for the comparison of epidemic models. We first introduce the deterministic model and then extend it to its stochastic equivalent. While the deterministic model describes the dynamics of disease spread using mathematical differential equations, the stochastic model offers a more realistic approach involving random events and probabilities. The work, also analyses the impact of both models on the parameters using mathematical calculations based on real data and MATLAB software. The effects of important model parameters such as transmission rate and recovery rate on the models and their results were compared and discussed.

Keywords: SIR model, Brownian motion, Stochastic differential equations, Parameter estimation

REFERENCES

1. Moujahid, Abdelmalik, and Fernando Vadillo, A Comparison of Deterministic and Stochastic Susceptible-Infected-Susceptible (SIS) and Susceptible-Infected-Recovered (SIR) Models, Open Journal of Modelling and Simulation 93,1,246-258,2021.
2. Olabode, Damilola, et al., Deterministic and stochastic models for the epidemic dynamics of COVID-19 in Wuhan, China, Mathematical Biosciences and Engineering, 18,1,950-967, 2021.
3. Allen, Linda JS, and Amy M. Burgin, Comparison of deterministic and stochastic SIS and SIR models in discrete time, Mathematical biosciences 163,1, 1-33,2000.
4. He, Sha, Sanyi Tang, and Libin Rong. "A discrete stochastic model of the COVID-19 outbreak: Forecast and control." *Math. Biosci. Eng* 17.4. 2792-2804, 2020.

GPLM for Regression of Complex Systems

Ozlem Defterli^{1,*}, Ayse Ozmen²

¹Department of Mathematics, Cankaya University, Ankara, Turkiye

defterli@cankaya.edu.tr,

² National University of Singapore, Centre for Maritime Studies, Singapore

ayseozmen19@gmail.com

Abstract

Complex systems in a network structure arise in several real-life phenomena where the complexity of such systems can increase due to the existence of a large number of variables and unknown parameters. Linear models are generally used to find the unknown parameters which describe the interactions between the components, but in real-life problems they are non-linearly implied. Therefore, more appropriate models need to be investigated to represent accurately both the linear and nonlinear interactions appearing simultaneously in such systems. In this study, the Generalized Partial Linear Model (GPLM) approach (see [1-3] and references therein) is presented for the dynamical modelling of complex regulatory networks to improve the system identification.

Keywords: Mathematical modelling; Regression; Computational biology.

REFERENCES

1. M. Müller, Estimation and Testing in Generalized Partial Linear Models: A Comparative Study, Statistical Computing, 11, 299–309, 2001.
2. O. Defterli, Modern Mathematical Methods in Modeling and Dynamics of Regulatory Systems of Gene-Environment Networks, Publisher: Graduate School of Natural Sciences, Department of Mathematics, Middle East Technical University, Published: Ankara, 2011.
3. A. Ozmen, G.-W. Weber, Z. Cavusoglu and O. Defterli, The new robust conic GPLM method with an application to finance: prediction of credit default, Journal of Global Optimization, 56(2), 233–249, 2013.

*Corresponding/Presenting Author

PLANAR CONGRUENT CURVES ACCORDING TO CAPUTO FRACTIONAL DERIVATIVE

Ece ATLAN¹ and Handan ÖZTEKİN¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

atlanecce@gmail.com and handanoztekin@gmail.com

Abstract

In this paper, a new definition of the Frenet vector of planar curves by means of the Caputo fractional derivative is given. Then planar congruent curves with respect to the new Frenet vectors, are defined and their properties under some special cases are investigated.

Keywords: Caputo fractional derivative, Plane curves, Congruent curves

REFERENCES

1. Lopez, F.R. ve Rubio, O., (2023) A New Fractional Curvature Of Curves Using The Caputo's Fractional Derivative, Advanced Mathematical Models & Applications Vol.8, No.2, 2023, pp.157-175.
2. Dağışan H. ve Tunçer Y., (2019) Düzlem Eğrilerinin Kendi Frenet Vektörlerine Göre Kongrüent Eğrileri, Uşak Üniversitesi Fen ve Doğa Bilimleri Dergisi, (2), 65-79.
3. Aydın, M.E., Bektas, M., Ogrenmis, A.O. & Yokus, A. (2021). Differential geometry of curves in Euclidean 3-space with fractional order. IEJG, (14), 132-144.
4. Öğrenmiş M. (2022). Geometry of Curves with Fractional Derivatives in Lorentz Plane. Journal of New Theory, (38), 88-98.
5. Aydın, M.E., Mihai, A. & Yokus, A. (2021). Applications of fractional calculus in equiaffine geometry: Plane curves with fractional order. Math. Meth. Appl. Sci., (44), 13659–13669.
6. Yajima, T, Oiwa S, & Yamasaki, K. (2018). Geometry of curves with fractional-order tangent vector and Frenet-Serret formulas. Fractional Calculus and Applied Analysis, 21(6), 1493-1505.

DYNAMICS OF A PLANT-HERBIVORE MODEL SUBJECT TO ALLEE EFFECTS WITH LOGISTIC GROWTH OF PLANT BIOMASS

Emin Beso¹, Senada Kalabušić¹, Esmir Pilav¹, Arzu Bilgin²

¹Department of Mathematics and Computer Sciences, University of Sarajevo, Sarajevo,
Bosnia and Herzegovina

²Department of Mathematics, Recep Tayyip Erdogan University, Rize, Turkey

arzu.bilgin@erdogan.edu.tr

This paper explores the relationship between herbivores and plants with a strong Allee effect, using the logistic equation to model plant growth. We analyze equilibrium points and their stability, identifying several bifurcations. We discover an Allee threshold below which both populations face extinction, but above which coexistence is stable. Numerical simulations suggest that this range of stability can be extended. The system is found to be highly parameter sensitive. A comparison with a system without strong Allee effects is an enrichment of our understanding.

Keywords: Strong Allee effect; bifurcations; logistic growth; plant-herbivore model; stability

REFERENCES

1. Allee, W. C. [1931] Animal aggregations: Study in general sociology, (University of Chicago, Press Chicago).
2. Allen, L. [2006] An introduction to mathematical biology, (Prentice-Hall, Upper Saddle River, NJ).
3. Bešić, E., Kalabušić, S., Pilav, E. [2023] "Dynamics of a plant-herbivore system with Ricker plant growth and the strong Allee effects on plant population," Discrete and Continuous Dynamical Systems Series B (DCDS-B), 2023:108.
4. Bonsall, M. B., van der Meijden, E., Crawley, M. J. [2003] "Contrasting dynamics in the same plant-herbivore interaction," Proceedings of the National Academy of Sciences, 100, 14932–14936.
5. Çelik, C., Duman, O. [2009] "Allee effect in a discrete-time predator–prey system," Chaos, Solitons Fractals, 90, 1952–1956.
6. Chow, Y., Jang, S. R.-J. [2014] "Allee effects in a Ricker-type predator-prey system," Journal of Difference Equations and Applications, 20, 1350–1371.
7. Comins, H. N., McMurtrie, R. E. [1993] "Long-term response of nutrient-limited forests to CO₂ enrichment; equilibrium behavior of plant-soil models," Ecological Applications, 3, 666–681.
8. Elaydi, S. [2005] An introduction to difference equations, 3rd Ed. (Springer).
9. Elaydi, S., Sacker, R. J. [2010] "Population models with Allee effect: A new model," J. Biol. Dyn., 4, 397–408.
10. Feng, Z., DeAngelis, D. L. [2018] "Mathematical models of plant-herbivore interactions," (Chapman & Hall/CRC).

DECODING STRUCTURAL ISOMER: AN ARTIFICIAL INTELLIGENCE APPROACH TO CLUSTER DETECTION

Ömer Akgüller¹, Mehmet Ali Balcı¹

¹Department of Mathematics, Mugla Sitki Kocman University, Mugla, Turkey

oakguller@mu.edu.tr, mehmetalibalci@mu.edu.tr

Abstract

This study utilizes artificial intelligence to decipher the complex network of isomer connections utilizing sophisticated community detection methods. We utilize artificial intelligence methods to examine molecular networks created from several structural and chemical distance metrics, including Euclidean and Levenshtein distances. Through the utilization of community recognition techniques rooted in deep learning, we are able to discern discrete clusters among the isomers, revealing cohorts that possess common structural and chemical characteristics. The network analysis uncovers notable patterns throughout these isomer communities, providing insights into the molecular traits that support their biological roles and chemical activity. Statistical tests comparing molecular properties across identified clusters demonstrate large disparities in important attributes such as lipophilicity, molecular refractivity, hydrogen bonding capacity, molecule shape, and polar surface area. The results emphasize the practical consequences of the variety of structures in isomer networks and emphasize the ability of AI to uncover intricate connections in chemical systems. Our findings provide a structure for future investigation of molecular variety through the utilization of artificial intelligence-based methods for identifying communities.

Keywords: Deep learning; Mathematical chemistry; Complex networks.

REFERENCES

1. Fisicaro, Giuseppe, Bastian Schaefer, Jonas A. Finkler, and Stefan Goedecker. Principles of isomer stability in small clusters. *Materials Advances* Vol. 4, No. 7, 1746-1768, 2023.
2. Modee, Rohit, Sheena Agarwal, Ashwini Verma, Kavita Joshi, and U. Deva Priyakumar. DART: deep learning enabled topological interaction model for energy prediction of metal clusters and its application in identifying unique low energy isomers. *Physical Chemistry Chemical Physics* Vol. 23, No. 38, 21995-22003, 2021.

MEASURING AND ASSESSING ORGANIZATIONAL DATA MATURITY

Merve Ak¹, Senem Şahan Vahaplar¹, M. Hakkı Ersoy¹, Ahmet Feyzioğlu²

¹Experilabs (SahaBT Yazılım), R&D Center, İstanbul, Türkiye

²Department of Mechanical Engineering, Marmara University, İstanbul, Türkiye

senem.vahaplar@experilabs.com

Abstract

In today's competitive business environment, data is a crucial factor. Infinite amount of data is produced from many different sources for almost all business types, but these data are meaningless unless analyzed and transformed into significant results. Therefore; as the organizations improve their ability to be data driven, the more competitive they become. The importance of data maturity concept emerges at this point.

Data maturity is defined as a journey towards improvement and increased capacity in data use, and it is a measure of how advanced organizations are in their data-driven way of doing business. It can be measured not by how much data a business collects, but by how much it processes, analyzes and uses this data in its business processes to make the right decisions. In this study, we study a framework to collect data through sectoral questions specific to organizations, make changes on the question sets autonomously with dynamic learning methods depending on the answers received, determine the level of data maturity based on the outputs of the analyses. Also, the system will be supported with various suggestions for end-users to make decisions accordingly.

Keywords: Data maturity; Maturity assesment; Data management.

REFERENCES

1. Miftahu Rahmatika, Dewi Krismawati, Sinta Denovi Rahmawati, Assaf Arief, Dana Indra Sensuse, Muhammad Fadhil Dzulfikar, An Open Government Data Maturity Model: A Case Study in BPS-Statistics Indonesia, *2019 7th International Conference on Information and Communication Technology (ICoICT)*, Kuala Lumpur, Malaysia, 1-7, 2019.
2. Didier Grimaldi, Javier Diaz, Hugo Arboleda, Vicenc Fernandez, Data maturity analysis and business performance. A Colombian case study, *Heliyon*, Vol:5, 1-9, 2019.

The Novel Numerical Solutions of the Cahn-Hilliard Equation via the Novel Hybrid Method

Aslı Alkan¹, Tolga Aktürk², Hasan Bulut³

¹Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

²Department of Mathematics and Science Education, Faculty of Education, Ordu University, Ordu, Turkey

³Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

alkanasli47@gmail.com, tolgaakturkk@gmail.com, hbulut@firat.edu.tr

Abstract

The objective of this study is to explore innovative numerical techniques for solving the Cahn-Hilliard equation. A new hybrid approach is employed to solve the equation. Furthermore, the solutions are visualized using Maple software through two-dimensional and three-dimensional graphs. Computer simulations are performed to verify the effectiveness and dependability of the proposed method.

Keywords: Novel hybrid method, Numerical solution, Cahn-Hilliard equation.

REFERENCES

1. Alkan, A., Aktürk, T., Bulut, H. 2024. The Traveling Wave Solutions of the Conformable Time-Fractional Zoomeron Equation by Using the Modified Exponential Function Method. *Eskişehir Technical University Journal of Science and Technology A-Applied Sciences and Engineering*, 25(1), 108-114.
2. Miranville, A. (2003). Generalized Cahn-Hilliard equations based on a microforce balance. *Journal of Applied Mathematics*, 2003, 165-185.
3. Cahn, J. W., & Hilliard, J. E. (1958). Free energy of a nonuniform system. I. Interfacial free energy. *The Journal of chemical physics*, 28(2), 258-267.
4. Benattia, M. E., & Belghaba, K. 2021. Shehu conformable fractional transform, theories and applications. *Cankaya University Journal of Science and Engineering*, 18(1), 24-32.

The New Numerical Solutions of the Navier-Stokes Equation with the New Hybrid Method

Aslı Alkan¹, Tolga Aktürk², Hasan Bulut³

¹Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

²Department of Mathematics and Science Education, Faculty of Education, Ordu University, Ordu, Turkey

³Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

alkanasli47@gmail.com, tolgaakturk@gmail.com, hbulut@firat.edu.tr

Abstract

The objective of this study is to explore the new numerical solutions to the Navier-Stokes equation. The new hybrid method is utilized to solve Navier-Stokes equation. Additionally, two and three-dimensional graphs of the obtained solutions were drawn in Maple software. The purpose of the computer simulations was to verify the effectiveness and dependability of the suggested approach.

Keywords: New hybrid method, Numerical solution, Navier-Stokes equation.

REFERENCES

1. Alkan, A., Aktürk, T., Bulut, H. 2024. The Traveling Wave Solutions of the Conformable Time-Fractional Zoomeron Equation by Using the Modified Exponential Function Method. *Eskişehir Technical University Journal of Science and Technology A-Applied Sciences and Engineering*, 25(1), 108-114.
2. Burqan, A., El-Ajou, A., Saadeh, R., & Al-Smadi, M. (2022). A new efficient technique using Laplace transforms and smooth expansions to construct a series solution to the time-fractional Navier-Stokes equations. *Alexandria Engineering Journal*, 61(2), 1069-1077.
3. Mukhtar, S., Shah, R., & Noor, S. (2022). The numerical investigation of a fractional-order multi-dimensional Model of Navier–Stokes equation via novel techniques. *Symmetry*, 14(6), 1102.
4. Benattia, M. E., & Belghaba, K. (2021). Shehu conformable fractional transform, theories and applications. *Cankaya University Journal of Science and Engineering*, 18(1), 24-32.

The Novel Numerical Solutions of the Rosenau-Hyman Equation via the Novel Hybrid Method

Aslı Alkan¹, Tolga Aktürk², Hasan Bulut³

¹Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

²Department of Mathematics and Science Education, Faculty of Education, Ordu University, Ordu, Turkey

³Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey

alkanasli47@gmail.com, tolgaakturkk@gmail.com, hbulut@firat.edu.tr

Abstract

The aim of this study is to investigate novel numerical methods for solving the Rosenau-Hyman equation. The Rosenau-Hyman equation is solved using a new hybrid approach. In addition, the solutions were graphed in Maple software using both two-dimensional and three-dimensional graphs. The computer simulations were conducted to validate the efficacy and reliability of the proposed approach.

Keywords: Novel hybrid method, Numerical solution, Rosenau-Hyman equation.

REFERENCES

1. Alkan, A., Aktürk, T., Bulut, H. 2024. The Traveling Wave Solutions of the Conformable Time-Fractional Zoomeron Equation by Using the Modified Exponential Function Method. *Eskişehir Technical University Journal of Science and Technology A-Applied Sciences and Engineering*, 25(1), 108-114.
2. Yasmin, H., Alshehry, A. S., Saeed, A. M., Shah, R., & Nonlaopon, K. 2023. Application of the q-Homotopy Analysis Transform Method to Fractional-Order Kolmogorov and Rosenau-Hyman Models within the Atangana-Baleanu Operator. *Symmetry*, 15(3), 671.
3. Wang, Y., Sun, Z. & Li, H. 2020. Numerical study of the fractional-order Rosenau-Hyman equation using finite difference methods. *J. Comput. Phys.* 417, 109697.
4. Benattia, M. E., & Belghaba, K. 2021. Shehu conformable fractional transform, theories and applications. *Cankaya University Journal of Science and Engineering*, 18(1), 24-32.

HOW TO DETERMINE THE OPTIMAL STRATEGIES TO ELIMINATE THE HARMFUL EFFECTS OF TECHNOLOGY ADDICTION?

Derya Avcı^{1,*} and Sanem Sakarya¹

¹Department of Mathematics, Balıkesir University, Balıkesir, Turkey

*dkaradeniz@balikesir.edu.tr, sanemsakarya45@gmail.com

Abstract

Addictions spread from person to person, just like epidemics. For this reason, various mathematical models have been developed and optimal control strategies have been realized in order to understand the dynamics of addictions and prevent the harmful effects they cause.

With the developing technology, technology addiction has become a factor affecting our lives. This study aims to control the level of an individual's technology addiction and examines the cost of combating addiction with different control strategies. The discussed SLHB model is developed as SLHBR model by considering the recovered compartment into the model.

Before formulating the optimal control problem for the model, the existence of optimal control is ensured. Then, the optimal system is obtained by using Hamilton's formalism. Numerical results are simulated with the Python programming language. In the graphs, the effect of the proposed control strategies and the uncontrolled model are compared. Thus, the importance of implementing optimal strategies to prevent technology addiction is revealed.

Keywords: Technology Addiction; Optimal Control; Mathematical modeling; Epidemiology.

REFERENCES

1. Zara, M. C., Monteiro, L. H., The negative impact of technological advancements on mental health: An epidemiological approach. Applied Mathematics and Computation, Vol:396,125905,2021.
2. Zhou, M., Zhu, W., Sun, X., Huang, L., Internet addiction and child physical and mental health: Evidence from panel dataset in China. Journal of affective disorders, Vol:309, 52-62,2022.
3. Kirk, D. E., Optimal Control Theory: An Introduction. Courier Corporation,2004.
4. Saman, A., Side, S., Pratama, M. I., Sanusi, W., Optimal control of the SEIR model of online game addiction using guidance and counseling. Engineering Letters, Vol:30, No:1,27-31,2022.

AN OPTIMAL TRAINING POLICY TO REDUCE CRIMINAL BEHAVIOR

Derya Avcı^{1,*} and Aylin Yetim¹

¹Department of Mathematics, Balıkesir University, Balıkesir, Turkey

*dkaradeniz@balikesir.edu.tr, ayetim72@gmail.com

Abstract

Crime and criminal behavior greatly affect countries socially and economically. Specifically, as the crime rate increases, countries increase their expenditures for prisoners. Reducing crime and criminal behavior ensures peace for the citizens of a country. In this study, we consider a mathematical model to reduce crime dynamics. The model divides society into four groups: susceptible individuals, criminally active individuals, criminals in prison, and reformed individuals. By adapting the control function to the model as a training parameter, we examine the behavior of the prescribed model under this control function. To obtain the effect of the training, we simulate the graphs with the help of the MATLAB software. The graphical results show that the suggested training strategy is optimally effective in reducing criminal behavior as desired.

Keywords: Mathematical model; Optimal Control Theory; Crime Model; Pontryagin's Maksimum Principle.

REFERENCES

1. Kwofie, Theophilus, Dogbatsey, Matthias, Moore, Stephen E., Curtailing crime dynamics: A mathematical approach. *Frontiers in Applied Mathematics and Statistics*, 8, 1086745, 2023.
2. Lenhart, Suzanne, and John T. Workman. *Optimal control applied to biological models*. Chapman and Hall/CRC, 2007.
3. Opoku, Nicholas Kwasi-Do Ohene, Georg Bader, and Edem Fiatsonu. "Controlling crime with its associated cost during festive periods using mathematical techniques." *Chaos, Solitons Fractals*, 145 (2021): 110801.

ON f -STATISTICAL CONVERGENCE VIA q -CALCULUS

Nazlım Deniz Aral¹, Hacer Şengül Kandemir² and Mikail Et³

¹Department of Mathematics, Bitlis Eren University, Bitlis, Turkey

²Faculty of Education, Harran University, Şanlıurfa, Turkey

³Department of Mathematics, Fırat University, Elazığ, Turkey

ndaral@beu.edu.tr

hacer.sengul@hotmail.com

mikailet68@gmail.com

Abstract

In this paper, we introduce the the concept of q – statistical convergence and q –strongly Cesàro summability by using a modulus function. We also give some inclusion relation between this concepts.

Keywords: Statistical convergence; q - Cesàro matrix, Modulus function.

REFERENCES

1. M Ayman Mursaleen, S Serra-Capizzano, Statistical convergence via q – Calculus and Korovkin’s Type Approximation Theorem, Axioms, Vol:11, No:2, 2022.
2. Vinod K. Bhardwaj, Shweta Dhawan, f –Statistical convergence of order α and strong Cesàro summability of order α with respect to a modulus, Journal of Inequalities and Applications, 2015, 1-14, 2015.

A RESEARCH ON THE QUALITATIVE BEHAVIOR OF SOLUTIONS OF NEUTRAL SYSTEMS WITH PERIODIC COEFFICIENTS

Yener ALTUN¹, Şakir İŞLEYEN²

¹Department of Business Administration, University of Yüzüncü Yıl, Van, Turkey

²Department of Econometrics, University of Yüzüncü Yıl, Van, Turkey

yeneraltun@yyu.edu.tr

sakirisleyen@yyu.edu.tr

Abstract

In this search, we consider a class of nonlinear neutral type systems (NNSs) with periodic coefficients. Some assumptions guaranteeing the exponential stability of the zero solution are pointed out by using the Lyapunov-Krasovskii functional. Under these assumptions, we give some estimates that characterizing the decay rate of solutions at infinity. We give two examples to demonstrate the applicability of the obtained results.

Keywords: Exponential decay, Lyapunov functional, neutral systems, periodic coefficient.

REFERENCES

1. Demidenko GV, Matveeva II. On estimates for solutions of systems of differential equations of neutral type with periodic coefficients. Siberian Mathematical Journal 2014; 55(5): 866–881.
2. Matveeva II. Exponential stability of solutions to nonlinear time-varying delay systems of neutral type equations with periodic coefficients. Electronic Journal of Differential Equations 2020; 2020(20): 1-12.
3. Altun Y. Improved results on the stability analysis of linear neutral systems with delay decay approach. Mathematical Methods in the Applied Sciences 2020; 43(3): 1467–1483.

Explicit Solutions of the Schrödinger Equation Using Fractional Analysis

Fatma Uzer¹, Resat Yilmazer¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

uzerfatma676@gmail.com, ryilmazer@firat.edu.tr,

Abstract

In this article, the Schrödinger equation with Coulomb potential in α -dimensional fractional space is considered and the radial and angular equations are obtained by the method of separation of variables. Appropriate transformations are made for the solution of the radial equation and the explicit solutions of the second order linear differential equation with singular coefficients are obtained with the help of fractional analysis.

Keywords: Fractional calculus; Schrödinger equation; Differential equations.

REFERENCES

1. Podlubny, I. Fractional Differential Equations; Academic Press: New York, NY, USA, 1999.
2. Eid, R., Muslih, S. I., Baleanu, D., Rabei, E., On Fractional Schrödinger Equation in α -dimensional Fractional Space, Nonlinear Analysis: RWA, 10, 1, 2009.
3. Yilmazer, R. N-fractional calculus operator Nm method to a modified hydrogen atom equation. Math. Commun. 15, 489–501, 2010.
4. Yilmazer, R., Ozturk, O., Explicit solutions of singular differential equation by means of fractional calculus operators. Abstr. Appl. Anal. 2013, 6, 2013.

DIRAC SYSTEMS THAT CONTAIN EIGENVALUE DEPENDENT BOUNDARY CONDITIONS

Mine Babaoglu

Department of Mathematics and Science Education, University of Kahramanmaras Sutcu
Imam, Kahramanmaras, Turkey

mnbabaoglu@gmail.com

Abstract

In this article, we deal with the Dirac operator with eigenvalue dependent boundary conditions. Then, we obtain essential and significant results by modifying some known techniques for the presented problem.

Keywords: Dirac system; Eigenvalue dependent bounday conditions; Sampling theory.

REFERENCES

1. B. M. Levitan, I. S. Sargsjan, Introduction to Spectral Theory: Selfadjoint Ordinary Differential Operators, American Mathematical Society, Providence, Rhode Island, 1975.
2. A. I. Zayed, Advances in Shannon's Sampling Theory, CRC Press, Boca Raton, 1993.

STRUCTURAL, OPTOELECTRONIC AND PHOTOVOLTAIC PROPERTIES OF NEW CONJUGATED SMALL MOLECULES FOR ORGANIC SOLAR CELLS

Imane El mhamedi¹, Anass El karkri¹, Zakaria El malki¹, Mohammed bouachrine²

¹High School of Technology, Moulay Ismail University, Meknes, Morocco

²High School of Technology, Sultan Moulay Slimane University, Khenifra, Morocco

anass.elkarkri@gmail.com,

Abstract

A series of new conjugated molecules with small band gaps were studied by DFT and TD-DFT methods to propose them as active layers for organic solar cells. The AMPS-1D program was used to simulate and determine the photovoltaic characteristics and cell performance. The organic compounds studied are the basis of the donor unit Triphenylamine, phenylenevinylene and different units such as 3, 4-ethylenedioxythiophene (EDOT), carbazole and thiophene. DFT and TD-DFT calculations show strong linearity of the molecules and the calculated values of different characteristic quantities of these molecular compounds suggest that they are good candidates for use in organic solar cells. The simulation results of solar cells based on our donor material and PCBM as acceptor confirm the conclusion obtained, especially since the power conversion efficiency (PCE) reaches a significant value of 8.58%.

Keywords: Organic materiel, DFT, Solar cell, Power conversion efficiency.

Numerical Solution of the Singularly Perturbed Cauchy Problem for an Ordinary Differential Equation

Asan OMURALIEV¹, Ella ABYLAEVA¹,

¹Department of Applied Mathematics and Informatics, Kyrgyzstan-Turkey Manas University,
Bishkek, Kyrgyzstan

ella.abylaeva@manas.edu.kg,

Abstract

For a visual description of Lomov's method, we consider the Cauchy problem for a scalar first-order ordinary differential equation:

$$\begin{aligned}L_\varepsilon u(x, \varepsilon) &\equiv \varepsilon u' + a(x)u = f(x), \quad x \in [0, 1], \\ u(0, \varepsilon) &= u^0,\end{aligned}\tag{1}$$

where $a(x) > 0 \quad \forall x \in [0, 1]$, $a(x), f(x) \in C^\infty[0, 1]$, $\varepsilon > 0$ is a small parameter.

Using the method of Lomov [1], we reduce the singularly perturbed problem (1) to a problem regular in ε when $\varepsilon \rightarrow 0$. To do this, let us introduce a regularizing variable in the following form:

$$\xi = \frac{\varphi(x)}{\varepsilon} = \frac{1}{\varepsilon} \int_0^x a(s) ds \tag{2}$$

and, instead of $u(x, \varepsilon)$ we will consider the extended function $\tilde{u}(x, \xi, \varepsilon)$ such that,

$$\tilde{u}(x, \xi, \varepsilon)|_{\xi=\varphi(x)/\varepsilon} \equiv u(x, \varepsilon). \tag{3}$$

Next, we will partially discretize the extended problem and construct a difference scheme.

Keywords: Lomov's method; Cauchy problem; Singularly perturbed problem.

REFERENCES

1. S. A. Lomov, Introduction to the general theory of singular perturbations, Nauka, Moscow, 1981.

INFECTIOUS DISEASE MODELS WITH PROPORTIONAL DERIVATIVES ON TIME SCALES

Gülcan TOKAY¹, Emrah YILMAZ²

¹ Department of Mathematics, Firat University, Elazığ, Türkiye

gulcan.tky99@gmail.com

Abstract

Fractional calculus is a generalization of ordinary calculus, which includes derivatives and integrals of non-integer order. This field dates back to the times of Leibniz, Gauss, and Newton, who invented such calculations centuries ago. For three centuries, the theory of fractional calculus developed solely as a theoretical area useful to mathematicians. However, in recent times, it has drawn the interest of many scientists and engineers. Furthermore, time scale theory was first introduced by Hilger to unify discrete and continuous phenomena in mathematics. The aim of this study is to establish and solve SIR (Susceptible-Infectious-Recovered) model, which is one of the infectious disease models containing proportional derivatives, on a time scale. While solutions for the classical case of the SIR model exist, solutions will be obtained using proportional derivatives on a time scale in this study. Finally, solutions will be a special case of classical solutions.

Keywords: Time Scales; Proportional Derivative; SIR Model.

REFERENCES

1. Hilger, S. (1988). Ein Makettenkalkl mit Anwendung auf Zentrumsmanigfaltigkeiten, Ph. D. Thesis, Universtat Wurzburg.
2. Anderson, D. R., Georgiev, S. G. (2020). Conformable Dynamic Equations on Time Scales. Chapman and Hall/CRC.
3. Mohajan, Haradhan. "Mathematical Analysis of SIR Model for COVID-19 Transmission." (2022): 1-18.
4. Bohner, M. and Georgiev, S. (2016). Multivariable dynamic calculus on time scales, Springer.
5. Gulsen, T., Yilmaz, E., Goktas, S. (2017). Conformable fractional Dirac system on time scales, Journal of Inequalities and Applications, 2017, 161.

The Concretization Process of the “Pyramid” Concept; Deaf Student Example

Nejla GÜREFE¹

¹ Department of Mathematics Education, Faculty of Education, Mersin University, Mersin,
Turkey

nejlagurefe@mersin.edu.tr

Abstract

In this investigation, we explored the utilization of sign language and gestures in concretizing the pyramid from 3-dimensional geometric objects particularly focusing on their integration into the mathematics learning process. Sign language, being the native language of deaf individuals, encompasses systematic and traditional movements of the hands, face, and body. Sign language (SL) primarily employs manual modality, serving as a substitute for speech among deaf or hard-of-hearing students. Gestures are defined as spontaneous movements of the arms and hands that are closely synchronized with speech. The participants in this case study consist deaf students. Data were gathered through semi-structured interviews and document analysis, which were subsequently subjected to content analysis. The analysis revealed that hand signals, including gestures and sign language, are predominantly employed in elucidating the pyramid and its attributes, while representations such as writing and drawing figures are less frequently utilized. Based on these findings, it is suggested that educators incorporate hand, arm, and body movements into their classroom practices to facilitate the concretization of mathematical concepts, given the significant utilization of hand signs by students.

Keywords: Sign Language; Gesture; Pyramid; Deaf Student

REFERENCES

1. Brentari, D. (2010). Sign Languages. Cambridge University Press.
2. Fay, N., Lister, C., Ellison, T.M. & Goldin-Meadow, S. (2014). Creating a communication system from scratch: Gesture beats vocalization hands down. In: Berent, I., Goldin-Meadow, S., editors. Language by Hand and by Mouth, *Frontiers in Psychology. Language Sciences*, 5, p. 354.
3. Goldin-Meadow, S. & Brentari, D. (2017). Gesture, sign and language: The coming of age of sign language and gesture studies. *The Behavioral and Brain Sciences*, 1–82. <http://doi.org/10.1017/S0140525X15001247>.
4. McNeill, D. (1992). Hand and mind: what gestures reveal about thought. Chicago: University of Chicago.

Modified Exponential Function Method for Two-Dimensional Nonlinear Mathematical Model

Yusuf GUREFE¹

¹Department of Mathematics, Faculty of Science, Mersin University, Mersin, Turkey

ygurefe@gmail.com,

Abstract

In this study, we use the modified exponential function method (MEFM) to investigate the solutions of 2DBSE. MEFM is a powerful mathematical technique for obtaining exact and approximate solutions to nonlinear PDEs, especially soliton solutions. In this way, we obtain traveling wave solutions belonging to the nonlinear mathematical model. We analyze the behavior of these solution functions with the help of the program.

Keywords: Modified Exponential Function Method (MEFM); The two-dimensional breaking soliton equation (2DBSE); Traveling wave solutions.

REFERENCES

1. Ablowitz, M. J., & Segur, H. (1981). Solitons and the Inverse Scattering Transform. SIAM.
2. Dai, H.-H., & Wu, Y.-S. (2008). New periodic wave solutions for the two-dimensional breaking soliton equation. Chaos, Solitons & Fractals, 36(1), 118–126.
3. Hirota, R. (1971). Exact solution of the Korteweg-de Vries equation for multiple collisions of solitons. Physical Review Letters, 27(18), 1192–1194.
4. Xu, Z., Zhang, H.-Q., & Li, Z.-B. (2019). Modified exponential function method for solving nonlinear partial differential equations. Communications in Nonlinear Science and

The Calculation of the Trace Formulas for Dirac Operator by Lax Method

E.Panakhov and F.Asadli

Institute of Applied Mathematics, Baku State University, Baku,
Azerbaijan

In this work the trace formula is calculated by Lax method [1] for Dirac operator in $[0, \pi]$. We consider Dirac operator [2] with self-conjugated boundary condition

$$Ly = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} + \begin{bmatrix} p(x) & 0 \\ 0 & q(x) \end{bmatrix} \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix} = \mu \begin{bmatrix} y_1(x) \\ y_2(x) \end{bmatrix}$$

where $p(x), q(x) \in C^2[0, \pi]$. Let L_0 is operator L in the case $p(x) = 0, q(x) = 0$.

Theorem .1 *Let $\{\mu_n\}, n \in \mathbb{Z}$ denote eigenvalues of operator L , $\{\mu_n(0)\}, n \in \mathbb{Z}$ denote eigenvalues of operator L_0 , then*

$$\sum_{n=-\infty}^{\infty} [\mu_n - \mu_n(0)] = \int_0^{\pi} p(x) S_1(x) dx + \int_0^{\pi} q(x) S_2(x) dx,$$

where

$$S_1(x) = \sum_{n=-\infty}^{\infty} [y_{n1}^2(x, 0) - y_{n2}^2(x, 0)], \quad S_2(x) = \sum_{n=-\infty}^{\infty} [2y_{n1}(x, 0)y_{n2}(x, 0)]$$

$(y_{n1}(x, 0)y_{n2}(x, 0))^T, n \in \mathbb{Z}$ eigenfunction of operator L_0 . We denote that the sums

$$\sum_{n=-\infty}^{\infty} [y_{n1}^2(x, 0) - y_{n2}^2(x, 0)], \quad \sum_{n=-\infty}^{\infty} [2y_{n1}(x, 0)y_{n2}(x, 0)]$$

are convergent.

Corollary .1 *In the case of periodic boundary conditions $y_1(0) = y_1(\pi), y_2(0) = y_2(\pi)$ trace formula has the following form:*

$$\sum_{n=-\infty}^{\infty} [\mu_{2n}(x, 0) + \mu_{2n+1}(x, 0) - 4n] = 0$$

References:

- [1] Lax P.D., Comm.Pure and Appl.Math., 1994, vol. XLVII
- [2] Lidskiy V.F., Sadovnichiy V.A., Dokl.AN USS, 1953, vol.88, no 4

The Calculation of the Regularization Trace of the Diffusion Equation by Lax's Method

E.S.Panakhov¹, I.Shikhaliyeva²

1. Institute of Applied Mathematics, Baku State University
2. Faculty of Mathematics and Mechanics of Baku State University

In this work, by using Lax's method [1], we calculate trace formulas of the equation

$$-y'' + [2\lambda p(x) + q(x)]y = \lambda^2 y, \quad x \in [0, \pi] \quad (1)$$

with boundary conditions

$$y(0) = y'(\pi) = 0 \quad (2)$$

$$y'(0) = y(\pi) = 0 \quad (3)$$

where

$$q(x) \in W_2^m[0, \pi], \quad p(x) \in W_2^{m+1}[0, \pi] \quad (m \geq 0)$$

Let

$$\dots, \chi_{-2}, \chi_{-1}, \chi_1, \chi_2, \dots$$

$$\dots, \nu_{-2}, \nu_{-1}, \nu_1, \nu_2, \dots$$

is a discrete spectrum corresponding to three boundary problems (1), (2) and (1), (3). For the problem (1), (2) and (1), (3) trace formula has the form respectively

$$\sum_{n=1}^{\infty} (\chi_n + \chi_{-n} - 2c_0) = \frac{p(0) - p(\pi)}{2}$$
$$\nu_0 + \sum_{n=1}^{\infty} (\nu_n + \nu_{-n} - 2c_0) = \frac{p(0) - p(\pi)}{2}$$

References:

- [1] Lax P. D. Trace formulas for the Schrodinger operator. Comm. Pure and Applied Math., vol XLVII, pp 503 – 512
- [2] Guseinov G. Sh. Vestnik MGU, 1984, № 3, pp. 14 – 21

EXAMINING THE RELATIONSHIP BETWEEN INTEGRAL EQUATIONS AND DIFFERENTIAL EQUATIONS

Cemil İNAN

¹Department of Business, University of Mardin Artuklu, Mardin, Turkey

cemilinan@artuklu.edu.tr

Abstract

In this study, Integral and Differential equations were introduced and the relations between them and how they could be converted to each other were investigated. It was examined on samples. **Keywords:** Exponential method; Kundu-Eckhaus; Complex exponential.

Keyword : *Differential equations, Integral equations, solved examples*

REFERENCES

1. Aksoy Yavuz, İntegral denklemleri Yıldız Teknik; Üniversitesi İatanbul 1998, pp.1-24
2. Çatalbaş, M.N “Pizetti expansion in terms of a singular elliptic operatör and some application”
3. Tricomi, F. G. İntegral Equations Newyork 1957. Pp. 32-46

ASYMPTOTICS OF SOLUTIONS TO A FIRST-ORDER PARTIAL DIFFERENTIAL EQUATION WITH A POWER-LAW BOUNDARY LAYER

Asan Omuraliev¹, Peyil Esengul kyzy²

¹Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan

asan.omuraliev@manas.edu.kg,

²Department of Mathematics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan

peyil.esengul@manas.edu.kg

Abstract

In the article, a regularized asymptotic of any order of a mixed problem for a first-order partial differential equation is constructed, when the limit equation has a regular singularity. The constructed asymptotic contains boundary-layer functions of two types: power, exponential, and angular functions. The asymptotic of the solution is constructed by a special class of function corresponding to the structure of the fundamental system of solutions. The asymptotic character of the constructed solution is established.

Keywords: a mixed problem; Lomov method; asymptotic of the solution; singularly perturbed problems.

REFERENCES

- [1]. Nesterov A.V., On the asymptotics of the solution of a singularly perturbed system of partial differential equations of the first order with a small nonlinearity in the critical case, Journal of Computational Mathematics and Mathematical Physics, 52(7), 1267 - 1276, 2012. (in Russian)
- [2]. Lomov S. A., Introduction to the general theory of singular perturbations, Moscow(Nauk), 1981. (in Russian)
- [3]. Lomov S. A., Power-law boundary layer in problems with singular perturbation, Izv. AN USSR. Ser. Mat., 30(3), 525-572, 1966. (in Russian)
- [4]. Omuraliev A. S., Abylaeva E. D. and Esengul kyzy P., A system of singularly perturbed parabolic equations with a power boundary layer, Lobachevskii Journal of Mathematics, 41(1), 71-79, 2020.

*Abstract Submission should be prepared only **1 page**.

ARMAMENT MODEL AND ITS ANALYSIS WITH PROPORTIONAL DERIVATIVE ON TIME SCALES

Zehra ÖZDEMİR¹, Emrah YILMAZ²

²Department of Mathematics, University of Firat, Elazig, Türkiye

2321zehra@gmail.com

Abstract

The time scale is a non-empty subset of the real numbers. Time scale theory is a relatively new research area. This theory was first developed in 1988 by Hilger in his doctoral thesis with the aim of unifying differential equations theory and difference equations. The idea of combining fractional calculus with calculations on time scales was first addressed in Bastos' doctoral thesis with the aim of developing fractional calculus on time scales. Recently, several studies involving nabla, delta, and symmetric fractional calculus on arbitrary time scales have been considered. However, Laplace transform plays an important role in solving differential equations where the integral of the function cannot be obtained. It is a simple and useful transformation for solving differential equations with given initial conditions. The Laplace transform transforms differential equations into algebraic equations, simplifying problems mathematically. In this study, the armament model will be solved using classical methods and proportional Laplace transforms on a time scale, and the results will be compared.

Keywords: Time Scales; Proportional Derivative; Proportional Laplace Transform; Armament Model.

REFERENCES

1. Hilger, S. (1988). Ein Makettenkalkl mit Anwendung auf Zentrumsmanigfaltigkeiten, Ph. D. Thesis, Universtat Wurzburg.
2. Anderson, D. R., Georgiev, S. G. (2020). Conformable Dynamic Equations on Time Scales. Chapman and Hall/CRC.
3. Özalp, N. Matematiksel modelleme. Gazi Kitabevi, 2006.
4. Bohner, M. and Georgiev, S. (2016). Multivariable dynamic calculus on time scales, Springer.
5. Yılmaz, E., and Goktas, S. (2021). "On the solution of a Sturm-Liouville problem by using Laplace transform on time scales". Cumhuriyet Science Journal, Vol. 42, No.1, pp. 132-140.

BLOW UP AT INFINITE TIME OF SOLUTIONS FOR THE VISCOELASTIC PLATE EQUATION WITH DISTRIBUTED DELAY AND SOURCE TERMS

Hazal Yüksekaya¹

¹Department of Mathematics, University of Hakkari, Hakkari, Turkey

hazalyuksekkaya@hakkari.edu.tr,

Abstract

In the eighteenth century, the first equations with delay were considered by brothers Leonard Euler and Bernoulli. By A. Myshkis and R. Bellman, systematical study started at the 1940s. Since 1960, there have been appeared many surveys on the subject. In the middle of 1990s, robust control of systems with uncertain delay was started and led to the "delay bloom" in the beginning of the twenty-first century. Time-delay systems are also named systems with aftereffect or dead-time, equations with deviating argument, hereditary systems, or differential-difference equations. They belong to the class of functional differential equations which are infinite-dimensional, as opposed to ordinary differential equations. In this paper, we consider the viscoelastic plate equation with time delay and source term. At first, we give the local and global existence results. Later, under suitable conditions, we prove the blow up of solutions at infinite time by using the energy method. Time delays often appear in many practical problems such as thermal, economic phenomena, biological, chemical, physical, electrical engineering systems, mechanical applications and medicine.

Keywords: Blow up; Plate equation; Time delay.

REFERENCES

1. Fridman, E., Introduction to Time-Delay Systems. Birkhäuser Basel, 2014.
2. Richard, J. P. (2003). Time-delay systems: an overview of some recent advances and open problem, Automatica, 39, 1667 -1694.
3. Yüksekaya, H., & Pişkin, E. (2022). Local existence, global existence and decay results of a logarithmic wave equation with delay term. Mathematical Methods in the Applied Sciences, 1-12.
4. Yüksekaya, H., Piskin, E., Kafini, M. M., & Al-Mahdi, A. M. (2023). Well-posedness and exponential stability for the logarithmic Lamé system with a time delay. Applicable Analysis, 1-13.
5. Yüksekaya, H., Piskin, E., Kafini, M. M., & Al-Mahdi, A. M. (2023). General energy decay estimate for a viscoelastic damped swelling porous elastic soils with time delay. Mathematical Methods in the Applied Sciences, 1-16.

BLOW UP RESULTS AT FINITE TIME FOR THE KIRCHHOFF-TYPE VISCOELASTIC EQUATION WITH TIME DELAY AND SOURCE TERM

Hazal Yüksekaya ¹

¹ Department of Mathematics, University of Hakkari, Hakkari, Turkey

hazalyuksekkaya@hakkari.edu.tr,

Abstract

In this paper, we consider a Kirchhoff-type viscoelastic equation with time delay and source term. Under suitable conditions, we prove the blow-up results at finite time. Controlling the behavior of solutions for partial differential equations with time delay effects has become an active research area. Generally, delay effects occur in many applications and practical problems such as physical, chemical, biological, thermal and economics. In many cases, delay is a source of instability, even an arbitrarily small delay may destabilize a system which is uniformly asymptotically stable in the absence of delay unless additional conditions or control terms have been used.

Keywords: Blow-up; Time delay; Viscoelastic equation.

REFERENCES

1. M. Kafini and S.A. Messaoudi, (2020). Local existence and blow-up of positive-initial-energy solutions of a nonlinear wave equation with delay, *Nonlinear Stud.*, 27(3), 865-877.
2. S. Nicaise and C. Pignotti, (2006). Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, *SIAM J. Control Optim*, 45(5), 1561-1585.
3. H. Yüksekaya, E. Pişkin, S. M., Boulaaras and B. B. Cherif, (2021). Existence, Decay, and Blow-Up of Solutions for a Higher-Order Kirchhoff-Type Equation with Delay Term. *Journal of Function Spaces*, 2021.
4. H. Yüksekaya, & E. Pişkin (2022). Local existence, global existence and decay results of a logarithmic wave equation with delay term. *Mathematical Methods in the Applied Sciences*, 1-12.
5. H. Yüksekaya, E. Pişkin, M. M. Kafini & A. M. Al-Mahdi, (2023). Well-posedness and exponential stability for the logarithmic Lamé system with a time delay. *Applicable Analysis*, 1-13.

ON PRİME SUBHYPERMODULES

Yılmaz Erol¹ and Ummahan Acar²

^{1,2} Department of Mathematics, University of Muğla Sıtkı Koçman, Muğla, Turkey

yilmazerol2@posta.mu.edu.tr, uacar@mu.edu.tr

Abstract

In this paper, we study generalizations obtained on the class of prime subhypermoudles of an R-hypermodule, where R is a Krasner hyperring. We give some new properties and examples of S- prime subhypermoudles. Also, the definition of almost S- prime subhypermoudles and examples are presented.

Keywords: Hyper submodules, S-prime hyper submodules, almost S-asal

REFERENCES

1. Ameri R., On The Prime Submodules of Multiplication Modules. International Journal of Mathematics and Mathematical Science 27: 1715-1724,2013
2. Davvaz B., Leoreanu-Fotea V., Hyperring Theory and Applications, International Academic Press, USA,2007
3. Davvaz B., A Walk Thourgh Hyper Structure, International Academic Press, USA,2013
4. Marty F. Sur une Generalizations de La Notion de Groupe 8th Congress Math. Scandenaves, Stockholm, 8: 45-49, 1934
5. Erol Y., Bozdaş M., Acar U., 35.Ulusal Matematik Semp., Asal Althipermodüllerin Bir Genellemesi, 118, 2023

AN APPROACH TO A FUZZY PROBLEM WITH VARIABLE COEFFICIENTS

Hülya Gültekin Çitil¹, Ayşe Nur Başar²

¹Department of Mathematics, Faculty of Arts and Sciences, Giresun University, Giresun, Turkey

² Institute of Science, Giresun University, Giresun, Turkey

hulya.citil@giresun.edu.tr, ay_se_nur_91@hotmail.com

Abstract

In this work, we consider a fuzzy problem with variable coefficients. We prove the existence of solutions. We illustrate the problem with examples.

Keywords: Fuzzy problem; Fuzzy sets; Fuzzy differential equation.

REFERENCES

1. F. S. Pedro, S. A. B. Salgado, D. E. Sánchez, E. Esmi, L. C. de Barros, On fuzzy Laplace transform in linearly correlated fuzzy space, *Soft Computing*, 27, 1425–1438, 2023.
2. H. Gültekin Çitil, Comparison results of linear differential equations with fuzzy boundary values, *Journal of Science and Arts*, 1(42), 33-48, 2018.
3. S. Karpagappriya, N. Alessa, P. Jayaraman, K. Loganathan, A Novel Approach for Solving Fuzzy Differential Equations Using Cubic Spline Method, *Mathematical Problems in Engineering*, Vol. 2021, Article ID 5553732, 1-9, 2021.



8th International Conference on Computational Mathematics and Engineering Sciences

17 – 19 May 2024,
Şanlıurfa – Türkiye

