

**9TH INTERNATIONAL
CONFERENCE ON
COMPUTATIONAL
MATHEMATICS AND
ENGINEERING SCIENCES**

17 – 19 May 2025,
Diyarbakır – Türkiye

**BOOK OF
ABSTRACTS**

Publish date: 12.05.2025

THE NINETH INTERNATIONAL CONFERENCE ON COMPUTATIONAL MATHEMATICS AND ENGINEERING SCIENCES (CMES- 2025), DİYARBAKIR/TÜRKİYE, MAY 17-19, 2025

The Ninth International Conference on Computational Mathematics and Engineering Sciences (CMES-2025) will be held in Dicle University from 17- to 19 May 2025 in Dicle, Türkiye. It provides an ideal academic platform for researchers and professionals to discuss recent developments in both theoretical, applied mathematics and engineering sciences. This event also aims to initiate interactions among researchers in the field of computational mathematics and their applications in science and engineering, to present recent developments in these areas, and to share the computational experiences of our invited speakers and participants.

The Organizing Committee

©All Rights Reserved. This conference is organized by a cooperation of several international organizations including Fırat University, Fırat International University, Moulay Ismail University, Private University of Fes, Dicle University, Harran University, Van Yüzüncü Yıl University, Manas University, Ordu University Erzurum Technic University and İnönü University. No part of this book can be reproduced or utilized in any forms or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval systems, without permission from the authors.

© The informations provided in the papers published in this book are under the responsibility of their author(s).

Committee Chairs

Hasan Bulut, Fırat University, Türkiye
Zakia Hammouch, Ecole Normale Supérieure de Meknes, Moulay Ismail University, Morocco
Ercan Çelik, Kyrgyz-Turkish Manas University, Kyrgyzstan
Hacı Mehmet Baskonus, Harran University, Türkiye

MESSAGE FROM THE GENERAL CHAIRS



Dear Conference Attendees,

We are honored to welcome you to the **Nineth International Conference on Computational Mathematics and Engineering Sciences (CMES-2025)** at Dicle University from 17 to 19 May 2025 in Diyarbakır City, Türkiye.

CMES, founded in 2016 at Faculty of Science and Techniques Errachidia Moulay Ismail University Morocco is an annual international conference, which was very successful in the past years by providing opportunities to the participants in sharing their knowledge and informations and promoting excellent networking among different international universities. This year, the conference includes 200 extended abstracts, several submissions were received in response to the call for papers, selected by the Program Committee. The program features keynote talks by distinguished speakers such as:

Dumitru Baleanu from Lebanese American University, Beirut, Lebanon; **Bayer Okutmustur** from Middle East Technical University, Türkiye; **Mehrdad Lakestani** from Tabriz University, Iran, **Ekrem Savas** from Usak University, Türkiye; **Ozlem Defterli** from Çankaya University, Türkiye; **Sedaghat Shahmorad** from Tabriz University, Iran. The conference also comprises contributed sessions, posters sessions and various research highlights.

We would like to thank the Program Committee members and external reviewers for volunteering their time to review and discuss submitted abstracts. We would like to extend special thanks to the Honorary, Scientific and Organizing Committees for their efforts in making CMES-2025 a successful event. We would like to thank all the authors for presenting their research studies during our conference. We are grateful to DUBAP(ZGEF.25.003) for research funds. We hope that you will find CMES-2025 interesting and intellectually stimulating, and that you will enjoy meeting and interacting with researchers around the world.

Hasan Bulut,

Firat University, Elazig, Türkiye.

Zakia Hammouch,

ENS Meknes, Moulay Ismail University Morocco

Thu Dau Mot University, Binh Duong Province, Vietnam

China Medical University Hospital Taichung 40402, Taiwan.

TOPICS

Control Theory,
Game Theory,
Applied Mathematics,
Financial Mathematics,
Artificial Intelligence,
Education Sciences,
Engineering Sciences,
Computer Science,
Information Technology,
Geometry and Its Applications,
Analysis and Its Applications,
Statistics and Its Applications,
Algebra and Its Applications,
Topology and Its Application,
Chaos and Dynamical Systems,
Cryptography and its Applications,
Fractional Calculus and Applications,
Economics and Econometric Studies,

Electrical and Electronic Engineering,
Defense industry and applications,
Mathematical Biology,
Computational Epidemiology,
Mathematical Chemistry,
Mathematics Education and Its Applications,
Numerical Methods and Scientific
Programming,
Linear and Nonlinear programming and
Dynamics,
Modeling of Bio-systems for Optimization
and Control,
Ordinary, Partial, Stochastic and Delay
Differential Equations,
Computational Fluids mechanics. Heat and
Mass Transfers,
Earth Sciences,
Applied Sciences

COMMITTEE CHAIRS

Prof.Dr.Hasan Bulut, Firat University, Elazığ, Türkiye

Prof.Dr.Zakia Hammouch, ENS Meknes, Moulay Ismail University, Morocco

Prof. Dr. Ercan Çelik, Kyrgyz-Turkish Manas University, Kyrgyzstan

Prof.Dr.Haci Mehmet Baskonus, Harran University, Türkiye

COMMITTEE CO-CHAIRS

Prof.Dr. Carlo Cattani, Tuscia University, Viterbo, Italy

Prof.Dr. Mohammed Ouazzani Jamil, University Privée de Fes, Morocco

Prof. Dr. Hassan Qjidaa, Sidi Mohammed Ben Abdellah University, Fes Morocco

Prof. Dr. El Mehdi El Khattabi, Moulay Ismail University Morocco

Assoc. Prof. Dr. Tolga Aktürk, Ordu University, Türkiye

HONORARY COMMITTEE

- Prof. Dr. Alpaslan Ceylan, Rector of Kyrgyz-Turkish Manas University, Kyrgyzstan
Prof. Dr. Aytaç Coşkun, Vice Rector of Dicle University, Türkiye
Prof. Dr. Ahmed Mouchtachi, Rector of Moulay Ismail University, Meknes, Morocco
Prof. Dr. Abdelmjid Abourriche, Vice Rector of Moulay Ismail University, Meknes, Morocco
Prof. Dr. Bülent Çakmak, Rector of Erzurum Technical University, Erzurum, Türkiye
Prof. Dr. Ekrem Savaş, Rector of Uşak University, Türkiye
Prof. Dr. Etibar Penahh, Baku State University, Baku, Azerbaijan
Prof. Dr. Fahrettin Göktaş, Rector of Fırat University, Elazığ, Türkiye
Prof. Dr. Haddad Abderrahman, Vice Rector of Moulay Ismail University Morocco
Prof. Dr. Hamdullah Şevli, Rector of Van Yüzüncü Yıl University, Türkiye
İbrahim Taşel, Final International University, Gime, Cyprus
İzzettin Toraman, Chairman of the Board of Toraman Tekstil
Prof. Dr. Kamuran Eronat, Rector of Dicle University, Diyarbakır, Türkiye
Prof. Dr. Mahmut Ergüt, Namık Kemal University, Tekirdağ, Türkiye
Prof. Dr. Mehmet Sıraç Özerdem, Vice Rector of Dicle University, Diyarbakır, Türkiye
Prof. Dr. Mehmet Tahir Güllüoğlu, Rector of Harran University, Sanliurfa, Türkiye
Prof. Dr. Mohammed Aziz Lahlou, President of Universite Privee of Fes, Morocco
Prof. Dr. Necdet Bildik, Retired Faculty Member, Celal Bayar University, Türkiye
Prof. Dr. Orhan Baş, Rector of Ordu University, Ordu, Turkey
Prof. Dr. Orhan Gemikonaklı, Rector of International Final University, Gime, Cyprus
Prof. Dr. Ouazzani Jamil Mohamed, Vice President of Universite Privee of Fez, Morocco
Prof. Dr. Rengin Ak, Rector of Kırklareli University, Kırklareli, Türkiye
Prof. Dr. Rifat Çolak, Retired Faculty Member, Fırat University, Türkiye
Şahin Otu, Chairman of Final Schools Diyarbakır
Prof. Dr. Şengül Kocaman, Dean of Education Faculty, Diyarbakır, Türkiye
Şevket Ertem, Final International University, Gime, Cyprus
Prof. Dr. Sezai Aşubay, Dean of Science Faculty, Diyarbakır, Türkiye
Prof. Dr. Subhan N. Namazov, Vice Rector, Azerbaijan Technical University, Baku, Azerbaijan
Prof. Dr. Velat Şen, Vice Rector of Dicle University, Türkiye
Prof. Dr. Vilayet M. Veliyev, Rector of Azerbaijan Technical University, Baku, Azerbaijan
Prof. Dr. Volkan Öngel, Rector of İstanbul Beykent University, İstanbul, Türkiye
Prof. Dr. Yusif Gasimov, Azerbaijan University, Baku, Azerbaijan

SCIENTIFIC COMMITTEE

- Abdellah Rababah (University of Science and Technology Irbid, Jordan)
Abdellah Rezzouk (Sidi Mohamed Ben Abdellah University, Morocco)
Abdelmjid Bouazizi (ENS Moulay Ismail University Morocco)
Abdelaziz Sbai (EST Moulay Ismail University, Morocco)
Abdelouahed Alla Hamou (FS, Mohamed V University Rabat, Morocco)

Abdulhamit Subaşı (University of Turku, Finland)
Abdulkadir Karakaş (Siirt University, Türkiye)
Abdullah Çağman (Ağrı İbrahim Cecen University, Türkiye)
Abdullah Kopuzlu (Atatürk University, Türkiye)
Abdullahi Yusuf (Biruni University, Istanbul, Türkiye)
Adnène Arbi (University of Carthage, Tunisia)
Agamirza Bashirov (Eastern Mediterranean University, Cyprus)
Ahmed Elngar (Beni Suef University, Egypt)
Ahmet Bedri Özer (Fırat University, Elazığ, Türkiye)
Ahmet Dumlu (Erzurum Technical University, Türkiye)
Ahmet Ocak Akdemir (Ağrı İbrahim Cecen University, Türkiye)
Ahmet Yıldız (İnönü University, Malatya, Türkiye)
Akbar Barati Chiyaneh (Yüzüncüyıl University, Bitlis, Van)
Ali Akgül (Siirt University, Türkiye)
Ali Çakmak (Bitlis Eren University, Bitlis, Türkiye)
Ali İhsan Mut (Dicle University, Türkiye)
Ali Karcı (İnönü University, Malatya)
Ali Rostami (University of Tabriz, Tabriz, Iran)
Ali Yousef (Kuwait College Of Science And Technology, Kuwait)
Alireza Khalili Golmankhaneh (IAU Urmia Branch, Iran)
Alla Belousova (Don State Technical University, Russian Federation)
Alev Dülger (Dicle University, Türkiye)
Alper Ekinci (Bandırma 17 Eylül University, Türkiye)
Alper Osman Öğrenmiş (Fırat University, Elazığ, Türkiye)
Alper Polat (Munzur University, Türkiye)
Aly Ramadan Seadawy (Taibah University, Saudi Arabia)
Anargül Urdaletova (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Aneta Barakoska (Faculty of Philosophy, University St. Cyril, Macedonia)
Anna Sandulyak (Université de Moscow, Russia)
Arife Atay (Dicle University, Türkiye)
Armando Ciancio (University of Messina, Italy)
Arzu Akbulut (Eskişehir Osmangazi University, Türkiye)
Arzu Denk Oguz (Atılım University, Türkiye)
Asan Ömüraliyev (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Atif A. Namazov (Azerbaijan Technical University, Baku, Azerbaijan)
Avıt Asanov (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Ayhan Şerbetçi (Ankara University, Ankara)
Aynur Keskin Kaymakçı (Selçuk University, Konya)
Ayşe Dilek Maden (Selçuk University, Konya)
Ayşe Metin Karakaş (Bitlis Eren University, Bitlis, Türkiye)
Ayşegül Çetinkaya (Ahi Evran University, Türkiye)
Badr Abou E Majd (Mohammed V University in Rabat, Morocco)
Bashar A. Al-Talip (University of Mosul, Iraq)
Bayram Çekim (Gazi University, Türkiye)
Bayram Şahin (Ege University, İzmir, Türkiye)

Behçet Öznacar (Near East University, Cyprus)
Berivan Polat (Munzur University, Türkiye)
Beytullah Yağız (Yüzüncü Yıl University, Türkiye)
Bilal Alatas (Fırat University, Türkiye)
Bilal Şenol (İnönü University, Malatya)
Bilgehan Polatoğlu (Atatürk University, Türkiye)
Burcu Gürbüz (Johannes Gutenberg-University Mainz, Germany)
Burçin Doğan (Malatya Turgut Özal University, Türkiye)
Burhan Tiryakioğlu (Marmara University, Türkiye)
Bülent Tavlı (Tobb Economics And Technology University, Türkiye)
Cameron Browne (University of Louisiana at Lafayette, USA)
Carla M.A. Pinto (ISEP, Porto, Portugal)
Carlo Cattani (Tuscia University, Viterbo, Italy)
Cemil İnan (Mardin Artuklu University, Türkiye)
Ceren Kına (Malatya Turgut Özal University, Türkiye)
Cesim Temel (Yüzüncüyıl University, Türkiye)
Chérif Ziti (Moulay Ismail University, Morocco)
Christos Volos (Aristotle University, Greece)
Çağrı Karaman (Atatürk University, Erzurum, Türkiye)
Çiğdem İnan (Mersin University, Türkiye)
Damla Yılmaz (Erzurum Technical University, Türkiye)
Danyal Soybaş (Erciyes University, Türkiye)
Davut Hanbay (İnönü University, Malatya)
Derya Arslan (Bitlis Eren University, Türkiye)
Derya Avcı (Balıkesir University, Türkiye)
Dilek Okuyucu (Erzurum Technical University, Türkiye)
Dumitru Baleanu (Cankaya University, Türkiye)
Dursun İrk (Eskişehir Osmangazi University, Eskişehir)
Dyyak Ivan (Ivan Franko National University, Lviv, Ukraine)
Ebru Bozkurt (Ataturk University, Türkiye)
Ecem Acar (Harran University, Şanlıurfa, Türkiye)
El Abbassi Ahmed (FSTE, Moulay Ismail University, Morocco)
El Ayyadi Rachid (FST, Sidi Mohammed Benabdellah University, Morocco)
El Hassan Ben Ahmed (Ens Meknes, Moulay Ismail University, Morocco)
El Houcine Lafhim (Sidi Mohamed Ben Abdellah University, Morocco)
El Khomssi Mohamed (Sidi Mohamed Ben Abdellah University, Morocco)
El Mehdi El Khattabi (Ens Meknes, Moulay Ismail University, Morocco)
Ella Abilayeva (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Elman Hazar (İgdir University, Türkiye)
Emel Biçer (Bingöl University, Türkiye)
Emrah Yılmaz (Fırat University, Elazığ)
Engin Özkan (Erzincan Binali Yıldırım University, Erzincan)
Ennahnahi Nouredine (Sidi Mohamed Ben Abdellah University, Morocco)
Erdal Baş (Fırat University, Elazığ, Türkiye)
Erdal Ekici (Çanakkale Onsekiz Mart University, Çanakkale)

Erdal Ulualan (Kutahya University, Türkiye)
Erdal Ünlüyol (Ordu University, Türkiye)
Erhan Deniz (Kafkas University, Türkiye)
Erhan Güler (Bartın University, Türkiye)
Erhan Pişkin (Dicle University, Diyarbakır, Türkiye)
Erhan Set (Ordu University, Türkiye)
Erol Yaşar (Mersin University, Türkiye)
Esin İlhan (Kırşehir Ahi Evran University, Türkiye)
Esra Karataş Akgül (Siirt University, Türkiye)
Eva Kaslik (University of Timisoara, Romania)
Evrin Toklu (Ağrı İbrahim Cecen University, Türkiye)
Ezgi Kaya (Iğdır University, Türkiye)
Fahd Jarad (Cankaya University, Türkiye)
Fahrettin Muhtarov (Bakü State University, Azerbaijan)
Fatma Ayaz (Gazi University, Türkiye)
Fatma Berna Benli (Erciyes University, Türkiye)
Fatma Bulut (Bitlis University, Türkiye)
Fatma Erdoğan (Firat University, Türkiye)
Fatma Sağsöz (Atatürk University, Türkiye)
Fatma Taşdelen Yeşildal (Ankara University, Türkiye)
Fethi Bin Muhammad Belgacem (Paaet, Shaamyia, Kuwait)
Fethiye Müge Sakar (Dicle University, Türkiye)
Feyza Esra Erdoğan (Ege University, İzmir, Türkiye)
Fikret Aliyev (Bakü State University, Azerbaijan)
Filiz Taşcan (Eskişehir Osmangazi University, Eskişehir)
Firat Evirgen (Balıkesir University, Türkiye)
Francesco Villecco (University Salerno, Italy)
Fulya Şahin (Ege University, İzmir, Türkiye)
Gonca Onargan (Final International University, Girne, Cyprus)
Gökhan Gökdere (Firat University, Elazığ)
Gülşat Muhametjanov (Kyrgyz-Turkish Manas University, Kyrgyzstan)
Gürhan İçöz (Gazi University, Türkiye)
Güven Kaya (Bingöl University, Türkiye)
Hacer Sengul (Harran University, Sanlı urfa, Türkiye)
Haldun Kucuk (Tarsus University, Mersin, Türkiye)
Hamit Akyüz (Gazi University, Ankara, Türkiye)
Hasan Can Kısakürek (İstanbul University, İstanbul, Türkiye)
Hasan Doğan (Karamanoğlu Mehmetbey University, Karaman)
Hasan Taş (Firat University, Elazığ, Türkiye)
Hasan Yusuf (The University of Nigeria, Nsukka)
Hatice Taş (Bitlis Eren University, Bitlis, Türkiye)
Hazım Sürer (Ataturk University, Erzurum, Türkiye)
Hidayet Eski (Mersin University, Türkiye)
Hussain Benazza (ENSAM, Moulay Ismail University Morocco)
Hüseyin Kalaycı (Mersin University, Türkiye)

Hüseyin Kaya (İnönü University, Malatya, Türkiye)
Hüseyin Yılmaz (Ankara University, Ankara)
İlyas Öztürk (Dicle University, Diyarbakır, Türkiye)
İbrahim Arslan (Fırat University, Türkiye)
İbrahim Korkmaz (Kırıkkale University, Türkiye)
İbrahim Okan Öztürk (Dicle University, Diyarbakır)
İsmail Yılmaz (Çanakkale Onsekiz Mart University, Türkiye)
İsmail Hakkı Keskin (Fırat University, Elazığ)
İsmail Küçük (Eskişehir Osmangazi University, Eskişehir)
Iván Reyes (Universidad Autónoma de Querétaro, México)
Jakub Tichy (Czech University of Life Sciences, Czech Republic)
Jean-Côme Toutain (University of Poitiers, France)
João Barbosa (University of Lisbon, Portugal)
João Luís (IST, University of Lisbon, Portugal)
John K. B. (University of York, UK)
Josep A. R. (University of Barcelona, Spain)
Joseph Asongu (University of Juba, Sudan)
Juan E. López (University of Chile, Chile)
Jürgen Noack (Brandenburg University of Technology, Germany)
Júlia Kovács (Technical University of Budapest, Hungary)
Kadir S. (University of Uludag, Türkiye)
Kamran E. (University of Tehran, Iran)
Kamran Yavuz (Ankara University, Ankara, Türkiye)
Kamil Kose (Gazi University, Türkiye)
Karim Jebli (University of Monastir, Tunisia)
Keith A. (University of Oxford, UK)
Kenan Kavasoğlu (Çankırı Karatekin University, Türkiye)
Khaled M. (King Fahd University of Petroleum and Minerals, Saudi Arabia)
Khoa T. (Vietnam National University)
Kohei T. (Kyoto University, Japan)
Kostas G. (University of Thessaloniki, Greece)
Krzysztof K. (Silesian University, Poland)
László Sz. (University of Szeged, Hungary)
Louay Al-Kharouf (University of Bahrain, Bahrain)
Maher Shoukry (Cairo University, Egypt)
Mohamed S. S. (Al-Mansoura University, Egypt)
Mohamed Tarek Hamza (Mansoura University, Egypt)
Mubariz Alvi (University of Sindh, Pakistan)
Murat Karakas (Bitlis University, Türkiye)
Mustafa Arslan (Gazi University, Türkiye)
Mustafa Tunc (Erzurum Technical University, Türkiye)
Nezih Gülkır (Akdeniz University, Türkiye)
Nima Mostafavi (Isfahan University of Technology, Iran)
Nural Özkaya (Dicle University, Diyarbakır, Türkiye)
Ouberka Hanane (ENS, Moulay Ismail University Morocco)

Oğuz Karabacak (İstanbul University, İstanbul, Türkiye)
Ömer Gökhan Kaya (Sakarya University, Türkiye)
Ömer Yılmaz (İnönü University, Malatya, Türkiye)
Ouarghi Khalid (ENS, Moulay Ismail University Morocco)
Rafik Ben Fadhel (University of Carthage, Tunisia)
Ramazan Alper (İnönü University, Malatya, Türkiye)
Rattan Kumar Sahu (IIT Bhubaneswar, India)
Samir Ben Lakhdar (University of Carthage, Tunisia)
Selma R. (University of Graz, Austria)
Sergiy R. (Ukrainian Academy of Science, Ukraine)
Sinan Deniz (Yüzüncü Yıl University, Türkiye)
Sipahi, D (Bilkent University, Türkiye)
Skouri Rachid (EST, Moulay Ismail University Morocco)
Soledad Torregrosa (University of Murcia, Spain)
Srikant Mishra (Biju Patnaik University of Technology, India)
Stephen T. (University of Maryland, USA)
Sulaiman Al-Harbi (King Saud University, Saudi Arabia)
Şakir İşleyen (Yüzüncü Yıl University, Türkiye)
Şule Kocabaş (İstanbul University, İstanbul, Türkiye)
Tamara T. (Kyiv National University, Ukraine)
Tantaoui Mouad (ENS, Moulay Ismail University Morocco)
Tial Mohammed (FSTE, Moulay Ismail University Morocco)
Tomasz Z. (Warsaw University, Poland)
Valter S. (University of Porto, Portugal)
Viktor Barbu (University of Oradea, Romania)
Yılmaz K. (Ege University, İzmir, Türkiye)
Yener Altun (Yüzüncü Yıl University, Türkiye)
Zafer E. (Tunceli University, Türkiye)
Zain Siham (ENS, Moulay Ismail University Morocco)
Zaki Mohammed (ENS, Moulay Ismail University Morocco)
Zekeriya H. (Malatya University, Türkiye)
Zekeriya M. (Atatürk University, Türkiye)
Zubair M. (Ministry of Higher Education, Pakistan)

ORGANIZING COMMITTEE

Prof. Dr. Hasan Bulut, (Chair), Fırat University, Türkiye
Prof. Dr. Zakia Hammouch, (Chair), ENS, Moulay Ismail University Morocco
Prof. Dr. Ercan Çelik, (Chair), Kyrgyz-Turkish Manas University, Kyrgyzstan
Prof. Dr. Hacı Mehmet Baskonus, (Chair), Harran University, Türkiye
Prof. Dr. Hatun Özlem Güney, (Chair), Dicle University, Diyarbakır, Türkiye
Prof. Dr. Erhan Pişkin, (Chair), Dicle University, Diyarbakır, Türkiye
Prof. Dr. Ouazzani Jamil Mohammed, (Co-Chair), Université Privée de Fes Morocco
Prof. Dr. Hassan Qjidaa, (Co-Chair), Sidi Mohammed Ben Abdellah University, Fes Morocco
Prof. Dr. El Mehdi El Khattabi, (Co-Chair), Moulay Ismail University Morocco

Assoc. Prof. Dr. Tolga Aktürk, (Co-Chair), Ordu University, Türkiye
Prof. Dr. Mikail Et, Fırat University, Türkiye
Prof. Dr. Vedat Asil, Fırat University, Türkiye
Prof. Dr. Alper Öğrenmiş, Fırat University, Türkiye
Prof. Dr. Muhittin Evren Aydın, Fırat University, Türkiye
Prof. Dr. Fevzi Erdoğan, Yüzüncüyıl University, Türkiye
Prof. Dr. Alaattin Esen, İnönü University, Türkiye
Prof. Dr. Yusuf Pandır, Bozok University, Türkiye
Prof. Dr. Yusuf Güreffe, Mersin University, Türkiye
Prof. Dr. Bahaddin Sinsoysal, İstanbul Beykent University
Assoc. Prof. Dr. Mahmut Modanlı, Harran University, Türkiye
Assoc. Prof. Dr. Gülnur Yel, Final International University, Girne, Cyprus
Assoc. Prof. Dr. Muhammed Yiğider, Erzurum Technic University, Türkiye
Dr. Nuray Eroğlu, Tekirdağ Namık Kemal University, Türkiye

LOCAL COMMITTEE

Sedat İlhan (Dicle University, Türkiye)
Zehra Yücedağ (Dicle University, Türkiye)
Kemal Özgen (Dicle University, Türkiye)
Mehmet Aydın (Dicle University, Türkiye)
Tuğba Örnek (Dicle University, Türkiye)
Seçil Yalaz (Dicle University, Türkiye)
Arife Atay (Dicle University, Türkiye)
Ömer Oruç (Dicle University, Türkiye)
Damla Barlak (Dicle University, Türkiye)
Alev Dülger (Dicle University, Türkiye)
Yasin Kaya (Dicle University, Türkiye)
Nebi Yılmaz (Dicle University, Türkiye)
Ayşe Fidan (Dicle University, Türkiye)
Gülistan Butakın (Dicle University, Türkiye)
Erkan Sancar (Dicle University, Türkiye)
Hasan Kandemir (Dicle University, Türkiye)
Muhteşem Demir (Dicle University, Türkiye)
Yunus Özişiçi (Dicle University, Türkiye)
Ayşe Demirhan (Dicle University, Türkiye)
Özlem Orhan (Bandırma Onyedil Eylül Ün,
Balıkesir)
Mehmet Karay (Final Int. University, Girne,
Cyprus)
Münevver Tuz (Fırat University, Elazığ, Türkiye)
Ebru Cavlak Aslan (Fırat University, Elazığ,
Türkiye)
Tüba Gülşen (Fırat University, Elazığ, Türkiye)
Bahar Acay (Fırat University, Türkiye)

Ahu Ercan (Fırat University, Elazığ, Türkiye)
Berat Karaagac (Adıyaman University, Türkiye)
Ezgi Kaya (Iğdır University, Iğdır, Türkiye)
Onur Saldır (Van Yüzüncü Yıl University, Van,
Türkiye)
Zelal Temel (Van Yüzüncü Yıl University, Van,
Türkiye)
İbrahim Enam İnan (Fırat University, Elazığ,
Türkiye)
Meltem Öğrenmiş (Fırat University, Elazığ,
Türkiye)
Hajar Farhan Ismael (University of Zakho, Iraq)
Karmina K. Ali (University of Zakho, Iraq)
Sibel Şehriban Ataş (Fırat University, Elazığ,
Türkiye)
Aysenur Akkılıç (Assistant, Fırat University,
Türkiye)
Aslı Alkan (Fırat University, Elazığ, Türkiye)
Beyhan Kemaloğlu (Fırat University, Elazığ,
Türkiye)
Mucahit Saydam (Fırat University, Elazığ,
Türkiye)
Çağrı Yılmaz (Fırat University, Elazığ,
Türkiye)
Merve Zeynep Gecmen (Ağrı İbrahim Çeçen
University, Ağrı, Türkiye)
Necati Erdoğan (Van Yüzüncü Yıl
University, Van, Türkiye)

Arif Özkul (Fırat University, Elazığ, Türkiye)
Süleyman Sarıkaya (Fırat University, Elazığ, Türkiye)
Şeyma Firdevs Korkmaz (Fırat University, Elazığ, Türkiye)
Tuğçem Partal (Recep Tayyip Erdoğan University, Türkiye)
Yasemin Bakır (Fenerbahçe University, Türkiye)
Funda Türk (Bartın University, Türkiye)
Ali Sercan Karakaş (İnönü University, Türkiye)
Neşe Dokumacı Sütçü (Dicle University, Türkiye)

Halis Yılmaz (Dicle University, Türkiye)
Mustafa İlhan (Dicle University, Türkiye)
Ali İhsan Mut (Dicle University, Türkiye)
Yılmaz Gündüzalp (Dicle University, Türkiye)
Mustafa Uğurlu Arslan (General Secretary of Dicle University, Türkiye)
Özkan Cığa (Vice Dean of Education Faculty, Diyarbakır, Türkiye, Türkiye)
Mehmet Serkan Şahin (Vice Dean of Education Faculty, Diyarbakır, Türkiye)

LANGUAGE EDITORIAL COMMITTEE

Suheyra Demirkol Orak (Fırat University, Elazığ, Türkiye)

International Communications Consultant

Rengin AK, Kırklareli University, Kırklareli, Türkiye
İkmet Kemaloğlu, Fırat University, Elazığ, Türkiye
İlkay Açıkgöz Erkaya, Ahi evran University, Türkiye
M. Evren Aydın, Fırat University, Elazığ, Türkiye

PROCEEDINGS

Full version of submitted papers will be published in Special Volumes of reputed journals. Procedure, Guidelines and Checklist for the preparation and submission of papers to the Proceedings of CMES-2025 can be found in the journals websites. The journals in which selected and peer-reviewed full papers of CMES-2025 will be published are as follows:

1. BOOK OF ABSTRACTS [Free of charge]

If Authors submit ABSTRACT TEXTS, then, after getting referees evaluations for these abstracts, they will be published in ABSTRACT PROCEEDING BOOK of CMES-2025. For FULL TEXT PAPERS, Authors have to submit their FULL TEXT PAPERS online via submission system of CMES-2025. These FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2025 after getting at least two positive reports.

2. CONFERENCE PROCEEDINGS [Free of charge]

At the beginning, if Authors submit FULL TEXT PAPERS, then, after getting at least two positive referee reports, FULL TEXT PAPERS will be published in FULL TEXT PROCEEDING BOOK of CMES-2025 with ISBN:77733 number. Therefore, Abstracts of these FULL TEXT PAPERS will **NOT** be published in ABSTRACT PROCEEDING BOOK of CMES-2025.

3. FRACTAL AND FRACTIONAL JOURNAL [SCI-E],

Selected papers from CMES-2025 will be published in a special issue dedicated to the Conference entitled "**Feature Papers for Mathematical Physics**

Section".

https://www.mdpi.com/journal/fractalfract/special_issues/1TAP5BBZ45

This journal is indexed by SCI-E.

4. PROCEEDINGS OF THE INSTITUTE OF MATHEMATICS AND MECHANICS [E-SCI]

Selected papers from CMES-2025 will be published by <https://proc.imm.az/special/>

This journal is indexed by E-SCI.

5.TURKISH JOURNAL OF SCIENCE, [FREE]

Participants of CMES 2025 can submit their good quality papers to Turkish Journal of Science. After the peer review process, the papers will be published at TJOS. The authors must write "CMES 2025" as comments to the editor.

(Editor in Chief: Dr. Ahmet Ocak AKDEMİR) For on-line submission: <https://dergipark.org.tr/tr/pub/tjos>

6.TURKISH JOURNAL OF INEQUALITIES, [FREE]

"Participants of CMES 2025 can submit their good quality papers to Turkish Journal of Inequalities. Selected papers will be published at TJI after the

peer review process. The participants can send their papers to erhanset@tjinequality.com. The authors must write "CMES 2025" as the subject. (Editor in Chief: Prof. Dr. Erhan SET)
<http://tjinequality.com/>

7. MATHEMATICS IN NATURAL SCIENCE (MNS)

Authors can submit their full text paper directly to the journal by using the following link
<https://www.isr-publications.com/mns>

8. MATHEMATICS IN ENGINEERING, SCIENCE AND AEROSPACE (MESA), [FREE, SCOPUS]

"Selected papers will be published after peer review in the Journal of Mathematics in Engineering, Science and Aerospace (MESA)"
(Editor in Chief: Prof. Seenith Sivasundaram)
<http://nonlinearstudies.com/index.php/mesa>

9. APPLIED MATHEMATICS AND NONLINEAR SCIENCES, [SCOPUS]

Participants of CMES 2025 can submit their high quality full text papers to Applied Mathematics and Nonlinear Sciences by selecting CMES-2025 under the Select Article Type Menu.
<https://www.editorialmanager.com/amns/default.aspx>

10. MATHEMATICAL MODELLING AND NUMERICAL SIMULATION WITH APPLICATIONS (MMNSA), [TR DİZİN]

The Special Issue on "Advanced Methods of Modelling and Numerical Computation in Science and Engineering". Authors can submit their full text paper directly to the journal by using the information provided in the following link
https://mmnsa.org/index.php/mmnsa/special_issues/SI-CMES2023

11. SYMMETRY [SCI-E] ; SPECIAL ISSUE "ADVANCES IN MATRIX TRANSFORMATIONS, OPERATORS AND SYMMETRY"

Authors can submit their full text paper directly to the journal by using the following link
https://www.mdpi.com/journal/symmetry/special_issues/Advances_Matrix_Transformations_Operators_Symmetry

12. YUZUNCU YIL UNIVERSITY JOURNAL OF THE INSTITUTE OF NATURAL AND APPLIED SCIENCES (TR-Dizin)

Authors can submit their full text paper directly to the journal by using the following link
<https://dergipark.org.tr/tr/pub/yyufbed>

13. PEDAGOGICAL PERSPECTIVE (PEDPER)

Pedagogical Perspective (PedPer) is an international, double blind reviewing, non-profit, professional scientific journal. PedPer is a journal that accepts manuscripts related to pedagogy and education.
<http://pedagogicalperspective.com/>

PLENARY & INVITED TALKS



MODIFIED FRACTIONAL OPERATORS: THEORY AND APPLICATIONS

Dumitru Baleanu¹

¹Department of Computer Science and Mathematics, Lebanese American University, Beirut, Lebanon and Institute of Space Sciences-Subsidiary of INFLPR,

Magurele-Bucharest, Romania

dumitru.baleanu@lau.edu.lb,

Abstract

Fractional Calculus deals with the study of so-called fractional order integral and derivative operators over real or complex domains, and their applications. In this talk I will discuss the modified ABC operator and I will explain the related properties and its real world applications.

Keywords: Fractional calculus; General kernel; Modified ABC operator.

REFERENCES

1. Mohammed Al-Refai, Dumitru Baleanu, On an extension of the operator with Mittag-Leffler kernel, *Fractals*, 30(51), 2022, Article number 2240129.



A COMPARATIVE STUDY OF ADAPTIVE GRID METHODS FOR BURGERS MODELS

Bayer Okutmustur¹

¹Department of Mathematics, Middle East Technical University (METU), Ankara, Turkey

bayer@metu.edu.tr

Abstract

This work presents comparative numerical study of shock solutions in classical and relativistic a Burgers models using two finite volume-based adaptive mesh refinement (AMR) strategies: *h*-adaptivity (local grid refinement) and *r*-adaptivity (mesh redistribution). The *h*-method dynamically adjusts resolution by adding/removing cells in shock regions, while the *r*-method maintains a fixed cell count but optimally redistributes points to sharpen shock resolution. We compare both approaches in terms of accuracy and efficiency, demonstrating their effectiveness for handling shocks and nonlinear waves in both classical and relativistic cases. This work is based on the joint work [4].

Keywords: Adaptive grid; h-refinement; r-refinement; Monitor function; Finite volume method; Relativistic Burgers equation

REFERENCES

1. J.M. Stockie, J.A. Mackenzie, R.D. Russell, A moving mesh method for one-dimensional hyperbolic conservation laws, **SIAM J. Sci. Comput.** 22 (2001) 1791–1813.
2. H.Z. Tang, T. Tang, Adaptive mesh methods for one- and two-dimensional hyperbolic conservation laws, **SIAM J. Numer. Anal.** 41 (2003) 487–515.
3. A. van Dam, P.A. Zegeling, A robust moving mesh finite volume method applied to 1D hyperbolic conservation laws from magnetohydrodynamics, *Journal of Computational Physics*, Volume 216, Issue 2, 2006, 526-546.
4. B. Okutmustur, K. Vuik, K. Yigit, Adaptive Mesh Refinement Methods for Relativistic Burgers-Schwarzshild Model, in preparation.



The development adventure of mathematics

Ekrem SAVAŞ

Uşak University, Uşak-Turkey

Email: ekremsavas@yahoo.com

Abstract:

Mathematics has always existed since the first steps of culturel history of the living creatures called human beings. It has always been a part of culture with numbers, figures, characteristics and with their applications according to the technical levels of the day. In this study, a general knowledge will be given concerning how math began and what phases it went through up to now. In getting the data, books and articles dealing with the issue have been studied. The findings we have show that it is not possible to say exactly where and when mathematics began. However, we can say, on the basis of findings, that math began in Epypt and Mesopotamia between 3000-2000 B.C. and then spread to other countries.



ADVANCED MATHEMATICAL MODELLING AND INFERENCE OF GENE REGULATORY NETWORKS WITH REAL DATA

Ozlem Defterli¹

¹Department of Mathematics, Faculty of Arts and Sciences,

Cankaya University, Ankara, Turkey

defterli@cankaya.edu.tr

Abstract

Systems biology addresses the challenge of understanding living systems in their entirety, as opposed to concentrating on individual biological elements. One approach to describing biological systems is through networks which are the graphical representations in which nodes denote entities of the system and edges signify the relationships between them. Given that the underlying structure of many networks remains partially or entirely unknown, a key objective of systems biology is to predict the complex and dynamic interactions among genes. This is called *network inference (NI)* that focuses on deducing network structures by leveraging high-throughput data in combination with reverse engineering methods. A fundamental challenge in network inference is the high dimensionality—often involving thousands of genes—contrasted with the relatively small number of available samples. Consequently, gene regulatory network (GRN) inference is inherently under-determined [1,2]. In this study, a time-series gene expression data-set derived from a micro-array chip experiment involving a model eukaryotic organism is used for illustrative purposes. The dataset's key characteristics are analyzed to gain insights into the structure and behavior of the underlying biological process. Subsequently, the temporal dynamics of the system are modeled in a discrete-time framework, employing advanced mathematical modeling techniques to capture the complexity and regulatory mechanisms of gene interactions [3,4].

Keywords: Network inference, Mathematical modelling, Data processing.

REFERENCES

1. Slonim K. Donna, From patterns to pathways: gene expression data analysis comes of age (Review), Nature Genetics, Vol:32(Suppl.), 502–508, 2002.



SOME EFFECTIVE NUMERICAL METHODS FOR FRACTIONAL DIFFERENTIAL EQUATIONS

S. Shahmorad¹

shahmorad@tabrizu.ac.ir

¹Department of Applied Mathematics, University of Tabriz, Tabriz,

East Azarbaijan, Iran

Abstract

Some efficient numerical methods such as block-by-block, fuzzy transform and interpolation based methods are studied for solving linear and nonlinear single term and multi-term fractional differential equations (MFDEs). The approaches involve converting the given linear and nonlinear MFDEs with some initial conditions into equivalent Volterra integral equations (VIE), and applying the mentioned numerical approaches to the obtained VIES. Error bounds and convergence theorems are discussed for each case, separately. Finally, illustrative and comparative examples are provided to demonstrate the applications of the proposed methods and verify the theoretical results.

Keywords: Multi-term fractional initial value problem, Block-by-block method, Fuzzy transform, Nonlinear Volterra integral equation, weak singularity.



NUMERICAL SOLUTION OF SINGULARLY PERTURBED INITIAL VALUE PROBLEMS WITH DELAY USING B-SPLINE WAVELETS

MEHRDAD LAKESTANI

Department of Applied Mathematics, University of Tabriz, Tabriz, Iran

mehrdadlakestani@ydu.edu.tr

ABSTRACT

In this paper, we solve a singularly perturbed initial value problem with delay by using c B-Spline wavelets. The properties of these functions are provided, and by employing the operational matrix of differentiation, a numerical method is generated over some subintervals that reformulates the problem into a system of algebraic equations. This system can be solved to find the approximate solution. Numerical results demonstrate the efficiency of the method.

Keywords: Singularly perturbed problem; B-Spline wavelets; Delay differential equation.

REFERENCES

1. M. Lakestani, M. Dehghan, The solution of a second-order nonlinear differential equation with Neumann boundary conditions using semi-orthogonal B-spline wavelets, International Journal of Computer Mathematics, Vol: 83, No:8- 9, 685–694, 2006.
2. G.M. Amiraliyev, F. Erdogan, A finite difference scheme for a class of singularly perturbed initial value problems for delay differential equations, Numerical Algorithms, Vol: 52, 663- 675, 2009.

TYPE	AUTHORS	TITLES	PAGE	TOPICS
Oral	Muhammad Imran, Brett Allen McKinney , Saira Batoo	Mathematical modeling and control analysis for the Whooping Cough using SVEITR compartmental Model	1	Computational Epidemiology
Online	İlayda Tarhanacı, Elif Başkaya	On A Heat Equation And Newton- Raphson Method	2	Applied Mathematics
Online	Cebraill Saka, Ahmet Öteleş	Eigenvalues Of Complex Skew-Symmetric Circulant Matrices	3	Algebra and Its Applications
Oral	Özlem Kırıcı, Hasan Bulut, Melik Şenyuva	Investigation Of Dynamics Of Magnets Through The Heisenberg Ferromagnetic Spin Chain Equation	4	Applied Mathematics
Online	Yavuz Dinç, Özlem Akyapı	Lower Bound Estimates for the Blow-up Time of the pLaplacian Equation	5	Applied Mathematics
Online	Yavuz Dinç, İrem Üzmez	Nonexistence of Global Solutions for the Hyperbolic Type Equation	6	Applied Mathematics
Oral	Memik Alper Aslan, Kemal Özgen	Investigation of Middle School Students' Real-Life Problem Posing Processes	7	Mathematics Education and Its Applications
Online	Melike Karta	Numerical Solutions Of The Generalized Regular Long Wave (GrLw) Equation Using Four Different Operator Splitting Algorithms Combined With The Finite Element Method	8	Applied Mathematics
Oral	Nuri Murat Yağmurlu, Ali Sercan Karakaş, Selçuk Kutluay	The Newest Numerical Investigation Of The Equal-Width Wave Equation With Quintic Hermite B-Spline Basis Functions	9	Applied Mathematics
Oral	Selçuk Kutluay, Ali Sercan Karakaş, Nuri Murat Yağmurlu	A Robust numerical investigation of the MEW equation with FEM based on Hermite B-spline basis functions	10	Applied Mathematics
Online	Merve İlkhan Kara	On New Paranormed Sequence Spaces	11	Analysis and Its Applications
Oral	Dilara Altan Koç, Gül Gözde Biçer Şarlak, Hasan Bulut	Wave Dynamics In The Lonngren Equation: Exact Solutions And Analysis	12	Applied Mathematics
Oral	Gül Gözde Biçer Şarlak, Özkan Koç, Sibel Paşalı Atmaca, Hasan Bulut	Exploring Nonlinear Wave Behavior In The Strain Equation	13	Applied Mathematics
Online	Sinem Tarsuslu Yılmaz	Intuitionistic Fuzzy Be-Algebras W.R.T. (T, St)-Norm	14	Algebra and Its Applications
Online	Mitat Uysal, Aynur Uysal, Elif Erçelik	Migrating Birds Optimization-Based Radial Basis Function Method To Find the Optimal Surface from a Given Data Set	15	Applied Mathematics
Online	Şeyma Firdevs Hızal, Hasan Bulut, Abdullahi Yusuf	M-Truncated Derivative Applied To The Nonlinear Model With (N+1)-Dimension	16	Numerical Methods and Scientific Programming
Oral	Ulviye Demirbilek, Hasan Bulut	Chaotic Dynamics of the Fractional Improved mKdV Equation via Atangana Conformable Derivative	17	Applied Mathematics

Online	Artion Kashuri, Rozana Liko	Novel Tempered Fractional Integral Inequalities Of Hermite-Hadamard-Mercer Type In The Frame Of Multiplicative Calculus	18	Applied Mathematics
Oral	Mehrdad Lakestani, Fevzi Erdogan	Numerical Solution Of Singularly Perturbed Initial Value Problems With Delay Using B-Spline Wavelets	19	Applied Mathematics
Online	Hazal Yüksekaya	Well-Posedness And Decay Of Solutions for The Porous System with time delay	20	Applied Mathematics
Online	Hazal Yüksekaya	Blow Up of Solutions For The Delayed Beam Equation	21	Applied Mathematics
Oral	Esin İlhan, Adel Sufyan Hussain, Hasan Bulut	Machine Learning And Classical Estimation Methods For Extreme-Value Generated Log-Logistic Distributions: Properties And Performance Analysis	22	Applied Mathematics
Oral	Aslı Alkan, Hasan Bulut, Ercan Çelik	The Novel Numerical Solutions For Time fractional Fishers Equation	23	Fractional Calculus and Applications
Oral	İlker Küçüköğlü, Hacer Özden Ayna, Aysun Yurttaş Güneş, İsmail Naci Cangül	A New Topological Index For Supplier Selection In Procurement Plans	24	Applied Mathematics
Oral	Mücahit Aybas, Yeşim Akbulut	An Optimal Coefficient Control Problem For A Linear Ordinary Differential Equation	25	Control Theory
Oral	Abdullah Kaplan, Zafer Bekiryazici	Analysis Of Win-Probabilities In Turkish style Rummikub Tile Game From A Statistical Perspective	26	Applied Mathematics
Online	Sadullah Bulut, Muhammed Yiğider	Numerical Solution of the Buckley-Leverett Equation with $\mathbb{R}$ -Conformable Fractional Derivative Using Artificial Neural Networks	27	Applied Mathematics
Oral	Gülay Oğuz, Ercan Çelik	On The Structures Of Soft Ideals Over Dirings	28	Topology and Its Application
Oral	Muhammed Fatih Şenocak, Fatma Sağsöz	Coefficient Inequalities For A New Subclass Of Bi-Univalent Functions Defined By Gegenbauer Polynomials	29	Analysis and Its Applications
Oral	Gülay Oğuz, Hüseyin Manap, Ahmet Taş	An Introduction to soft epigroups	30	Topology and Its Application
Online	Muhammed Burak Gul, Emrah Yilmaz	Proportional Tumor Growth Model On Time Scales	31	Fractional Calculus and Applications
Online	Esra Ünal Yılmaz, Yeşim Sağlam Özkan	Exact solutions and modulation instability analysis of the $(3+1)$ -dimensional Gross-Pitaevskii equation with periodic potential	32	Applied Mathematics
Online	Alperen Kızılay, Atakan Tuğkan Yakut	The New Modified Saban Frame And The Evolution Of Curves Based On Modified Saban Frame On $S^{1,2}$ And $H^{2,0}$	33	Geometry and Its Applications
Online	Latifa Agamalieva, Yusif Gasimov	Some Problems For The Eigen values Of The Elliptic Operators	34	Control Theory
Oral	Safar Irandoust-pakchin, Somaiyeh Abdi-mazraeh	Some new numerical methods for solving fractional stochastic differential equations	35	Stochastic and Delay Differential Equations
Online	Aleyna Damar, Hıfı Altınok	Generalized $\mathbb{R}$ -Statistical Convergence Of Order $\mathbb{R}$	36	Analysis and Its Applications

Online	Mithat Kasap, Hıfı Altınok	Some Properties Of The Sequence Class	37	Analysis and Its Applications
Online	Seyfettin Vural, Veyis Turut	Numerical Comparisons for solving systems of equation of Emden–Fowler type	38	Applied Mathematics
Oral	Ali Algan, Faruk Düşünceli	Numerical Solution of One-Dimensional Wave Equation Using Artificial Neural Networks	39	Applied Mathematics
Online	Hayatem Hamal	Parametric Extension of New Generalization of SzászMirakjan Operators	40	Analysis and Its Applications
Oral	Aslıhan Dilek,Nuray Altınöz,Faruk Düşünceli	Numerical Solution of Time–Space Fractional–Order Elliptic Partial Differential Equations Using Artificial Neural Network	41	Applied Mathematics
Online	Hilal Şengür, Hatice Aslan	Degree Of Approximation By Matrix Means Of Hexagonal Fourier Series	42	Analysis and Its Applications
Oral	Taylan Demir, Fahd Jarad	Fractional Model of Human Metapneumovirus (HMPV)	43	Fractional Calculus and Applications
Online	Mehmet Cengiz Fidancı, Yasin Yazlık, Murad Khan Hassani	Stability Analysis Of A Three-Dimensional System Of Difference Equations With Quadratic Terms	44	Applied Mathematics
Online	Ümmühan Yıldız, Emrah Yılmaz	Supply And Demand Model With Proportional Derivatives On Time Scales	45	Analysis and Its Applications
Oral	Fatma Almaz	On The Special Surfaces In Semi-Euclidean 4-Space	46	Geometry and Its Applications
Oral	Muhteşem Demir, Erhan Pişkin	Growth Of Solutions For The Kirchhoff Type Equation	47	Applied Mathematics
Oral	Ayşe Demirhan, Erhan Pişkin	Existence Of Solutions For The Hyperbolic Type Equation	48	Applied Mathematics
Oral	Güldar Efe, Erhan Pişkin	Local Existence Of Solutions For A Delayed Logarithmic Wave Equation	49	Applied Mathematics
Oral	Nebi Yılmaz	Nonexistence Of Global Solutions For The Logarithmic Kirchhoff–Type Hyperbolic Equation With Variable Exponent	50	Applied Mathematics
Oral	Hakan Kutlu, Aynur Yalçın	Error and Erasure Correction Using Parikh Matrices of Uniformly Parikh–Friendly Words	51	Algebra and Its Applications
Online	Gülden Kapusuz, Ahmet Şahiner, Nurullah Yılmaz	A New Approach To The l_p ($0 < p \leq 1$) Minimization Problem With Tight Frame And it is Image Processing Application	52	Analysis and Its Applications
Online	Eyüp Buğra Yalçın, Mehmet Çağrı Yılmaz, Mikail Et	On Pell Calculus	53	Analysis and Its Applications
Online	Süleyman Şengül	The Element-Free Galerkin Method for Solving the Fractional Schrödinger Equation	54	Applied Mathematics
Online	Mustafa Cemil Bişgin	Some New Sequence Spaces Derived By The Composition Of Binomial Matrix And Quintet Band Matrix	55	Analysis and Its Applications
Online	Mustafa Cemil Bişgin	New Sequence Spaces Defined By The Domain Of Sextuple Band Matrix	56	Analysis and Its Applications
Oral	Pakize Uygun, Mehmet Atçeken, Tuğba Mert	Some Special Condition For Pseudosymmetric Kenmotsu Manifolds	57	Geometry and Its Applications
Online	Ahmet Şahiner, Sena Ülgen Çetinkaya	Increasing Fruit Productivity with Modern Modeling Methods	58	Analysis and Its Applications

Online	Yunus Aydogdu, Resat Yilmazer	Fractional Solutions to the k-Hypergeometric Differential Equation	59	Applied Mathematics
Oral	İnci Gültekin, Yasemin Taşyurdu, Hiranur Karaca	The Gaussian Pell Numbers Via Permanents and Determinant of Tridiagonal Matrices	60	Algebra and Its Applications
Oral	Ümmügölsüm Günay	The Approximation of Fixed Point of Endriched Φ -Contraction Mappings	61	Analysis and Its Applications
Oral	Canıl Talantbekova, Ercan Çelik	Estimating The Air Pollution Value of Bishkek City With Arima Using Time Series Method	62	Statistics and Its Applications
Online	Oktay Şahin, Seydi Battal Gazi Karakoç	Numerical Investigations of the Sawada-Kotera Eqaution with Septic B-spline Functions	63	Applied Mathematics
Oral	Ugur Bayrakci, Seyma Tuluçe Demiray, Huseyin Yıldırım	New soliton solutions with modified generalized exponential rational function method	64	Applied Mathematics
Online	Aslı Öner, Sertac Goktas, Yusuf Gurefe	Application of Extended Weierstrass Transform Method to Beta Time Differential Equations	65	Applied Mathematics
Online	Ajay Kumar, Hacı Mehmet Baskonus	Dynamical behavior of analytical soliton solutions for nonlinear partial differential equation	66	Applied Mathematics
Online	Ümmügölsün Çağlayan, İbrahim Gökcan, Ali Hikmet Değer	Lorentz Matrix Multiplication: Exploring Its Relationship With Fibonacci and Lucas Numbers	67	Algebra and Its Applications
Oral	Ayşe Fidan, Erhan Pişkin	Decay And Blow Up of Solutions for a Weighted Biharmonic Equation	68	Applied Mathematics
Oral	Nursemin Kas, Tolga Akturk	Modified Exponential Function Method for Traveling Wave Solutions of the Nonlinear Mathematical Model	69	Applied Mathematics
Oral	Haydar Alıcı	Solutions of Two Exponential Type Potentials for Diatomic Molecules	70	Mathematical Chemistry
Oral	Haydar Alıcı	Explicit Representations for a Class of Gauss Hypergeometric Functions	71	Applied Mathematics
Online	Selahattin Aslan	Tool Frame of Robot End-Effector Using Two Ruled Surfaces Corresponding to the Hyper-Dual Curve	72	Geometry and Its Applications
Online	Muhammet Emin Aydın	Generalized Lacunary Statistical Convergence of Modulus Function Sequences	73	Topology and Its Application
Online	Meltem Öğrenmiş	Multiplicative Equiaffine Curves and Their Properties	74	Geometry and Its Applications
Online	Meltem Öğrenmiş	Characterization of Plane Curves With Constant Multiplicative Equiaffine Curvature	75	Geometry and Its Applications
Online	Çiğdem Süsoy Atalar, Tefik Şahin	Fractional Derivatives: Definitions and Properties	76	Geometry and Its Applications
Oral	Süheyla Elmas, Ferdî Çınar	On Brief Note Regarding F-Contractions In Complete S-Metric Spaces	77	Analysis and Its Applications
Oral	Süheyla Elmas, Aleyna Özel	A Study on Wardowski Contraction In A-Metric Spaces	78	Analysis and Its Applications
Oral	Sezgin Akbulut, İsa Yıldırım, Büşra Melike Kör	Investigation of Applications of Fixed Point Iteration Methods With The Aid of Green's Function	79	Analysis and Its Applications
Online	Berat Süer, Zehra Yücedag	Existence Result of Weak Solution For A Singular kirch hof type Problem With Steklov Boundary Condition	80	Applied Mathematics

Oral	M. Mustafa Beydağı, İlhan İçen, A. Fatih Özcan	Use of Rough Set Theory In Data Mining	81	Topology and Its Application
Oral	Orhan Amirov, Mesut Karabacak	Solution of An Optimization Problem Using Grover's Algorithm For Qutrit Systems	82	Numerical Methods and Scientific Programming
Oral	Ercan Tunç	Oscillation results of second-order noncanonical neutral differential equations with distributed deviating arguments	83	Applied Mathematics
Online	Kübra Tunc, Alper Osman Öğrenmiş	Curves And Their Properties In Dynamic Geometry	84	Geometry and Its Applications
Online	Yusuf Özdemir, Mustafa Yeneroğlu	On The Manganetic Curves According To The Ribbon Frame In Minkowski Space	85	Geometry and Its Applications
Oral	Merve Özkan	Fixed Point Theorems For Four-Point Contractions	86	Analysis and Its Applications
Oral	Semih Geçen, A. Fatih Özcan, İlhan İçen	Soft Set Theory And Various Application Areas	87	Topology and Its Application
Oral	Özlem Öcal, Durmuş Yarımpabuç	Elastic Analysis Of Rotating Fg Pressure Vessels Using The Power Series Method	88	Applied Mathematics
Online	Canan Ödemiş, Abdullah Akkurt	Ostrowski-Gruss Type Inequalities Via Katugampula Fractional Integrals	89	Fractional Calculus and Applications
Online	Ercan Tunç, Adil Kaymaz	A Study On Oscillation Of Neutral Differential Equations With Damping Term	90	Applied Mathematics
Oral	Kevser Soytürk, Mustafa Gezek, Mahmut Ergüt	Unital Designs Admitting Blocking Sets	91	Algebra and Its Applications
Oral	Orhan Özdemir, Ercan Tunç	On The Oscillation Of Fourth-Order Neutral Differential Equations	92	Applied Mathematics
Oral	Hülya Altınok, İsa Yıldırım	On Finite Families of Hemiccontractive Mappings	93	Analysis and Its Applications
Online	Amine Mahzum, Mehmet Çitil	Some Applications of Fuzzy Logic Using Fuzzy Data	94	Applied Sciences
Oral	Himmet Korkmaz, Ayhan Çinici Ali İhsan Mut	A Comparative Analysis Of The Problem solving Processes Of Prospective Elementary Mathematics And Science Teachers In The Context Of A Real-Life Problem	95	Mathematics Education and Its Applications
Oral	Özlem Kırıcı, Arif Özkul, Hasan Bulut	A Collection Of Solitary Waves For The Fractional Kadomtsev-Petviashvili modified-Equal-Width Equation	96	Applied Sciences
Oral	Derya Deniz, Ebru Cavlak Aslan	Various Optical Soliton Solutions for Combined Model Using Analytical Technique	97	Applied Mathematics
Online	Sultan Ceritli, Münevver Yıldırım Yılmaz	Pseudo Null And Null Smarandache Curves In E^3 And Their Characterizations	98	Geometry and Its Applications
Online	Gamze Yaprak, Münevver Yıldırım Yılmaz	Generalized V-Mannheim Curves In E^4 And Their Characterizations	99	Geometry and Its Applications
Oral	Ali İhsan Mut, Muhammed Ahmet Pekyen, Himmet Korkmaz	Analysis Of The Triangles In The Context Of Prototypicality In The Türkiye Century Maarif Model Mathematics Textbooks	100	Mathematics Education and Its Applications
Online	Gül Sinem Keleş, Birol Altın	A Note On Demi Weak Almost Limited Operators	101	Analysis and Its Applications

Oral	Bedrettin Savaş, Efkan Çatıker, Temel Öztürk, Abdullah Karanfil	Synthesis and Characterization of a Series of Poly(2-hydroxypropyl acrylate)-based Copolymers	102	Engineering Sciences
Online	Beytullah Yağız, Erdal Eker, Sinan Deniz, Yener Altun	The Performance Advantage Of The Levygto Hybrid Algorithm Compared To Gto	103	Statistics and Its Applications
Online	Yusuf Gürsoy, Nilüfer Topsakal, Figen Kangalgil	Mathematical Modeling Of Crimeancongo Hemorrhagic Fever Spread: An Approach Based On The Sird Model	104	Applied Mathematics
Oral	Yener Altun, Şakir İşleyen	A Neutral System Approach To Asymptotic Stability Of Q-Fractional Singular Delay Systems	105	Applied Mathematics
Online	Duygu Yıldız, Merve Kara, Yasin Yazlık	Global Stability Of An Exponential Difference Equation System	106	Applied Mathematics
Online	Meral Süer, Mehmet Şirin Sezgin	Ideals and Gröbner Bases Of Some Pseudosymmetric Numerical Semigroup Families	107	Algebra and Its Applications
Online	Berfin Karadağlı, Saliha Erboğa, Nurettin Bağırılmaz	The Analysis of Lgs Success Via Rough Set Theory: The Case Of Siverek And Savur	108	Algebra and Its Applications
Online	Hasan Kandemir, Erhan Pişkin	Global Existence, Decay and Nonexistence of Solutions For Lamé Type Equation With a Fractional Time Delay Term	109	Applied Mathematics
Online	Ali Yılmaz, Abdullah Akkurt, Hüseyin Yıldırım	Fractional Hadamard's Type Inequalities for Exponentially Co-Ordinated Convex Functions	110	Fractional Calculus and Applications
Oral	Özden Taşdemir, Muhammed Emin Kula, Ömer Gündoğdu	Evaluation of Biomechanical and Physiological Effects of Vehicle Vibrations on Pregnant Drivers	111	Engineering Sciences
Online	Mehmet Sait Alakuş, Sedat İlhan	On Some Perfect Numerical Semigroups	112	Algebra and Its Applications
Online	Sinan Deniz, Beytullah Yağız, Yener Altun, Murat Canayaz	A Comparative Study of Iterative and Noniterative Optimization Methods For Nonlinear Least Squares Problems	113	Statistics and Its Applications
Oral	Fahreddin Melih Dursun, Ali Öz	Strength Performance of Alkali Activated Mortars Produced with Waste Brick Powder Depending on NS/NH Ratio and Curing Temperature	114	Engineering Sciences
Online	Kübra Elif Akbaş, Mahmut Işık	On Asymptotically Equivalent Sequences in Probability	115	Statistics and Its Applications
Online	Ayşegül Kızıllarlan, Ayşegül Çetinkaya	k-Lauricella Hypergeometric Functions	116	Applied Mathematics
Online	Ali İhsan Mut, Şeyda Bircan	An Analysis of 9th Grade Mathematics Textbook Prepared In Accordance With Türkiye Century Maarif Model Curriculum In The Context of Congruence and Similarity	117	Mathematics Education and Its Applications
Online	Dila Karataş, Fulya Yörük Dere	Existence and Uniqueness of Solutions For Fractional Boundary Value Problems Involving Caputo-Hadamard Fractional Derivatives	118	Analysis and Its Applications
Oral	İbrahim Boubacr Mounkaila, Gökhan Kaplan	An Experimental Study on the Durability of One-Part Geopolymer Foam Concrete Under NaCl Effect	119	Engineering Sciences
Oral	Nejla Gürefe	Conceptual Perceptions of Prospective Teachers Regarding the Concept of Unit in Mathematics	120	Mathematics Education and Its Applications

Online	Zozan Oktan, Cemil Tunç	On The Boundedness And Stability Properties of Solutions For Fourth Order Nonlinear Stochastic Differential Equations With Variable Delays	121	Applied Mathematics
Online	Zozan Oktan, Cemil Tunç	On The Stability of Fifth-Order Nonlinear Stochastic Delay Differential Equations	122	Applied Mathematics
Oral	Koray İbrahim Atabey, Mikail Et, Muhammed Çınar	Pell Statistical Convergence	123	Analysis and Its Applications
Oral	Adem Irmak, Arzu Aykut, Merve Özer, Sıdıka Şule Şener Kılıç	Numerical Calculations Using Newmark's Method to Solve a Hyperbolic Problem	124	Applied Mathematics
Online	Ghassan Mohammed Abdullah Alomaisi, Nurettin Bağırılmaz	On The Separation Axiom of Topological Spacen M-Space and Tm-Space	125	Topology and Its Application
Online	Elif Yaşar	Analyzing a Nonlinear Schrödinger Type Equation: Dynamical Analysis and Solitary Wave Features	126	Applied Mathematics
Oral	Ahu Ercan, Zehra Özcanoğlu	Euler Method In Numerical Solution of Fractional Differential Equations	127	Applied Mathematics
Oral	Fatma Sağsöz	Second Hankel Determinant for a Subclass of Bi-Univalent Functions Associated With Gegenbauer Polynomials and Subordination	128	Analysis and Its Applications
Oral	Bera Aydın, İsmail Yıldız	Production of Foam Concrete with Hasankale Pumice and Expanded Micronized Perlite	129	Engineering Sciences
Oral	Yasin Kaya	On Generalized Proportional Fractional Integral	130	Analysis and Its Applications
Oral	Oğuzhan Demirel, Durmuş Yarımpabuç	Stress Analysis of a Rotating Fg Polar Orthotropic Disk	131	Applied Mathematics
Oral	Özlem Cerit, Durmuş Yarımpabuç	Deformation Analysis of Fg Longitudinal Rods By Pseudospectral Chebyshev Method	132	Applied Mathematics
Oral	Gazal Okşas, Ali Yıldırım, Durmuş Yarımpabuç	Thermal Efficiency of Fgm Annular Fins	133	Applied Mathematics
Oral	Kerem Kaçar, Durmuş Yarımpabuç	Thermal Analysis of Fg Porous Fin By Adomian Decomposition Method	134	Applied Mathematics
Oral	Zuhal Bilir, Mehmet Ali Sabihaoğlu, Durmuş Yarımpabuç	Application of Inverse Laplace Transform Methods on Heat Conduction Equation	135	Applied Mathematics
Oral	Barış Yapışkan	Pfaffian Formulation of Nonlinear Differential Equations	136	Applied Mathematics
Oral	Esra Pamukçu	Robustness 2.0: A Transformational Story Between Artificial Intelligence and Statistics	137	Artificial Intelligence
Online	Muhittin Evren Aydın, Nesibe Tüzün Yılmaz	On Gaussian Curvature of the Cyclic Surfaces	138	Applied Mathematics
Oral	Sibel Özer, Yusuf Uçar, Halil Beyazıt	Numerical Solution of Grlw Equation By Cubic Hermite B-Spline Collocation Method	139	Applied Mathematics

Online	Sedat Karadağ, Senem Şahan Vahaplar, M. Hakkı Ersoy, Ahmet Feyzioğlu	Development of a Domain-Specific AI Platform Based On Small Language Models	140	Artificial Intelligence
Oral	Yıldırım Demir, Çetin Görür, Saniye Demir	Investigation of Wind Directions By Circular Data Analysis	141	Statistics and Its Applications
Online	Zelal Temel	Finite Difference Method for Linear Volterra Integro-Differential Equations	142	Applied Mathematics
Online	Okan Dardağan, Hakan Altunay	Multi-layer Artificial Neural Networks in Optimization of Predictive Maintenance Processes	143	Artificial Intelligence
Oral	Erkan Günerhan, Hatıra Günerhan, Ercan Çelik	Analytical Solution of a Caputo Fractional-Order Computer Virus Propagation Model	144	Applied Mathematics
Online	Zelal Temel	Numerical Approach of Singularly Perturbed Delay Differential Equations With Nonlocal Conditions	145	Applied Mathematics
Online	Sezer Erdem, Serkan Demiriz	On the Domain of 4-dimensional Catalan Matrix in the Space of Absolutely Summable Double Sequences	146	Analysis and Its Applications
Oral	Hatıra Günerhan	An Analytical Study of a Fractional Differential Model for COVID-19	147	Applied Mathematics
Oral	Hatıra Günerhan	Analytical Solution of a Mathematical Model for the Spread of the Zika Virus	148	Applied Mathematics
Online	Arzu Akbulut	Applications of bifurcation analysis, sensitivity analysis and modified Kudryashov method	149	Applied Mathematics
Oral	Ayşe Nur Akkilic, Hasan Bulut, Zulqurnain Sabir	An artificial neural network procedure for the nonlinear typhoid fever disease system	150	Artificial Intelligence
Online	Osman Tufan, Okan Dardağan, Hakan Altunay	Prediction of Concrete Compressive Strength Using Mamdani-Type Fuzzy Inference System	151	Artificial Intelligence
Online	Ayşe Metin Karakaş, Aslıhan Demir, Sinan Çalık	Topp-Leone Power Garima Distribution: Properties and Applications	152	Statistics and Its Applications
Oral	Begüm Sağlam, Elif Nisa Sert, Zeliha Geçimli, Mazlum Gezer, Halis Yılmaz	Travelling-wave Solutions of Evolution Equations	153	Applied Mathematics
Online	Mine Babaoğlu	Some Spectral Problems Of Diffusion Operator With High Order Approximations	154	Applied Mathematics
Online	Ayşe Metin Karakaş, Fatma Bulut, Murat Karakaş	New Lindley Distribution and Applications	155	Statistics and Its Applications
Oral	Cemil İnan	Comparison of Solution Methods In Nonlinear Equations With One Unknown	156	Applied Mathematics
Oral	Cemil İnan	Investigation of Iterative Approximate Solutions of Linear Equation Systems	157	Applied Mathematics
Oral	Sevda Ozdemir Calikusu, Fevzi Erdogan	Regime-Based Modeling of USD/TL via Stochastic Differential Equations and Change Point Estimation	158	Financial Mathematics
Oral	Erkan Sancar, Erhan Pişkin	Blow Up of Solutions for a Fractional Petrovsky Equation	159	Applied Mathematics
Oral	Tümer Akbayır	Used in Financial Time Series Mathematical Technical Indicators Foundations	160	Financial Mathematics

Oral	İremnur Yüzer Boynikoğlu, Handan Öztekin	The Curves With Modified Orthogonal Frame of in The Minkowski 3-Space	161	Geometry and Its Applications
Oral	Hakan Faik Er, Ekrem Kadioğlu	A Look at The Analysis of Coefficients of Univalent Functions with Artificial Intelligence	162	Analysis and Its Applications
Online	Yasemin Demirel, Oya Mert Coşkun	Determining The Shortest Route For Air Transport After A Possible Fire in Muğla Province Using Artificial Bee Colony Algorithm	163	Applied Mathematics
Online	Sibel Şehriban Ataş	Optical solutions for Fokas system in optical fiber	164	Applied Mathematics
Oral	Davood Ahmadian , Vahid Norouzi , Fevzi Erdogan, Sevda Ozdemir Calikusu	Hybrid Jump-Heston and Bidirectional LSTM Model with Attraction Mechanism for S&P 500 Forecasting	165	Financial Mathematics
Online	Murad khan Hassani	On a solvable difference equations system of second order its solutions are related to a generalized Mersenne sequence	166	Applied Mathematics
Oral	Ümmügülsüm Günay, Murat Özdemir	The Approximation of Fixed Point of Endriched Φ -Contraction Mappings	167	Analysis and Its Applications
Online	Kübra Elif Akbaş, Mahmut Işık	On Asymptotically Equivalent Sequences in Probability	168	Analysis and Its Applications
Oral	Yasemin Taşyurdu, Beyza Aydan	Generalized Bivariate Fibonacci and Lucas Polynomials by Hessenberg Matrices	169	Algebra and Its Applications
Oral	Yasemin Taşyurdu, Beyza Aydan	Bi-Periodic Generalized Bivariate Fibonacci Polynomials	170	Algebra and Its Applications
Oral	İrem Akbulut Arık, Seda İğret Araz	Epidemiological Modeling with Seasonal Environmental Effects	171	Applied Mathematics
Oral	Seda İğret Araz	Investigating Crossover Behaviors in Chaotic Systems	172	Applied Mathematics
Oral	Bilgesu Çifçi, Samet Gündüz	Examining The Golden Ratio-Aesthetic Relationship in Historical Bridges in Diyarbakır	173	Geometry and Its Applications

Mathematical modeling and control analysis for the Whooping Cough using SVEITR compartmental Model

Muhammad Imran¹, Brett Allen McKinney², Saira Batool³

¹ Tandy School of Computer Science, The University of Tulsa, USA

muhammad-imran@utulsa.edu,
brett-mckinney@utulsa.edu,
saira-batool@utulsa.edu,

Abstract

Whooping cough, or pertussis, is a highly contagious disease that poses serious health risks to people of all ages, particularly infants and young children worldwide [1,2]. Given its significant impact, there is a need for a mathematical model to predict the disease's future dynamics and provide strategies for the best controls and elimination [1,3]. In this study, we developed a new susceptible-vaccinated-exposed-infected-treatment-recovered (SVEITR) model for whooping cough to predict its spread and recommend effective strategies for disease control. We demonstrate that the model is well-posed, with positive and bounded solutions within a feasible region. Using the Next-Generation Matrix approach, we computed the basic reproduction number and analyzed the stability of equilibrium points.

Our research also aimed to evaluate the effectiveness of different pharmaceutical and non-pharmaceutical interventions for controlling disease. We expanded the standard SEIR model by incorporating vaccination and treatment compartments, allowing us to assess pharmaceutical (vaccination and treatment) and non-pharmaceutical (self-precaution and transmission reduction) measures. Numerical simulations revealed that all control strategies significantly influence the disease dynamics. Increasing the vaccination and treatment rates led to a reduction in exposed and infected individuals. Notably, combining pharmaceutical and non-pharmaceutical controls proved to be the most effective strategy for managing whooping cough. All the theoretical results and control analysis are validated through numerical simulations.

Keywords: Whooping cough; SVEITR model; Stability and sensitivity analysis; Pharmaceutical and non-pharmaceutical; Control strategies.

REFERENCES

1. AIK. Butt, Atangana-Baleanu Fractional Dynamics of Predictive Whooping Cough Model with Optimal Control Analysis, Symmetry 2023, 15, 1773.
2. Suleman, M. An optimal control of vaccination applied to whooping cough model. J. Math. 2019, 51, 121–136.
3. Ameri, K., Cooper, K.D. A Network-Based compartmental model for the spread of whooping cough in Nebraska. AMIA Summits Transl. Sci. Proc. 2019, 2019, 388.

ON A HEAT EQUATION AND NEWTON- RAPHSON METHOD

İlayda Tarhanacı¹, Elif Başkaya²

^{1,2}Department of Mathematics, Karadeniz Technical University, Trabzon, Turkey

432131@ogr.ktu.edu.tr,

elifbekar@ktu.edu.tr

Abstract

In this paper, we study a heat conduction problem in a rod that a boundary condition, which involves a linear combination of dependent variable and its derivative, arises when heat is lost from the end of the rod due to radiation into the surrounding medium. When we apply the separation of variables method to solve the problem, Newton- Raphson method is used to calculate the eigenvalues of the equations we encounter.

Keywords: Heat equation; Eigenvalue problem; Newton- Raphson method.

REFERENCES

1. Larry C. Andrews, Elementary Partial Differential Equations with Boundary Value Problems, Academic Press Inc., London, 1986.
2. Rao V. Dukkipati, Numerical Methods, New Age International Publishers, New Delhi, 2010.

EIGENVALUES OF COMPLEX SKEW-SYMMETRIC CIRCULANT MATRICES

Cebrail Saka¹, Ahmet Öteleş²

¹Siverek Karacadağ Anatolian High School, Ministry of Education, Şanlıurfa, Turkey

²Department of Mathematics, Faculty of Education, Dicle University, Diyarbakır, Turkey

c.saka63@gmail.com, aoteles@dicle.edu.tr

Abstract

Skew circulant matrix, which is a family of circulant matrices, has many applications such as in graph theory, mechanics, mathematical chemistry, signal processing, coding theory and image processing, etc. They arise in applications involving the discrete Fourier transform and the study of cyclic codes for error correction. They also play a crucial role for solving various differential equations. Numerical solutions of certain types of elliptic and parabolic partial differential equations with periodic boundary conditions often involve linear systems associated with circulant matrices

In this paper, we derive the eigenvalues of any complex skew-symmetric circulant matrix, depending on the Chebyshev polynomials of the first kind. Then, some illustrative examples are given by using maple software, one of computer algebra systems (CAS).

Keywords: Skew circulant matrix; Eigenvalues; Chebyshev polynomial.

REFERENCES

1. N. L. Tsitsas, E.G. Alivizatos, G.H. Kalogeropoulos, A recursive algorithm for the inversion of matrices with circulant blocks, *Applied Mathematics and Computation*. 188, 877-894, 2007.
2. G. Zhao, The improved nonsingularity on the r-circulant matrices in signal processing, in: *International Conference On Computer Technology and Development-ICCTD*, Kota Kinabalu, 564-567, 2009.
3. W. Zhao, The inverse problem of anti-circulant matrices in signal processing, in: *Pacific-Asia Conference on Knowledge Engineering and Software Engineering-KESE*, Shenzhen, 47-50, 2009.

INVESTIGATION OF DYNAMICS OF MAGNETS THROUGH THE HEISENBERG FERROMAGNETIC SPIN CHAIN EQUATION

Özlem Kırıcı¹, Hasan Bulut², Melik Şenyuva²

¹Department of Mathematics, Kırklareli University, Türkiye

²Department of Mathematics, Fırat University, Elazığ, Türkiye

ozlem.isik@klu.edu.tr, hbulut@firat.edu.tr, mlksnyv@gmail.com

Abstract

This paper is related with the Heisenberg ferromagnetic spin chain equation (HFSCE) that has a significant task in theory of magnets, spintronics, plasma physics, and condensed matter physics. The $(m + 1/G')$ -expansion method has been successfully utilized to investigate novel wave structures of the aforementioned equation. This approach has provided a series of new exact wave solutions with diverse physical structures. Various wave patterns have been also observed through the 3D graphs and contour plots. This search into the wave propagation may encourage to examine the nonlinear properties of magnetic materials.

Keywords: $(m + 1/G')$ -expansion method; Heisenberg ferromagnetic spin chain.

REFERENCES

1. Latha, M.M., Vasanthi, C.C., An integrable model of $(2+ 1)$ -dimensional Heisenberg ferromagnetic spin chain and soliton excitations. Phys. Scr. 89(6), 065204 (2014).
2. Devnath, S., Khatun, M.M., Akbar, M.A., Analytical solutions and soliton behaviors in the space fractional Heisenberg ferromagnetic spin chain equation. Partial Differ. Equ. Appl. Math. 11, 100783 (2024).
3. Ma, Y.L., Li, B.Q., Fu, Y.Y., A series of the solutions for the Heisenberg ferromagnetic spin chain equation. Math. Method. Appl. Sci. 41(9), 3316-3322 (2018).
4. Sulaiman, T.A., Aktürk, T., Bulut, H., Baskonus, H.M., Investigation of various soliton solutions to the Heisenberg ferromagnetic spin chain equation. J. Electromagn. Waves Appl. 32(9), 1093-1105 (2018).
5. Bashar, M.H., Islam, S.R., Kumar, D., Construction of traveling wave solutions of the $(2+ 1)$ -dimensional Heisenberg ferromagnetic spin chain equation. Partial Differ. Equ. Appl. Math. 4, 100040 (2021).
6. Osman, M.S., Tariq, K.U., Bekir, A., Elmoasry, A., Elazab, N. S., Younis, M., Abdel-Aty, M., Investigation of soliton solutions with different wave structures to the $(2+ 1)$ -dimensional Heisenberg ferromagnetic spin chain equation. Commun. Theor. Phys. 72(3), 035002 (2020).

Lower Bound Estimates for the Blow-up Time of the p-Laplacian Equation

Yavuz Dinç¹, Özlem Akyapı²

¹ Department of Business Faculty of Economic and Administrative Sciences, Mardin Artuklu University, Mardin, Turkey,

² The Institute for Graduate Educational studies, Department of Mathematics, Mardin Artuklu University, Mardin, Turkey,

yavuzdinc@artuklu.edu.tr, ozlmakyp@gmail.com,

Abstract

In this presentation, we deal with the p-Laplacian wave equation with damping terms in a bounded domain. The study investigates the behavior of solutions and focuses on obtaining lower bounds for the blow-up time under suitable conditions. These conditions are derived based on the nonlinearity of the equation and the properties of the damping terms. Our result extends the recent results obtained by Baghaei (2017) and Zhou (2015), specifically for $p > 2$, and provides new insights into the blow-up dynamics of the p-Laplacian wave equation with damping.

Keywords: Blow-up, Damping term, p-Laplacian equation.

REFERENCES

1. Erhan Pişkin, Yavuz Dinç, Cemil Tunç, Lower bounds for the blow up time to a coupled nonlinear hyperbolic type equations, Communications in Advanced Mathematical Sciences, 3 (1) (2020), 53–56.
2. Khadijeh Baghaei, Lower bounds for the blow-up time in a superlinear hyperbolic equation with linear damping term, Computers and Mathematics with Applications, 73 (2017), 560–564.
3. Salim A. Messaoudi, Ala A. Talahmeh, Jamal H. Al-Shail, Nonlinear damped wave equation: Existence and blow-up. Comp. Math. Appl. 74 (2017) 3024-3041.

Nonexistence of Global Solutions for the Hyperbolic Type Equation

Yavuz Dinç¹, İrem Üzmez²

¹ Department of Business Faculty of Economic and Administrative Sciences, Mardin Artuklu University, Mardin, Turkey,

² The Institute for Graduate Educational studies, Department of Mathematics, Mardin Artuklu University, Mardin, Turkey.

yavuzdinc@artuklu.edu.tr, iremuzmez@hotmail.com,

Abstract

In this presentation, we consider the Kirchhoff-type equation with a variable exponent, a significant equation in the context of evolution problems. The Kirchhoff-type equation belongs to the broader class of evolution equations, which are partial differential equations with time t as one of the independent variables. These types of equations are crucial in modeling various dynamic processes over time. The Kirchhoff equation, in particular, has found wide applications in many areas of applied science, such as the modeling of electrorheological fluid flows, thin liquid films, and other complex systems that exhibit time-dependent behavior. In this work, we focus on proving the upper bound for the blow-up time under suitable conditions, providing a deeper understanding of the time evolution of such systems. This result has important implications for the analysis of stability and the prevention of catastrophic blow-up phenomena in these models.

Keywords: Blow up, Kirchhoff-type equation, Variable exponent.

REFERENCES

1. Erhan Pişkin, Finite time blow up of solutions of the Kirchhoff-type equation with variable exponents. Int. J. Nonlinear Anal. Appl. 11 (2020) 37-45.
2. Yavuz Dinç, Erhan Pişkin, Cemil Tunç, Upper bounds for the blow up time for the Kirchhoff-type equation. Commun.Fac.Sci.Univ.Ank.Ser. A1 Math. Stat. Vol. 72, No. 2 (2023) 352-362.
3. Salim A. Messaoudi, Ala A. Talahmeh, Jamal H. Al-Shail, Nonlinear damped wave equation: Existence and blow-up. Comp. Math. Appl. 74 (2017) 3024 3041.

INVESTIGATION of MIDDLE SCHOOL STUDENTS' REAL-LIFE PROBLEM POSING PROCESSES

Memik Alper Aslan ¹ and Kemal Özgen ²

¹ Master's student at Dicle University, Diyarbakır, Turkey

23978009@ogr.dicle.edu.tr

² Department of Mathematics Education, University of Dicle, Diyarbakır, Turkey

kemal.ozgen@dicle.edu.tr

Abstract

This study aimed to examine the problem-posing processes of middle school students within real-life contexts. The research analyzes students' ability to relate problems to real-life situations and how they integrate real-life elements into their problem-posing processes (Chen & Cai, 2020; Palm, 2006). A teaching experiment was employed as a qualitative research method, and the study was conducted with four eighth-grade students (Cobb, 2000). During the data collection process, students posed a total of four problems-two during the problem-posing phase and two during the lesson extension phase of two problem-posing-based learning task. A two-stage analysis process was carried out. In the first stage, the problems were examined in terms of contextuality, solvability, data quantity, and grammar, to determine their status as valid problems (Kwek, 2015; Nedaei et al., 2022; Özgen et al., 2017; Silver & Cai, 1996; Ulusoy, 2023). In the second stage, the problems were analyzed based on components such as event, question, information/data, presentation, solution strategies, circumstances, solution requirements, and purpose (Palm, 2006). The findings revealed how students incorporate authenticity components into their problem-posing processes and highlight areas requiring further development. The evaluation of four tasks showed that three were classified as fully authentic, while one was classified as partially authentic. These results indicated that although students generally construct problems based on real-life contexts, certain deficiencies prevent them from achieving full authenticity. According to the study's results, students' problem-posing performance was at a satisfactory level. Consequently, their ability to connect real-life contexts with problem-posing was generally positive, except for a few components. Additionally, the analysis of the posed problems indicated that this process contributes positively to students' overall learning experiences. In this regard, it was recommended that problem-posing activities be integrated into the mathematics curriculum. Furthermore, students should be encouraged to connect mathematical concepts with real-life situations by drawing on their own experiences or real-world scenarios.

Keywords: Connection with real life, Problem posing, Problem posing tasks.

REFERENCES

1. Chen, T., & Cai, J. (2020). An elementary mathematics teacher learning to teach using problem posing: A case of the distributive property of multiplication over addition. *International Journal of Educational Research*, 102, 101420.
2. Cobb, P. (2000). Conducting teaching experiments in collaboration with teachers. In A. E. Kelly & R. A. Lesh (Eds.), *Handbook of Research Design in Mathematics and Science Education* (pp. 307–333). Erlbaum.
3. Özgen, K., Aydın, M., Geçici, M. E. ve Bayram, B. (2017). Sekizinci sınıf öğrencilerinin problem kurma becerilerinin bazı değişkenler açısından incelenmesi. *Turkish Journal of Computer and Mathematics Education*, 18(2), 323-351.
4. Mahboubeh Nedaei, Farzad Radmehr & Michael Drake (2022) Exploring undergraduate engineering students' mathematical problem-posing: the case of integral-area relationships in integral calculus, *Mathematical Thinking and Learning*, 24 (2), 149-175, DOI: 10.1080/10986065.2020.1858516
5. Palm, T. (2006). Word problems as simulations of real-world situations : A proposed framework. *for the learning of mathematics*, 26(1), 42-47.
6. Silver, E. A., & Cai, J. (1996). An analysis of arithmetic problem posing by middle school students. *Journal for Research in Mathematics Education*, 27(5), 521–539. DOI:10.2307/749846
7. Ulusoy, F. (2023). Değerlendirme aracı olarak problem kurma. In K. Özgen, T. Kar, S. Çenberci, & Y. Zengin (Eds.), *Matematikte Problem Çözme ve Problem Kurma* (pp. 283–305). Pegem Akademi.
8. Kwek, M. L. (2015). Using problem posing as a formative assessment tool. In F. M. Singer, N. Ellerton, & J. Cai (Eds.), *Mathematical Problem Posing. Research in Mathematics Education* (pp. 273–292). Springer, New York, NY. DOI: 10.1007/978-1-4614-6258-3_13

NUMERICAL SOLUTIONS OF THE GENERALIZED REGULAR LONG WAVE (GRLW) EQUATION USING FOUR DIFFERENT OPERATOR SPLITTING ALGORITHMS COMBINED WITH THE FINITE ELEMENT METHOD

Melike KARTA¹

¹Department of Mathematics, University of Ağrı İbrahim Çeçen, Ağrı, Turkey

mkarta@agri.edu.tr

Abstract

The one-dimensional nonlinear generalized regular wave (GRLW) equation with physical boundary conditions will be taken into consideration in this article. This study will employ operator splitting algorithms in conjunction with the finite element-based Galerkin method, which is a practical and highly adaptable method for the numerical solution of the equation. The study's goal is to generate more precise and appealing results. B-splines serve as the foundation for the application of Galerkin methods in this study. Four distinct numerical schemes—Lie-Trotter splitting techniques $L_{\Delta t} = A - B$, $L_{\Delta t} = B - A$ and Strang splitting techniques $S_{\Delta t} = A - B - A$, $S_{\Delta t} = B - A - B$ are utilized for the numerical results obtained throughout the article and it is evaluated which of these methods yields more accurate results. The work take into consideration the algorithm's output that yields the best results. Using the suggested techniques, the study has been put into practice to solve the single solitary wave problem. Error norms have been calculated in this article, and the results have been compared with the solutions discovered by some researchers in the literature to show the precision and appeal of the numerical techniques employed. Furthermore, The current method is shown to be unconditionally stable through the use of the Von Neumann analysis. This study, which will be carried out after the literature review, will create a new and more effective solution method for the (GRLW) equation.

Keywords: Generalized regularized long wave; B-splines; Galerkin method; Operator splitting.

REFERENCES

1. Roshan, T. A, Petrov-galerkin method for solving the generalized regularized long wave (GRLW) equation, Computers and Mathematics with Applications, 63,943–956,2012.
2. Karakoç, S.B.G and Zeybek, H. Solitary-wave solutions of the GRLW equation using septic B-spline collocation method, Appl. Math. Comput. 289, 159–171, 2016.
3. Khalifa, A.K., Raslan, K.R., and Alzubaidi, H.M., A collocation method with cubic B-splines for solving the MRLW equation, J Comput Appl Math., 212 (2), 406–418, 2008.
4. Karakoç, S.B.G., Liguan, M., and Khalid, K.A. Two efficient methods for solving the generalized regularized long wave equation, Applicable Analysis, 101 (13), 4721–4742, 2022.

THE NEWEST NUMERICAL INVESTIGATION OF THE EQUAL-WIDTH WAVE EQUATION WITH QUINTIC HERMITE B-SPLINE BASIS FUNCTIONS

Nuri Murat YAĞMURLU^{1*}, Ali Sercan KARAKAŞ^{2*}, Selçuk KUTLUAY^{3*}

*Department of Mathematics, University of İnönü, Malatya, Türkiye

murat.yagmurlu@inonu.edu.tr¹, 36203614003@ogr.inonu.edu.tr²,
selcuk.kutluay@inonu.edu.tr³

ORCID: 0000-0003-1593-0254¹, 0000-0001-8622-1127², 0000-0001-9610-504X³

ABSTRACT

In this paper, a numerical investigation of the equal-width wave equation by collocation finite element method using Rubin-Graves type linearization based on quintic Hermite B-spline basis functions under homogeneous boundary conditions will be presented. In order to investigate the effectiveness and reliability of the scheme obtained by the present method, five test problems are selected and the results are presented and evaluated. Only one of the test problems has an analytical solution. In the simulation results of this test problem for which an analytical solution is available, the values of the error norms L_2 and L_∞ will be calculated at the same mesh points available in the literature. Since the other four test problems do not have analytical solutions, the results will be tabulated by giving the values of the conservation constants. Another observation in comparing the results is to examine the graphs of the simulation results at certain times. Graphs will be drawn at certain times and presented in the study. The obtained results show the efficiency and accuracy of the presented method and gives a sight for its further usage for other problems.

Keywords: Equal Width Wave Equation; Quintic Hermite B-spline; Finite Element Method.

REFERENCES

1. Arora, S., Kaur, I. (2016). An efficient scheme for numerical solution of Burgers' equation using quintic Hermite interpolating polynomials. *Arab. J. Math.* **5**, 23–34 <https://doi.org/10.1007/s40065-015-0137-6>.
2. Kutluay, S., Yağmurlu, M., & Karakaş, A. S. (2024). A Robust Quintic Hermite Collocation Method for One-Dimensional Heat Conduction Equation. *Journal of Mathematical Sciences and Modelling*, 7(2), 82-89. <https://doi.org/10.33187/jmsm.1475294>

This study is supported by Inonu University Scientific Research Project with project number FDK-2023-3402.

A Robust numerical investigation of the MEW equation with FEM based on Hermite B-spline basis functions

Selçuk KUTLUAY^{1*}, Ali Sercan KARAKAŞ^{2*}, Nuri Murat YAĞMURLU^{3*}

**Department of Mathematics, University of İnönü, Malatya, Türkiye*

selcuk.kutluay@inonu.edu.tr¹ 36203614003@ogr.inonu.edu.tr²
murat.yagmurlu@inonu.edu.tr³

ORCID: 0000-0001-9610-504X¹, 0000-0001-8622-1127², 0000-0003-1593-0254³

ABSTRACT

The problem considered in the paper is the modified equal-width wave equation with Dirichlet-type boundary conditions. A numerical investigation of the considered problem will be presented by means of a powerful new numerical approach based on quintic Hermite B-spline basis functions and collocation finite element method. First, the solution domain will be divided into sub-regions of equal length. A typical element will be selected from these sub-regions and the element equation will be derived on this typical element. Then the scheme of the problem over the whole region will be found and a solvable system of algebraic equations will be obtained by applying the boundary conditions given together with the problem. The program code of the scheme will be written with the help of symbolic programming language MATLAB. In order to investigate the efficiency and reliability of the scheme obtained by applying the numerical method, four test problems will be examined. The results obtained as a result of simulations for various values of spatial and temporal sizes will be presented in the study with the help of tables and graphs. When the results are analyzed, one can clearly see that the method can be easily applied to other ordinary, partial and fractional differential equations.

Keywords: Finite Element Method; Modified Equal-Width Wave Equation; Linearization.

REFERENCES

1. Arora, S., Jain, R. & Kukreja V.K. Solution of Benjamin-Bona-Mahony-Burgers equation using collocation method with quintic Hermite splines, 154 1-16.
<https://doi.org/10.1016/j.apnum.2020.03.015>
2. Kutluay S., Yağmurlu N.M., & Karakaş A.S. (2024) A novel perspective for simulations of the Modified Equal-Width Wave equation by cubic Hermite B-spline collocation method, Wave Motion, 129, 20-31. <https://doi.org/10.1016/j.wavemoti.2024.103342>

This study is supported by Inonu University Scientific Research Project with project number FDK-2023-3402.

ON NEW PARANORMED SEQUENCE SPACES

Merve İlkhan Kara¹

¹Department of Mathematics, Düzce University, Düzce, Türkiye

merveilkhan@duzce.edu.tr,

Abstract

In general, since paranormed spaces are a generalization of normed spaces, they offer the possibility of using classical functional analysis methods in a broader context. In the present study, it is aimed to introduce certain paranormed sequence spaces as the domain of a newly derived matrix in the Maddox spaces. Further, α, β, γ -duals of the resulting spaces are computed and the characterization of some matrix classes are given.

Keywords: Paranormed sequence space; Matrix domain; Dual spaces.

REFERENCES

1. Bilal Altay, Feyzi Başar, Some paranormed sequence spaces of non-absolute type derived by weighted mean, Journal of Mathematical Analysis and Applications, Vol:319, 494–508, 2006.
2. Bilal Altay, Feyzi Başar, Generalization of the sequence space $\ell(p)$ derived by weighted mean, Journal of Mathematical Analysis and Applications, Vol:330, 174–185, 2007.
3. Merve İlkhan, Serkan Demiriz, Emrah Evren Kara, A new paranormed sequence space defined by Euler totient matrix, Karaelmas Science and Engineering Journal, Vol:9, No:2, 277-282, 2019.
4. Hacer Bilgin Ellidokuzoğlu, Serkan Demiriz, Some new paranormed sequence spaces derived by q -second difference matrix, Journal of Mathematical Extension, Vol:17, No:9, 1-18, 2023.

WAVE DYNAMICS IN THE LONNGREN EQUATION: EXACT SOLUTIONS AND ANALYSIS

Dilara ALTAN KOÇ¹ Gül Gözde Biçer Şarlak¹ Hasan BULUT²

¹Department Mathematics, Mugla Sitki Kocman University, Mugla, Turkey,

²Department of Mathematics, Firat University, Elazig, Turkey,

dilaraaltan@mu.edu.tr, gozdebicersarлак@mu.edu.tr, bulut@firat.edu.tr,

Abstract

In this study, we investigate the Lonngren wave equation using the generalized $(m + 1/G')$ -expansion method to obtain exact travelling wave solutions under different parameter conditions. The method is applied to derive new exact solutions, providing insights into the dynamic behavior of the equation. The obtained solutions are verified by direct substitution into the original equation. Additionally, graphical representations, including three-dimensional and contour, plots, illustrate the physical properties of the derived waveforms. The results demonstrate the effectiveness of the generalized expansion method in solving nonlinear partial differential equations and contribute to the theoretical understanding of wave propagation in applied fields.

Keywords: Lonngren wave equation; generalized expansion method; travelling wave solution.

REFERENCES

1. Kırıcı Ö., Aktürk T., Bulut H., Simulation of Wave Solutions of a Mathematical Model Representing Communication Signals, Journal of the Institute of Science and Technology, Vol:11, No:4, 3086-3097, 2021.
2. Altan Koç, D. Kılbitmez, S., Bulut H., The new complex travelling wave solutions of the simplified modified camassa holm equation, Optical and Quantum Electronics, 56: 215, 2024.
3. Alharbi, A., Abdelrahman, M.A.E., Almatraf, M.B., Analytical and numerical investigation for the DMBBM equation, Computer Modeling in Engineering and Science, Vol:122, No:2, 743–756, 2020.
4. Abbasova, Kh.E., The schrodinger operator with growing potential, Advanced Mathematical Models and Applications, Vol:8, No:1, 14–18, 2023.

EXPLORING NONLINEAR WAVE BEHAVIOR IN THE STRAIN EQUATION

Gül Gözde Biçer Şarлак¹ Özkan Koç¹ Sibel Paşalı Atmaca¹ Hasan Bulut²

¹Department Mathematics, Mugla Sitki Kocman University, Mugla, Turkey,

²Department of Mathematics, Firat University, Elazig, Turkey,

gozdebicersarлак@mu.edu.tr, ozkannkoc@gmail.com, sibela@mu.edu.tr,
hbulut@firat.edu.tr,

Abstract

In this study, we analyze the Strain equation and obtain its travelling wave solutions under various parameter conditions using a generalized analytical approach. Exact solutions are derived in different functional forms, including hyperbolic, trigonometric, and rational expressions, providing deeper insights into the nonlinear wave dynamics of the system. To visualize the wave behavior, three-dimensional surface plots and contour diagrams are generated. The results confirm the effectiveness of the proposed method in solving nonlinear partial differential equations and contribute to the understanding of wave propagation phenomena in applied mathematical models.

Keywords: train equation; travelling wave solutions; nonlinear partial differential equations; wave propagation; exact solutions; graphical analysis.

REFERENCES

1. Sarfaraz A., · Badr Saad T. A. · Sara Salem A., Wave Propagation and Soliton Behaviors for the Strain Equation by Using the Sub-ODE Method and Expansion Technique, International Journal of Applied and Computational Mathematics, 10:122, 2024.
2. Ismael, H.F., Baskonus, H.M., Bulut, H., Abundant novel solutions of the conformable Lakshmanan-Porsezian-Daniel model, Discrete and Continuous Dynamical Systems, Vol:14, No:7, 2311–2333, 2021.
3. Alharbi, A., Abdelrahman, M.A.E., Almatraf, M.B., Analytical and numerical investigation for the DMBBM equation, Computer Modeling in Engineering and Science, Vol:122, No:2, 743–756, 2020.
4. Abbasova, Kh.E., The schrodinger operator with growing potential, Advanced Mathematical Models and Applications, Vol:8, No:1, 14–18, 2023.

INTUITIONISTIC FUZZY BE-ALGEBRAS w.r.t. (T, S_T) -NORM

Sinem Tarsuslu (Yılmaz)¹

¹Department of Natural and Mathematical Sciences, Faculty of Engineering, Tarsus University, Tarsus, Turkey

sinemtarsuslu@tarsus.edu.tr,

Abstract

In this study introduces the concept of intuitionistic fuzzy BE-algebra as a generalization of BCK-algebra, with respect to t-norms and t-conorms. It thoroughly investigates the algebraic properties of triangular norm-based intuitionistic fuzzy BE-algebras and explores the characteristics of norm-based α -cuts. The paper also defines and analyzes the notion of intuitionistic fuzzy filters in the context of triangular BE-algebras. A novel concept of t-norm-based intuitionistic fuzzy BE-algebra is introduced and provides a distinct example to illustrate the structure. It examines the algebraic properties of this new framework and analyzes the role of triangular norms in shaping its behavior. Additionally, the concept of a t-norm-based intuitionistic fuzzy filter is formally defined and explored, with theoretical results supported by concrete examples, highlighting the key characteristics of the proposed algebraic structure.

Keywords: Intuitionistic fuzzy sets, intuitionistic fuzzy BE-algebra, t-norms, intuitionistic fuzzy filter.

REFERENCES

1. Atanassov K.T., Intuitionistic fuzzy sets, Fuzzy Sets and Systems, Volume 20, Issue 1, 1986, Pages 87-96. [https://doi.org/10.1016/S0165-0114\(86\)80034-3](https://doi.org/10.1016/S0165-0114(86)80034-3)
2. Dymek G., Walendziak A., Fuzzy Filters of BE – Algebras, Math. Slovaca 63(5), 935-946, 2013. <https://doi.org/10.2478/s12175-013-0145-y>
3. Elnair, E.M., More Results on Intuitionistic Fuzzy Ideals of BE – Algebras, European Journal of Pure and Applied Math., 17(1), 426-434, 2024. <https://doi.org/10.29020/nybg.ejpam.v17i1.5030>
4. Janis V., t-Norm based cuts of intuitionistic fuzzy sets, Information Sciences 180(7), 2010, 1134--1137. <https://doi.org/10.1016/j.ins.2009.11.039>
5. Klement E. P., Mesiar R., Pap E., Triangular Norms, ISBN 978-90-481-5507-1, Springer-Science+Business Media, B.V. 2000. <https://doi.org/10.1007/978-94-015-9540-7>
6. Kim H.S., Kim H.Y., On BE – Algebras, Scientiae Mathematicae Japonicae Online, 1299-1302, e-2006.
7. Mukkamala, S. R., A Course in BE – algebras, Springer Nature Singapore Pte Ltd. 2018. <https://doi.org/10.1007/978-981-10-6838-6>
8. Parveen P.A., Begum M.H.J., Intuitionistic Fuzzy Ideals in BE – Algebras, American Int. J. of Research in Science, Technology, Engineering&Mathematics, 26(1), March-May 2019, 14-18.

Migrating Birds Optimization-Based Radial Basis Function Method To Find the Optimal Surface from a Given Data Set

Mitat Uysal¹, Aynur Uysal², Elif Erçelik³

^{1,2,3}Department of Software Engineering, Doğuş University, Istanbul, Turkey

muysal@dogus.edu.tr, auysal@dogus.edu.tr, eercelik@dogus.edu.tr

Abstract

In this study, we propose a method to determine the optimal surface from a given set of data points using the Radial Basis Function (RBF) method. The accuracy of the RBF interpolating surface largely depends on the choice of its shape parameter, which controls the smoothness and generalization of the surface. To efficiently determine the optimal shape parameter, we apply the Migrating Birds Optimization (MBO) algorithm, a metaheuristic inspired by the V-shaped flight formations of migratory birds. By utilizing MBO, we minimize the error between the original data and the interpolated surface. Our experiments demonstrate that MBO achieves superior performance in determining the optimal parameters when compared to traditional methods.

Keywords: Radial Basis Function; Migrating Birds Optimization; Surface Fitting

REFERENCES

1. Buhmann M.D., Radial Basis Functions: Theory and Implementations, Cambridge University Press, 2003.
2. Duman, E., Uysal, M., Alkaya, A.F., Migrating Birds Optimization: A New Metaheuristic Approach and Its Performance on Quadratic Assignment Problem. Information Sciences, 217:65-77, 2012.
3. Rippa, S., An Algorithm for Selecting a Good Value for the Parameter in Radial Basis Function Interpolation. *Advances in Computational Mathematics*, 11: 193-210, 1999.

M-TRUNCATED DERIVATIVE APPLIED TO THE NONLINEAR MODEL WITH $(n+1)$ -DIMENSION

Şeyma Firdevs Hızal¹, Hasan Bulut¹, Abdullahi Yusuf¹

¹Department of Mathematics, University of Firat, Elazığ, Turkey

sfkorkmaz@firat.edu.tr,

hbulut@firat.edu.tr,

yusufabdullahi@fud.edu.ng

Abstract

In this paper, the optical solitons with M-truncated derivative and other solutions to the $(n + 1)$ -dimensional nonlinear Schrodinger equation (NLSE) with Kerr and power laws nonlinearities are obtained. We apply the undetermined coefficient method to achieve such novel solutions. Study on the new soliton control structure has been found to explain specific physical problems. These findings have been usefully extended to transmitting long-wave and high-power telecommunication systems. The necessary conditions for the existence of these solutions are presented.

Keywords: $(n + 1)$ -dimensional nonlinear Schrodinger equation; Optical solitons; Undetermined coefficient; Kerr law; Power law; M-truncated derivative.

REFERENCES

1. Younis M., Optical solitons in $(n + 1)$ -dimensions with Kerr and power law nonlinearities, Modern Physics Letters B, Vol:31, 1750186, 2017.
2. Sousa J. V. D. C., De Oliveira E. C., A New Truncated M-Fractional Derivative Type unifying Some Fractional Derivative Types with Classical Properties, International Journal of Analysis and Applications, Vol:16, No:1, 83-96, 2018.
3. Yusuf A., Sulaiman T. A., Mirzazadeh M., Hosseini K., M-Truncated Optical Solitons to a Nonlinear Schrödinger Equation Describing the Pulse Propagation through a Two-Mode Optical Fiber, Optical and Quantum Electronics, Vol: 53, 558, 2021.

Chaotic Dynamics of the Fractional Improved mKdV Equation via Atangana Conformable Derivative

Ulviye Demirbilek¹ Hasan Bulut²

¹Department of Mathematics, University of Mersin, Mersin, Turkey

²Department of Mathematics, University of Firat, Elazig, Turkey

udemirbilek@mersin.edu.tr, hbulut@firat.edu.tr

Abstract

This study examines the chaotic dynamics and bifurcation structures of the fractional improved modified Korteweg–de Vries equation, which includes a space-time dispersion term and a conformable fractional derivative of the Atangana type in time. The dynamics of the system are analyzed using phase portraits, bifurcation diagrams, and Lyapunov exponents. The investigation reveals that changes in key parameters can shift the system from periodic regimes to chaotic behavior. This highlights the nonlinear characteristics and memory effects introduced by the fractional derivative operator. The results show that the system displays complex and sensitive behaviors, such as period-doubling cascades and chaotic attractors, which are typical of nonlinear systems of fractional order. These findings enhance our understanding of complex dynamical processes in dispersive media governed by time-fractional models.

Keywords: Bifurcation analysis; Chaos; Lyapunov exponents; Fractional derivative.

REFERENCES

1. M. Alquran, Novel Methods for Exploring Fractional Solitary Waves in the Improved mKdV Equation with Time-Fractional Derivative and Space-Time Dispersion, Romanian Reports in Physics, pp. 11-7, 2024.
2. B. S. T. Alkahtani, Exploration of Lie Symmetry, Bifurcation, Chaos and Exact Solution of the Geophysical KdV Equation, International Journal of Theoretical Physics, Vol: 64, No: 3, pp. 1–20, 2025.
3. A. Elmandouh, Bifurcation, Quasi-Periodic, Chaotic Pattern, and Soliton Solutions to Dual-Mode Gardner Equation, Mathematics, Vol: 13, No: 5, p. 841, 2025.

NOVEL TEMPERED FRACTIONAL INTEGRAL INEQUALITIES OF HERMITE-HADAMARD-MERCER TYPE IN THE FRAME OF MULTIPLICATIVE CALCULUS

Artion Kashuri¹, Rozana Liko²

¹Department of Mathematical Engineering, Polytechnic University of Tirana,
1001 Tirana, Albania

²Department of Mathematics, University Ismail Qemali, 9400 Vlora, Albania

a.kashuri@fimif.edu.al, rozana.liko@univlora.edu.al

Abstract

In this paper, authors generalize Hermite-Hadamard-Mercer type inequalities using tempered fractional integrals within the framework of multiplicative calculus. Some new multiplicative tempered fractional integral inequalities are established via Jensen-Mercer inequality and several special cases are obtained. Moreover, for suitable choices of parameters some known results are recaptured.

Keywords: Hermite-Hadamard inequality; Hermite-Hadamard-Mercer inequality; Jensen-Mercer inequality; Tempered fractional integral operators; Multiplicatively convex function.

REFERENCES

1. Abdul Mateen, Zhiyue Zhang, Serap Özcan, Muhammad Aamir Ali, Generalization of Hermite-Hadamard, trapezoid, and midpoint Mercer type inequalities for fractional integrals in multiplicative calculus, Boundary Value Problems, Vol:2025, No:22, 1-30, 2025.
2. A.M. Mercer, A variant of Jensen's inequality, Journal of Inequalities in Pure and Applied Mathematics, Vol:4, No:4, 1-2, 2003.
3. Thabet Abdeljawad, Michael Grossman, On geometric fractional calculus, Journal of Semigroup Theory and Applications, Vol:2016, No:2, 1-14, 2016.
4. Agamirza E. Bashirov, Emine Misirli Kurpinar, Ali Özyapici, Multiplicative calculus and its applications, Journal of Mathematical Analysis and Applications, Vol:337, No:1, 36-48, 2008.
5. Silvestru Sever Dragomir, Further inequalities for log-convex functions related to Hermite-Hadamard result, Proyecciones Journal of Mathematics, Vol:38, No:2, 267-293, 2019.

NUMERICAL SOLUTION OF SINGULARLY PERTURBED INITIAL VALUE PROBLEMS WITH DELAY USING B-SPLINE WAVELETS

Mehrdad Lakestani^{1,2} and Fevzi Erdogan³

¹Faculty of Engineering, Van Yüzüncü Yıl University, Van, Türkiye

²Department of Applied Mathematics, University of Tabriz, Tabriz, Iran

mehrdadlakestani@yyu.edu.tr

³Department of Econometrics, Van Yüzüncü Yıl University, Van, Türkiye

ferdogan@yyu.edu.tr

Abstract

In this paper, we solve a singularly perturbed initial value problem with delay by using c B-Spline wavelets. The properties of these functions are provided, and by employing the operational matrix of differentiation, a numerical method is generated over some subintervals that reformulates the problem into a system of algebraic equations. This system can be solved to find the approximate solution. Numerical results demonstrate the efficiency of the method.

Keywords: Singularly perturbed problem; B-Spline wavelets; Delay differential equation.

REFERENCES

1. M. Lakestani, M. Dehghan, The solution of a second-order nonlinear differential equation with Neumann boundary conditions using semi-orthogonal B-spline wavelets, International Journal of Computer Mathematics, Vol: 83, No:8- 9, 685–694, 2006.
2. G.M. Amiraliyev, F. Erdogan, A finite difference scheme for a class of singularly perturbed initial value problems for delay differential equations, Numerical Algorithms, Vol: 52, 663-675, 2009.

WELL-POSEDNESS AND DECAY OF SOLUTIONS FOR THE POROUS SYSTEM WITH TIME DELAY

Hazal Yüksekaya¹

¹ Department of Mathematics, University of Hakkari, Turkey

hazalyuksekkaya@hakkari.edu.tr,

Abstract

In this paper, we study the Porous system with time delay. First, we prove the well-posedness results. Then, by using energy method and by constructing a suitable Lyapunov functional, we establish the decay of solutions under suitable conditions. Time delay effects arise in many applications and practical problems such as physical, chemical, biological, thermal and economic phenomena.

Keywords: Well-posedness, Decay, Porous system, Time delay.

REFERENCES

1. R. Datko, J. Lagnese, M.P. Polis, An example on the effect of time delays in boundary feedback stabilization of wave equations, SIAM J. Control Optim, 22, 152-156, 1986.
2. E. Fridman, Introduction to Time-Delay Systems. Birkhäuser Basel, 2014.
3. H. Yüksekaya, & E. Pişkin, Local existence, global existence and decay results of a logarithmic wave equation with delay term. Mathematical Methods in the Applied Sciences, 46(11), 11802-11813, (2023).
4. S. Nicaise, C. Pignotti, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, SIAM J. Control Optim., 45(5), 1561-1585, 2006.

BLOW UP OF SOLUTIONS FOR THE DELAYED BEAM EQUATION

Hazal Yüksekaya¹

¹ Department of Mathematics, University of Hakkari, Turkey

hazalyuksekkaya@hakkari.edu.tr,

Abstract

Time delay effects arise in many applications and practical problems such as physical, chemical, biological, thermal and economic phenomena. In the eighteenth century, the first equations with delay were considered by brothers Leonard Euler and Bernoulli. By A. Myshkis and R. Bellman, systematical study started at the 1940s. Since 1960, there have been appeared many surveys on the subject. In the middle of 1990s, robust control of systems with uncertain delay was started and led to the "delay bloom" in the beginning of the twenty-first century. Time-delay systems are also named systems with aftereffect or dead-time, equations with deviating argument, hereditary systems, or differential-difference equations. They belong to the class of functional differential equations which are infinite-dimensional, as opposed to ordinary differential equations. In this paper, we study the beam equation with time delay. Under suitable conditions, we prove the blow up results for the beam equation.

Keywords: Blow up, Beam equation, Delay.

REFERENCES

1. R. Datko, J. Lagnese, M.P. Polis, An example on the effect of time delays in boundary feedback stabilization of wave equations, SIAM J. Control Optim, 22, 152-156, 1986.
2. E. Fridman, Introduction to Time-Delay Systems. Birkhäuser Basel, 2014.
3. H. Yüksekaya, & E. Pişkin, Local existence, global existence and decay results of a logarithmic wave equation with delay term. Mathematical Methods in the Applied Sciences, 46(11), 11802-11813, (2023).
4. S. Nicaise, C. Pignotti, Stability and instability results of the wave equation with a delay term in the boundary or internal feedbacks, SIAM J. Control Optim., 45(5), 1561-1585, 2006.
5. Richard, J. P. (2003). Time-delay systems: an overview of some recent advances and open problem, Automatica, 39, 1667-1694.

MACHINE LEARNING AND CLASSICAL ESTIMATION METHODS FOR EXTREME-VALUE GENERATED LOG-LOGISTIC DISTRIBUTIONS: PROPERTIES AND PERFORMANCE ANALYSIS

Esin İlhan¹, Adel Sufyan Hussain², Hasan Bulut^{3,4}

¹Faculty of Engineering and Architecture, Kirsehir Ahi Evran University, Kirsehir, Turkey.

² IT Department, Amedi Technical Institutes, University of Duhok Polytechnic, Duhok, Iraq.

³ Department of Mathematics, Firat University, Elazig, Turkey.

⁴ Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan.

eilhan@ahievran.edu.tr, adel.sufyan@dpu.edu.krd, hbulut@firat.edu.tr

Abstract

The development of new probability distribution models is essential for statistical modeling, as it enhances the accuracy of reproducing real-world data patterns. In this study, the Extreme Value Generated (EVG) family derived from the cumulative distribution function (CDF) of the Extreme Value distribution is introduced. This family includes a key distribution referred to as EVGLLog, characterized by four distribution parameters. A comprehensive statistical evaluation of the EVGLLog distribution is conducted, including derivations of its survival functions and moments, as well as methods for generating random samples. This paper investigates two approaches for parameter estimation of the EVGLLog model: Maximum Likelihood Estimation (MLE) and Long Short-Term Memory (LSTM), a machine learning technique, to determine the most effective and stable estimation method. The proposed EVGLLog model is applied to real-world datasets, including: (1) queue times observed in a banking system, and (2) failure data from accelerated life tests on 59 conductors assessed for electromigration failure. The goodness-of-fit of the EVGLLog distribution is evaluated using several statistical metrics, including chi-square test statistics, negative log-likelihood, Akaike Information Criterion (AIC), and Bayesian Information Criterion (BIC). Results show that the EVGLLog distribution provides a superior fit compared to traditional probability distribution models when analyzing real-world data.

Keywords: Reliability function; Extreme value, waiting time; Long Short-Term Memory; Simulation; Maximum Likelihood Estimations.

REFERENCES

1. David Collett, Modeling Survival Data in Medical Research, CRC Press, Chapman and Hall, New York, USA, 2015.
2. German Rodriguez, Parametric Survival Models, Princeton University, Rapport technique, Princeton, 2010.

THE NOVEL NUMERICAL SOLUTIONS FOR TIME-FRACTIONAL FISHERS EQUATION

Aslı ALKAN¹, Hasan BULUT¹, Ercan ÇELİK²

¹Department of Mathematics, Faculty of Science, Fırat University, Elazığ, Turkey,

²Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University,
Bishkek, Kyrgyzstan

alkanasli47@gmail.com, hbulut@firat.edu.tr, ercan.celik@manas.edu.kg,

Abstract

A new method for solving time-fractional partial differential equations (TFPDEs) is proposed in the paper. It is known as the fractional Kamal transform decomposition method. Time-fractional PDEs are approximated using the fractional Kamal transform decomposition method. The Caputo sense is used to define fractional derivatives. Also, MAPLE software displays the 2D and 3D graphs of the solutions to a few nonlinear TFPDEs. FKTDm is a highly effective method, as evidenced by the numerous applications.

Keywords: Fishers Equation, Fractional Kamal Transform, Adomian Polynomial.

References

- [1] R. Hilfer, Applications of fractional calculus in physics, World scientific, 2000.
- [2] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and applications of fractional differential equations, elsevier, 2006.
- [3] K.S. Miller, B. Ross, An introduction to the fractional calculus and fractional differential equations, Wiley, 1993.
- [4] A. Jajarmi, D. Baleanu, A new fractional analysis on the interaction of HIV with CD4+ T-cells, Chaos, Solitons & Fractals. 113 (2018) 221–229. <https://doi.org/10.1016/j.chaos.2018.06.009>.

A NEW TOPOLOGICAL INDEX FOR SUPPLIER SELECTION IN PROCUREMENT PLANS

İlker Küçükoglu¹, Hacer Özden Ayna², Aysun Yurttaş Güneş², İsmail Naci Cangül²

¹ Department of Industrial Engineering, Bursa Uludag University, Bursa, Turkey

² Department of Mathematics, Bursa Uludag University, Bursa, Turkey

ikucukoglu@uludag.edu.tr, hozden@uludag.edu.tr, ayurttas@uludag.edu.tr,
cangul@uludag.edu.tr

Abstract

Supplier selection is one of the challenging operations for the procurement plans when there exist several alternative markets to be selected. Moreover, planning the order of the markets to be visited makes the planning decisions more complex when the transportation cost is additionally considerable for the decision-maker. In this context, the traveling purchaser problem (TPP) as one of the well-known routing problems that fits with the supplier selection problem and routing decision. The TPP involves satisfying a set of product demands from a set of markets. In TPP, the purchaser selects a number of markets from a set of available markets to buy a number of products. In addition to the procurement plan, the purchaser has to decide on the route to visit selected markets. The aim of the problem is to minimize the total procurement and transportation cost of the purchaser. Since TPP has been shown as an NP-Hard problem, several heuristic and exact methods have been introduced in the literature. To solve the problem, one of the critical issues is the selection of the markets to be visited. When the markets to be visited are known, the procurement plan and route plan can be efficiently obtained by either an exact or heuristic approach. Based on the aforementioned motivation, this study introduces a new topological index that presents the affordability of each market regarding the available products at the market and their prices. After the affordability index of each market is determined, the markets with high index value are selected one by one until all demands are satisfied. The performance of the proposed topological index is tested on a well-known TPP problem set by comparing the selection made by the affordability index and the optimum solution of the problem. The results show that the introduced topological index has the potential to make supplier selection near to optimal solution.

Keywords: Graph theory; Topological indices; Traveling purchaser problem.

REFERENCES

1. Daniele Manerba, Renata Mansini, Jorge Riera-Ledesma, The traveling purchaser problem and its variants, European Journal of Operational Research, Vol:259, No:1, 1-18, 2017.
2. T. Ramesh, Traveling Purchaser Problem, Opsearch, Vol: 18, No:1-3, 78-91, 2011.

AN OPTIMAL COEFFICIENT CONTROL PROBLEM FOR A LINEAR ORDINARY DIFFERENTIAL EQUATION

Mücahit Aybas¹ and Yeşim Akbulut²

¹Graduate School of Natural and Applied Sciences, Atatürk University, Erzurum, Turkey

mucahit.aybas18@ogr.atauni.edu.tr,

²Department of Mathematics, Atatürk University, Erzurum, Turkey

ysarac@atauni.edu.tr

Abstract

Optimal control theory plays a crucial role in various fields of science and technology. Inverse problems and control problems related to ordinary differential equations have been extensively studied due to their wide range of applications [1–4]. In this paper, we address an optimal control problem for the Euler-Bernoulli beam equation. Our main objective is to establish the optimal controllability of a coefficient in the equation using a given boundary data. We provide a theoretical framework and establish the conditions under which optimal controllability can be achieved. The results contribute to the broader understanding of structural control and inverse problems in beam dynamics.

Keywords: Differential equations; Optimal control; Euler-Bernoulli beam.

REFERENCES

1. Chen-Hung Huang, A non-linear inverse vibration problem of estimating the time-dependent stiffness coefficients by conjugate gradient method, International Journal for Numerical Methods in Engineering, Vol: 50, 1545-1558, 2001.
2. D. Lesnic, The comparison model method for determining the flexural rigidity of a beam, J. Inv. III-Posed Problems, Vol:18, 577-590, 2010.
3. Tchavdar T. Marinov, Rossitza S. Marinova, Coefficient identification in Euler-Bernoulli equation from over-posed data, Journal of Computational and Applied Mathematics, Vol: 235, 450-459, 2010.
4. Yeşim Saraç, S. Şule Şener, Identification of the transverse distributed load in the Euler-Bernoulli beam equation from boundary measurement, International Journal of Modelling and Optimization, Vol:8, No:1, 1-4, 2018.

ANALYSIS OF WIN-PROBABILITIES IN TURKISH-STYLE RUMMIKUB TILE GAME FROM A STATISTICAL PERSPECTIVE

Abdullah Kaplan^{1,2,*} and Zafer Bekiryazici²

¹Rize Centre Atatürk Secondary School, Ministry of National Education, Rize, Turkey

²Department of Mathematics, Recep Tayyip Erdogan University, Rize, Turkey

abdullah_kaplan24@erdogan.edu.tr

Abstract

In this paper, the Turkish version of the tile game Rummikub is investigated from a probabilistic approach using simulations of the game in MATLAB. Several scenarios for winning the game, such as winning through doubles only or normal wins with sets of tiles, are simulated using an original algorithm and the outcomes are further studied with different ability levels for the program. The results of the simulations are analyzed statistically in terms of the goodness of fit and comparison of win probabilities for different players in each scenario. Chi-squared and Kolmogorov-Smirnov tests, Monte-Carlo simulations and probabilistic analysis of game scenarios are used to perform an in-depth investigation of the simulation results.

Keywords: Goodness of fit; Simulation; Tile game; Estimation; Win ratio; Chi-Squared Test.

REFERENCES

1. Den Hertog, D. and Hulshof, P.B., Solving Rummikub problems by integer linear programming, The Computer Journal, vol. 49, no. 6, pp. 665–669, 2006.
2. Ekmekçi, Ü., Çataltepe, Z., Güngör, G. and Kahya, D.C., EigenBots for the okey social board game, The Computer Journal, vol. 59, no. 9, pp. 1426–1432, 2016.
3. Uygun, İ., Playing the Turkish tile game okey with deep reinforcement learning, Master's thesis, Sosyal Bilimler Enstitüsü, Boğaziçi Üniversitesi, 2019.
4. Rummikub, Rummikub-history, <https://rummikub.com/history/>, Accessed: 19.12.2024.
5. Balakrishnan, N., Voinov, V. and Nikulin, M.S., Chi-squared goodness of fit tests with applications, Academic Press, 2013.

Numerical Solution of the Buckley-Leverett Equation with β –Conformable Fractional Derivative Using Artificial Neural Networks

Sadullah Bulut¹ Muhammed YİĞİDER²

¹⁻²Department of Mathematics, University of Erzurum Technical, Erzurum, Turkey

sadullah.bulut@erzurum.edu.tr myigider@erzurum.edu.tr

Abstract

In this paper, the classical Buckley-Leverett equation, widely used for modeling two-phase flow in porous media, is reformulated using the β -conformable fractional derivative. Replacing the traditional time derivative with a β -conformable derivative introduces a more flexible and generalized modeling framework. The resulting fractional model is approximately solved using a Physics-Informed Neural Network (PINN) approach, which does not require observational data but instead learns the solution based on the underlying physical law. Implemented in PyTorch, the model is trained over randomly sampled collocation points, and the saturation distribution at $t = 0.5$ is successfully visualized. The proposed method offers an effective and extendable alternative for solving a broader class of physical problems involving fractional-order dynamics.

Keywords: Buckley-Leverett equation, β –conformable derivative, fractional calculus, artificial neural network, physics-informed neural network (PINN)

REFERENCES

1. Rodriguez-Torrado, R., Ruiz, P., Cueto-Felgueroso, L., et al. (2022). Physics-informed attention-based neural network for hyperbolic partial differential equations: application to the Buckley–Leverett problem. *Scientific Reports*, 12, 7557.
2. Zang, J., & Yizhuo, L. (2024). Physics-Informed Neural Network for Solving Buckley-Leverett Equation of Plane Radial Flow. In *Proceedings of the International Conference on Computational Methods*.
3. Ma, L., Li, R. X., Zeng, F., Guo, L., & Karniadakis, G. E. (2023). Bi-orthogonal fPINN: A physics-informed neural network method for solving time-dependent stochastic fractional PDEs. *arXiv preprint arXiv:2303.10913*.

ON THE STRUCTURES OF SOFT IDEALS OVER DIRINGS

Gülay OĞUZ¹ and Ercan ÇELİK²

¹Department of Mathematics, Art and Science Faculty, Harran University, Sanliurfa, Turkey

²Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan

gulay.oguz@harran.edu.tr, ercan.celik@manas.edu.kg

Abstract

Molodtsov introduced the theory of soft sets, which serves as a powerful mathematical tool for addressing uncertainties, as it avoids many of the challenges associated with traditional theoretical methods. In this study, we apply the de notion of ideals over dirings to the concept of soft sets, and introduce the concept of soft ideals over dirings. Also, we explore the properties of soft ideals and demonstrate that the structure is preserved under diring epimorphisms. The main objective of this study is to further the theoretical study of soft ideals over dirings.

Keywords: Soft set; Diring; Ideal; Soft Ideal.

REFERENCES

1. Molodtsov, D. (1999). Soft set theory—first results. *Computers and mathematics with applications*, 37(4-5), 19-31.
2. Biyogmam, G. R., Nganou, J. B., Tchamna, S. (2023). An introduction to dirings. *Asian-European Journal of Mathematics*, 16(03), 2350050.
3. Celik, Y., Ekiz, C. (2011). A New View on Soft Rings. *Hacettepe Journal of Mathematics and Statistics*, 40(2), 273-286.
4. Oguz, G. (2020). Idealistic soft topological hyperrings. *Scientific Inquiry and Review*, 4(3), 61-70.
5. Koyuncu, F., Tanay, B. (2016). Some soft algebraic structures. *Journal of new results in science*, 5(10), 38-51.

COEFFICIENT INEQUALITIES FOR A NEW SUBCLASS OF BI-UNIVALENT FUNCTIONS DEFINED BY GEGENBAUER POLYNOMIALS

Muhammed Fatih Şenocak¹ and Fatma Sağsöz²

¹Graduate School of Natural and Applied Sciences, Atatürk University, Erzurum, Turkey

muhammedfatih.senocak18@ogr.atauni.edu.tr,

²Department of Mathematics, Atatürk University, Erzurum, Turkey

faltuntas@atauni.edu.tr

Abstract

The study of bi-univalent functions has recently attracted considerable attention due to their rich geometric structure and applications in complex analysis. In this paper, we introduce and investigate a novel subclass of bi-univalent functions that are defined via Gegenbauer polynomials in the open unit disk U . The primary objective of this study is to obtain estimates for the initial coefficients a_2 and a_3 in the Taylor-Maclaurin expansion of functions belonging to this new subclass. In addition to the coefficient bounds, we also examine the Fekete-Szegő problem concerning this subclass.

Keywords: Analytic functions; Bi-univalent functions; Coefficient estimates; Gegenbauer polynomials; Fekete-Szegő problem.

REFERENCES

1. Isra Al-Shbeil, Abbas Kareem Wanas, Abdelkader Benali, Adriana Cataş, Coefficient bounds for a certain family of bi-univalent functions defined by Gegenbauer polynomials, Hindawi Journal of Mathematics, Vol:2022, 2022
2. Morteza Moslemi, Ahmad Motamednezhad, Coefficient estimates for a subclass of bi-univalent functions associated with the Horadam polynomials and subordination, Palestine Journal of Mathematics, Vol:13, No:4, 647-656, 2024.
3. Waleed Al-Rawashdeh, Fekete-Szegő functional of a subclass of bi-univalent functions associated with Gegenbauer polynomials, European Journal of Pure and Applied Mathematics, Vol:17, No:1, 105-115, 2024

AN INTRODUCTION TO SOFT EPIGROUPS

Gülay OĞUZ¹, Hüseyin MANAP² and Ahmet TAŞ³

^{1,2,3}Department of Mathematics, Art and Science Faculty, Harran University, Sanliurfa, Turkey

gulay.oguz@harran.edu.tr, manaphuseyin63@gmail.com, ahmet.tas_63@hotmail.com

Abstract

This study investigates the intersection of soft set theory and epigroups, proposing the concept of soft epigroups and examining its foundational properties. Additionally, it obtains characterisations related to the family of soft epigroups and establishes the category of soft epigroups. Finally, it describes the concept of soft subepigroups and studies several structural features.

Keywords: Soft set; Epigroup; Subepigroup; Soft Epigroup; Soft Subepigroup.

REFERENCES

1. Molodtsov, D. (1999). Soft set theory—first results. *Computers and mathematics with applications*, 37(4-5), 19-31.
2. Borralho, M., Kinyon, M. (2020). Variants of epigroups and primary conjugacy. *Communications in Algebra*, 48(12), 5465-5473.
3. Liu, J. (2013). Epigroups in which the operation of taking pseudo-inverse is an endomorphism. In *Semigroup Forum*, 87(3), 627-638. Boston: Springer US.
4. Oguz, G., Icen, I., Gursay, M. H. (2019). Actions of soft groups. *Communications Faculty of Sciences University of Ankara Series A1 Mathematics and Statistics*, 68(1), 1163-1174.
5. Shaprynskiĭ, V.Y., Skokov, D. V., Vernikov, B. M. (2016). Special elements of the lattice of epigroup varieties. *Algebra universalis*, 76(1), 1-30.

PROPORTIONAL TUMOR GROWTH MODEL ON TIME SCALES

Muhammed Burak Gul¹, Emrah Yilmaz²

¹Institute of Natural and Applied Sciences, Firat University, Elazig, Türkiye

²Department of Mathematics, Faculty of Science, Firat University, Elazig, Türkiye

burakgul23@outlook.com¹, eyilmaz@firat.edu.tr²

Abstract

In this study, tumor growth model is constructed on some common time scales. Tumor growth model is a mathematical framework used to describe and predict the growth behavior of tumors over time. These models help researchers and medical professionals understand how tumors develop, spread, and respond to treatments. Because of this importance, the tumor growth model was solved on a time scale by including proportional derivatives and some numerical examples were concretized and comparisons were made.

Keywords: Time scales; Proportional derivative; Tumor growth model.

REFERENCES

1. Claret L., Girard P., Hoff P.M., Cutsem E.V., Zuideveld K.P., Jorga K., Fagerberg J., Bruno R., Model-based prediction of phase III overall survival in colorectal cancer on the basis of phase II tumor dynamics. *Journal of Clinical Oncology* 27, 4103–4108 (2009).
2. Barillot E, Calzone L, Hupe P, Vert J-P, Zinovyev A. *Computational systems biology of cancer*. Boca Raton: CRC Press; 2013.
3. D. R. Anderson, S. G. Georgiev, *Conformable dynamic equations on time scales*, CRC Press: United States, (2020).
4. Anyue Yin, Dirk Jan A.R. Moes, Johan G.C. van Hasselt, Jesse J. Swen and Henk-Jan Guchelaar, A Review of Mathematical Models for Tumor Dynamics and Treatment Resistance Evolution of Solid Tumors, *CPT Pharmacometrics Syst. Pharmacol.* (2019) 8, 720–737
5. Hilger, S. (1988). *Ein Makettenkalkl mit Anwendung auf Zentrumsmannigfaltigkeiten*, Ph.D. Thesis, Universtat Wurzburg.

Exact solutions and modulation instability analysis of the (3+1)-dimensional Gross-Pitaevskii equation with periodic potential

Esra Ünal Yılmaz,¹ Yeşim Sağlam Özkan²

¹Department of Mathematics, University of Bolu Abant İzzet Baysal, Bolu, Turkey

²Department of Mathematics, University of Bursa Uludağ, Bursa, Turkey

esraunal@ibu.edu.tr, ysaglam@uludag.edu.tr

Abstract

In this paper we examine the 3D-Gross-Pitaevskii equation, which describes phenomena such as wave propagation. The solitonic wave solutions of the underlying problem are examined using two efficient techniques: the modified auxiliary equation approach and the $\exp(w(\xi))$ – method. This approach yields solutions that are stated as exponential, rational, trigonometric, and hyperbolic functions. To have a better understanding of their dynamic behavior, several of the distinct types of solitons, including dark waves, have also been shown using 3D visuals for varying parameter values. The stability of the obtained results is confirmed by investigating the modulation instability for the governing model.

Keywords: Exponential method; Modified auxiliary equation method; Gross-Pitaevskii equation.

REFERENCES

1. Qawaqneh H., Altalbe A., Bekir A., Tariq K., Investigation of soliton solutions to the truncated M-fractional (3+1)-dimensional Gross-Pitaevskii equation with periodic potential, AIMS Mathematics 2024, Volume 9, Issue 9: 23410-23433.
2. Hosseini, K., Bekir, A., Ansari, R. Exact solutions of nonlinear conformable time-fractional Boussinesq equations using the $\exp(w(\xi))$ –expansion method. Optical and Quantum Electronics, 2017 49(4), 131..
3. Nadia Mahak, Ghazala Akram, The modified auxiliary equation method to investigate solutions of the perturbed nonlinear Schrödinger equation with Kerr law nonlinearity, Optik, Volume 207, 2020, 164467, ISSN 0030-4026.

THE NEW MODIFIED SABAN FRAME AND THE EVOLUTION OF CURVES BASED ON MODIFIED SABAN FRAME ON S_1^2 AND H_0^2

Alperen Kızılay¹, Atakan Tuğkan Yakut²

^{1 2} Department of Mathematics, University of Niğde Ömer Halisdemir, Niğde, Turkey

alperenkizilay@ohu.edu.tr,

Abstract

In this paper, we define a new modified orthogonal Saban frame for timelike and spacelike curves with geodesic curvature on S_1^2 and the Saban frame for hyperbolic curves with geodesic curvature on H_0^2 . We study the evolution of curves depending on the modified orthogonal Sabban frame and we obtain the necessary conditions for the inextensible flow of curves on modified orthogonal Saban frame. Thus this study presents theoretical examples using a modified orthogonal Saban frame based on the intrinsic property of time.

Keywords: Evolution of curve; Inextensible flow; Modified Saban Frame.

REFERENCES

1. Dariush Latifi, Asadollah Razavi, Inextensible Flows of Curves in Minkowskian Space, Advanced Studies in Theoretical Physics, Vol:2 No:16, 761–768, 2008.
2. Takao Sasai, The fundamental theorem of analytic space curves and apparent singularities of Fuchsian differential equations, Tohoku Mathematical Journal, Second Series, Vol:36 No:1 17-24, 1982.
3. Jan Koenderink Solid shape, MIT Press, Cambridge, 1990.
4. Johan Walrave, Curves and surfaces in Minkowski space, Ph.D. thesis, Leuven University, 1995

SOME PROBLEMS FOR THE EIGENVALUES OF THE ELLIPTIC OPERATORS

Latifa Agamalieva^{1,2}, Yusif Gasimov^{1,2}

¹ Azerbaijan University, Jeyhun Hajibeyli str., 71, AZ1007, Baku, Azerbaijan

²Institute for Physical Problems, Baku State University, Z. Khalilov 23, AZ1148, Baku, Azerbaijan

yusif.gasimov@au.edu.az, latifa.aghamaliyeva@au.edu.az

Abstract

We consider some minimization problems for the eigenvalues of the elliptic operators relatively domains. Finding the optimal domain for the considered functionals is reduced to the solution of the variational problem. The equivalency classes of these problems are defined. Consider the following eigenvalue problem

$$-Au = \lambda u, \quad x \in D, \quad (1)$$

$$u(x) = 0, \quad x \in S_D, \quad (2)$$

where A is some elliptic operator, $D \subset R^n$ is a convex bounded domain with smooth boundary S

Keywords: Eigenvalue Problem; Elliptic operator; variational problem.

REFERENCES

1. Niftiyev, A. A., & Gasimov, Y.S. (2004). Control by Boundaries and Eigenvalue Problems with Variable Domains Publishing House of Baku State University. *Baku, Azerbaijan*.
2. Sokołowski J., Zolésio J.-P. (1992). Introduction to Shape Optimization. Shape Sensitivity Analysis. Springer Series in Computational Mathematics, Vol. 16, Springer-Verlag, Berlin, 1992, 250 p.
3. Gasimov, Y.S. (2003). On some properties of the eigenvalues when the domain varies. *Mathematical Physics, Analyses and Geometry*, 10(2), 249-255.

Some new numerical methods for solving fractional stochastic differential equations

Safar Irandoust-pakchin, Somaiyeh Abdi-mazraeh¹

¹Department of Applied Mathematics, Faculty of Mathematics, Statistics and Computer Sciences, University of Tabriz, Tabriz, Iran

s.irandoust@tabrizu.ac.ir, s.abdi_m@tabrizu.ac.ir

Abstract

In this presentation, we introduce some new numerical works to solve fractional stochastic differential equations (FSDE). Most of these methods are based on the construction of operational stochastic, fractional matrices which the FSDEs convert to the algebraic systems that can be solved easily. Additionally, the approach employs thresholding to significantly reduce the computational workload in linear problems. Thresholding involves setting a threshold value to differentiate between relevant and irrelevant data. Then, the method can avoid unnecessary calculations and focus only on the relevant data. Thresholding can help filter out noise or unwanted elements in a matrices, improving the quality of the data. This results in significantly faster and highly efficient problem-solving. The recently developed approaches have revealed remarkable results in terms of both accuracy and efficiency, following a rigorous assessment of its error bounds and convergence analyses. A thorough analyses show that the schemes perform well demonstrates significant potential as a reliable and efficient methods. The findings indicate that the produced methods provide exceptional precision and operates with remarkable efficiency. These results are promising and suggest that these are an effective solutions.

Keywords: B-spline wavelets, flatlet oblique multiwavelets; fractional differential equation.

REFERENCES

1. S. Abdi-mazraeh, S. Irandoust-pakchin, M. Adel, The construction of stochastic operational matrix via sparsity properties using B-spline wavelets to solve fractional stochastic integro-differential equations, International Journal of Dynamics and Control, In press 2025.
- S. Abdi-mazraeh, S. Irandoust-pakchin, M. Adel, Application of flatlet oblique multiwavelets to solve the fractional stochastic integro-differential equations using Galerkin method, Mathematical Methods in the Applied Sciences, 47(11), (2024), 8342-8365.

GENERALIZED λ -STATISTICAL CONVERGENCE OF ORDER α

Aleyna Damar and Hıfı Altınok

Department of Mathematics, Firat University, Elazığ, Turkey

aleynadmr73@gmail.com hifsialtinok@gmail.com

Abstract

In this study, using the generalized difference operator Δ^m , we generalize the concepts of λ -statistical convergence of order α and p -Cesàro summability of order α by a non-decreasing sequence (λ_n) tending to infinity and give some properties of λ -statistical convergence of order α .

Keywords: Statistical convergence; Difference sequence; Cesaro summability.

REFERENCES

1. Connor, J. S. The statistical and strong p -Cesàro convergence of sequences, *Analysis* 8 (1988), 47-63.
2. Çolak, R. Statistical convergence of order α , *Modern Methods in Analysis and Its Applications*, Anamaya Pub., New Delhi, India (2010), 121-129.
3. Et, M. ; Çınar, M. and Karakaş, M. On λ -statistical convergence of order α of sequences of function, *J. Inequal. Appl.* 2013, 2013:204, 8 pp.
4. Mursaleen, M. λ -statistical convergence, *Mathematica Slovaca*, 50 (1) (2000), 111-115.
5. Šalát, T. On statistically convergent sequences of real numbers, *Math. Slovaca* 30 (1980), 139-150.
6. H. Kızmaz, On certain sequence spaces, *Canad. Math. Bull.* 24(2) (1981), 169-176.
7. Zygmund, A. *Trigonometric Series*, Cambridge University Press, Cambridge, UK, 1979.

SOME PROPERTIES OF THE SEQUENCE

CLASS $S_{\theta}^f(\Delta_F, b)$

Mithat KASAP¹ and Hıfı ALTINOK²

¹Department of Accounting, Sirnak University, Sirnak, Turkey

²Department of Mathematics, Firat University, Elazig, Turkey

fdd_mithat@hotmail.com hifsialtinok@gmail.com

Abstract

Lacunary statistical boundedness, which is mostly used to study the general behavior of sequences and operators, is a subject with wide applications in fields such as statistical analysis, functional analysis and summability theory. In the present paper, we generalize the concept of lacunary statistical boundedness of order β using a difference operator Δ for sequences of fuzzy numbers and give some inclusion relations with the help of many examples and figures. Furthermore, we study some properties like solidity, symmetricity, etc.

Keywords: Fuzzy sequence; Statistical boundedness; Lacunary sequence.

REFERENCES

1. Altinok, H. Et, M. and Altin, Y. Lacunary statistical boundedness of order β for sequences of fuzzy numbers, J Intell Fuzzy Syst 35 (2018), 2383—2390
2. Aytar, S. and Pehlivan, S. Statistically monotonic and statistically bounded sequences of fuzzy numbers, Inform. Sci., 176(6) (2006), 734-744.
3. Bhardwaj, VK. and Dhawan, S. f-statistical convergence of order α and strong Cesaro summability of order α with respect to a modulus, J. Inequal. Appl. (2015) 2015:332 DOI 10.1186/s13660-015-0850-x.
4. Çolak, R. Statistical convergence of order α , Modern Methods in Analysis and Its Applications, New Delhi, India: Anamaya Pub, (2010) 121--129.
5. Freedman, A. R., Sember, J. J. and Raphael, M., Some Cesaro-type summability spaces, Proc. Lond. Math. Soc. 37(3) (1978), 508-520.
6. Matloka, M. Sequences of fuzzy numbers, BUSEFAL 28 (1986) 28-37.

Numerical Comparisons for solving systems of equation of Emden–Fowler type

Seyfettin VURAL¹, Veyis TURUT²

¹Batman University, Instutity of Graduate Education, Batman, Türkiye.

²Batman University, Department of Mathematics, Faculty of Arts and Sciences, Batman, Türkiye.

vrl.seyfettin@gmail.com

veyisturut@gmail.com

Abstract

In this paper, we obtain some new numerical solutions to the linear and nonlinear systems of equation of Emden–Fowler type which seems in the fluid mechanics, relativistic mechanics and in the study of chemically reacting systems by using variational iteration method and Padé approximation.

Keywords: systems of equations; Emden–Fowler equation; variational iteration method, Padé approximation.

REFERENCES

1. G. Adomian, R. Rach, and N.T. Shawagfeh, On the analytic solution of Lane–Emden equation, *Found. Phys. Lett.* 8(2) (1995), pp. 161–181.
2. S. Chandrasekhar, *Introduction to the Study of Stellar Structure*, Dover Publications, NewYork, 1967.
3. H.T. Davis, *Introduction to Nonlinear Differential and Integral Equations*, Dover Publications, NewYork, 1962.
4. M. Dehghan and F. Shakeri, Approximate solution of a differential equation arising in astrophysics using the variational iteration method, *New Astron.* 13 (2008), pp. 53–59.
5. J.H. He, A variational iteration approach to nonlinear problems and its applications, *Mech. Appl.* 20(1) (1998), pp. 30–31.
6. J.R. Muatjetjeja and C.M. Khalique, First integrals for a generalized coupled Lane–Emden system, *Nonlinear Anal. RealWorld Appl.* 12 (2011), pp. 1202–1212.
7. K. Parand, M. Shahini, and M. Dehghan, Rational Legendre pseudospectral approach for solving nonlinear differential equations of Lane–Emden type, *J. Comput. Phys.* 228 (2009), pp. 8830–8840.
8. A.M. Wazwaz The variational iteration method for solving systems of equations of Emden–Fowler type, *International Journal of Computer Mathematics*, 88(16) (2011), pp. 3406–33415.
9. A.M. Wazwaz, A new method for solving differential equations of the Lane–Emden type, *Appl. Math. Comput.* 118(2/3) (2001), pp. 287–310.

Numerical Solution of One-Dimensional Wave Equation Using Artificial Neural Networks

Ali ALGAN¹, Faruk DÜŞÜNCELİ²

¹Department of Mathematics, The Institute for Graduate Educational Studies, Mardin Artuklu University, Türkiye

²Department of Economics, Mardin Artuklu University, Türkiye

math.alialgan@gmail.com, farukdusunceli@artuklu.edu.tr

Abstract

In this study, the authors developed an efficient solution to find the numerical solution of the 1D wave equation with the help of Artificial Neural Networks (ANN). Numerical solution of the wave equation by this method is a complex and computationally intensive process. The ANN method separates from all these processes and provides a faster numerical solution with very little error. In addition, the ANN method visualizes the solution. In ANN method, activation functions are used to increase the capabilities of the model and systematize the learning processes. Fibonacci polynomials are used as activation functions in the learning process of the network since they have a continuous and complex structure in themselves. Thus, the numerical solution of the one-dimensional wave equation with certain initial and boundary values is obtained by ANN method.

Keywords: Wave Equation; Fibonacci Polynomial; Artificial Neural Networks.

REFERENCES

1. Ihsan, A. F. (2021). On the Neural Network Solution of One-Dimensional Wave Problem. *Jurnal RESTI (Rekayasa Sistem dan Teknologi Informasi)*, 5(6), 1106-1112.
2. Sana, D. (2022). Approximating the wave equation via physics informed neural networks: various forward and inverse problems.
3. Armawanto, F. (2024). Persamaan gelombang satu dimensi menggunakan metode finite difference explicit dan neural network (doctoral dissertation, universitas PGRI Semarang)
4. Birol, F. (2018). Fibonacci, Lucas, Pell, Pell, Pell-Lucas, generalized Pell number sequences and polynomials and their relations with linear groups (Master's thesis, Balikesir University (Turkey)).

Parametric Extension of New Generalization of Szász-Mirakjan Operators

Hayatem Hamal¹

¹ Department of Mathematics, Tripoli University, Tripoli 22131, Libya

hafraj@yahoo.com,

Abstract

In this paper, we introduce a new extension of the generalized Kantorovich Type of Szász-Mirakjan operators based on a new parameter $\alpha > 0$ and $\eta \in \mathbb{R}^+$, as follows:

$$K_{m,\eta}^\alpha(h; y) = \sum_{j=0}^{\infty} S_{m,j}(y) \int_0^1 h\left(\frac{j+s^\eta}{m+\alpha}\right) ds \quad (1.1)$$

$$S_{m,j}(y) = e^{-ny} \frac{(ny)^k}{k!}, \quad (1.2)$$

and new operators are positive for arbitrary functions defined on the interval $[0, 1]$.

We obtain an essential auxiliary result for these new operators. Our subsequent study focuses on their qualitative aspects, including uniform convergence and asymptotic behavior. For $m \in \mathbb{N}$, $\alpha > 0$ and $\eta \in \mathbb{R}^+$, we have the following

$$\left(K_{m,\eta}^\alpha(t-y, y)\right)^2 \leq \frac{2}{(m+\alpha)^2} \gamma_{\eta,\alpha}^1(1+y^2), \quad \gamma_{\eta,\alpha}^1 > 0 \text{ for } y \in [0, \infty) \quad (1.3)$$

$$K_{m,\eta}^\alpha\left((t-y)^2, y\right) \leq \gamma_{\alpha,\eta}^2(m)(1+y)^2, \quad \gamma_{\alpha,\eta}^2(m) > 0 \text{ for } y \in [0, \infty) \quad (1.4)$$

$$K_{m,\eta}^\alpha\left((t-y)^4, y\right) \leq \varphi^*(\eta) \frac{A}{(m+\alpha)^2}, \quad \varphi^*(\eta) > 0, \text{ for } y \in [0, A]. \quad (1.5)$$

Keywords: Szász-Mirakjan operators; Linear positive operators; modulus of continuity.

REFERENCES

1. E. Baytunc., H. Aktuglu and N.I. Mahmudov, ~ A new generalization of Szász-Mirakjan Kantorovich operators for better error estimation, Fundam. J. Math. Appl., 6(4) (2023), 194-210. DOI 10.33401/fujma.1355254.

Numerical Solution of Time-Space Fractional-Order Elliptic Partial Differential Equations Using Artificial Neural Network

Aslıhan Dilek¹, Nuray Altınöz², Faruk Düşünceli³

^{1,2} Department of Mathematics, The Institute for Graduate Educational Studies, Mardin Artuklu University, Türkiye

³ Department of Economics, Mardin Artuklu University, Türkiye,

asliihandlk@gmail.com , nuray.altinz29@gmail.com, farukdusunceli@artuklu.edu.tr

Abstract

Classical numerical methods have been widely used for solving the heat equation. However, in order to obtain more accurate representations of complex physical phenomena, fractional order differential equations have received increasing attention in recent years. In this study, the solution of fractional heat equation is addressed through the application of artificial neural networks (ANN). A feed-forward neural network architecture is used, where the training process is driven by a composite loss function that imposes both the governing differential equation and associated initial and boundary conditions. The accuracy and efficiency of the proposed method are evaluated through comparisons with analytical and traditional numerical solutions.

Keywords: Fractional-Order Elliptic Partial Differential Equations; Heat equation; Artificial neural networks (ANNs); Numerical methods

REFERENCES

1. Rafi, S. A. (2024). *Analysis and construction of artificial neural networks for heat equations and their related parameters, depths, and accuracies* (Master's thesis, University of Arkansas). *Graduate Theses and Dissertations*. <https://scholarworks.uark.edu/etd/5277>
2. Ertuğ, S. (2023). *Damping problem for the fractional heat equation* (Doctoral dissertation, Ankara University). 64 pages.
3. Qu, H., Liu, X., & She, Z. (2020). Neural network method for fractional-order partial differential equations. *Neurocomputing*, 414, 225–237. <https://doi.org/10.1016/j.neucom.2020.07.063>
4. Mall, S., & Chakraverty, S. (2018). *Artificial neural network approach for solving fractional order initial value problems*. Department of Mathematics, National Institute of Technology Rourkela. <https://arxiv.org/abs/1810.04992>

DEGREE OF APPROXIMATION BY MATRIX MEANS OF HEXAGONAL FOURIER SERIES

Hilal Şengür* and Hatice Aslan**

Department of Mathematics, Firat University, Elazığ, Turkey

*sengurhilal7@gmail.com , ** haslan@firat.edu.tr

Abstract

The degree of approximation of the function f , which is periodic with respect to the hexagon lattice is estimated by matrix means $T_n^{(A)}(f)$ of its hexagonal Fourier series in the generalized Hölder metric where A is a lower triangular infinite matrix of nonnegative real numbers with nonincreasing row in this paper.

Keywords: Hexagonal domain, hexagonal Fourier series, generalized Hölder class, matrix mean

REFERENCES

1. J. Bustamante and M.A. Jimenez, *Trends in Hölder Approximation* in: Optimization and Mathematical Economics, Springer, 2001, 81-95
2. L. Condat, D.V. Ville and T. Blu, *Hexagonal versus orthogonal lattices: A new comparison using approximation theory*, IEEE International Conference on Image Processing. (2005).
3. R.A. DeVore and G.G. Lorentz, *Constructive Approximation* .Springer-Verlag, 1993.
4. B. Fuglede, *Commuting self-adjoint partial differential operators and a group theoretic problem*, *Journal of Functional Analysis* **16** (1974), 101-121.
5. A. Guven, *Approximation by means of hexagonal Fourier series in Hölder norms*, *Journal of Classical Analysis* **1** (1) (2012), 43-52.
6. A. Guven, *Approximation properties of hexagonal Fourier series in the generalized Hölder metric*. *Comput. Methods Funct. Theory* **13** (2013) 509-531.

Fractional Model of Human Metapneumovirus (HMPV)

Taylan Demir¹, Fahd Jarad²

¹Department of Mathematics, University of Ankara, Ankara, Turkey

²Department of Mathematics, University of Çankaya, Ankara, Turkey

taylandemir@ankara.edu.tr,

Abstract

Human Metapneumovirus (HMPV) is a major cause of acute respiratory infections, particularly affecting infants and young children, and often resulting in hospitalization. It also poses a considerable risk to elderly individuals and those with weakened immune systems. Gaining a deeper understanding of how this virus spreads is essential, and mathematical modeling proves to be an effective approach for this purpose. In this study, we introduce a novel model that uses fractional differential equations to examine the dynamics of HMPV transmission. These equations allow us to incorporate memory and non-local effects into the framework, offering a more realistic representation of disease progression. The model also reflects the delayed immune response and intricate interactions between the virus and the host. We conduct a thorough stability analysis and perform numerical simulations to explore how different parameters impact the spread of the infection. The findings offer valuable insights for refining public health strategies, including vaccination plans and antiviral treatment schedules, to better manage and contain HMPV outbreaks.

Keywords: Fractional calculus; Human Metapneumovirus (HMPV); Fractional differential equations; Disease modeling; Non-local effects; Epidemiology; Numerical simulations.

REFERENCES

1. Podlubny, I. (1998). *Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications* (Vol. 198). elsevier.
2. Van den Hoogen, B. G., de Jong, J. C., Groen, J., Kuiken, T., de Groot, R., Fouchier, R. A., & Osterhaus, A. D. (2001). A newly discovered human pneumovirus isolated from young children with respiratory tract disease. *Nature medicine*, 7(6), 719-724.
3. Diethelm, K., Ford, N. J., & Freed, A. D. (2002). A predictor-corrector approach for the numerical solution of fractional differential equations. *Nonlinear Dynamics*, 29, 3-22.

STABILITY ANALYSIS OF A THREE-DIMENSIONAL SYSTEM OF DIFFERENCE EQUATIONS WITH QUADRATIC TERMS

Mehmet Cengiz Fidancı, Yasin Yazlık, Murad Khan Hassani

Department of Mathematics, Nevşehir Hacı Bektaş Veli University, Nevşehir, Turkey

16.190man@gmail.com

Abstract

In this study, we analyze the dynamics of a nonlinear three-dimensional system of second-order difference equations involving quadratic terms. We establish the existence of fixed points and explore the properties of positive solutions such as boundedness, persistence, and invariance. We also investigate the local and global asymptotic stability of the equilibrium point and provide the rate of convergence of the solutions. Finally, numerical simulations are given to support and illustrate the theoretical findings.

Keywords: Difference equations; Asymptotic stability; Boundedness; Rate of convergence.

SUPPLY AND DEMAND MODEL WITH PROPORTIONAL DERIVATIVES ON TIME SCALES

Ümmühan YILDIZ¹, Emrah YILMAZ¹

¹ Department of Mathematics, Firat University, Elazig, Türkiye

ummuhan.yldz44@gmail.com, eyilmaz@firat.edu.tr

Abstract

In this study, supply and demand model, which explains how markets in the economy seek balance with their own internal dynamics and how the price mechanism directs resource allocation, will be defined and solved using proportional derivatives on time scales. Some comparisons will also be made with numerical examples.

Keywords: Time Scales, Supply and Demand Model, Proportional Derivative.

REFERENCES

1. Bohner, Martin, and Allan Peterson. Dynamic equations on time scales: An introduction with applications. Springer Science & Business Media, 2001
2. Anderson, Douglas R. and Georgiev, Svetlin G. Conformable dynamic equations on time scales. CRC Press, 2020.
3. Vasicek, O. (1977). An equilibrium characterization of the term structure. Journal of financial economics, 5(2), 177-188.
4. Anderson, Douglas R., Ulness, Darin J. Newly defined conformable derivatives, Advances in Dynamical Systems and Applications, 2015, 10.2: 109-137.
5. Sandeep S. Jaiswal, Economics Theory of Supply and Demand Outdated, Journal of Global Economics, Volume 12:02, 2024

ON THE SPECIAL SURFACES IN SEMI-EUCLIDEAN 4-SPACE

Fatma Almaz

Department of Mathematics, Faculty of Arts and Sciences, University of Batman,

Batman, TÜRKİYE

fatma.almaz@batman.edu.tr,

Abstract

In this work, the special surfaces, in which the surfaces that can be chosen to be tubular surface generated by normal curve in semi-Euclidean 4-space, are examined. Moreover, by using the Gaussian curvatures and mean curvatures of tube surfaces with normal curves generated by Frenet frame in semi-Euclidean 4-space, the conditions being Weingarten surface and HK-quadric surface, minimal surface, at, ... etc are examined in detail.

Keywords: Tubular surfaces, HK-quadric surface, weingarten surfaces, normal curves.

REFERENCES

1. Almaz, F., Kulahcı, M.A., The notes on rotational surfaces in Galilean space, International J. of Geometric Methods in Modern Physic , 18(2), 2021.
2. Almaz, F., Kulahcı, M.A., A Survey on Tube Surfaces in Galilean 3-Space, Journal of Polytechnic, 25(3), 1033-1042, 2022.
3. Ilarslan, K., Nesovic, E., Time-like and Null Normal Curves in Minkowski Space E^3_1 , Indian J. pure appl. Math., 35(7), 881-888, 2004.
4. Kuhnel, W., Differential geometry: curves- surfaces- manifolds. Braunschweig, Wiesbaden, 1999.
5. O'Neill, B., Semi-Riemannian Geometry with Applications to Relativity. Academic Press Inc., London, 1983.
6. Pressley, A., Elementary Differential Geometry, second edition London, UK. Springer-Verlag London Limited, 2010.

GROWTH OF SOLUTIONS FOR THE KIRCHHOFF TYPE EQUATION

Muhteşem Demir¹ and Erhan Pişkin²

¹Institute of Natural and Applied Sciences, Dicle University, Diyarbakır, Turkey

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

demirmuhtesem@gmail.com, episkin@dicle.edu.tr

Abstract

In this presentation, we consider the Kirchhoff type equation with initial and boundary conditions. We prove the growth of solutions under suitable conditions.

Keywords: Growth, Kirchhoff type equation, Parabolic type equation.

REFERENCES

1. E. Pişkin, F. Ekinçi, Blow up and growth of solutions to a viscoelastic parabolic type Kirchhoff equation, *Filomat*, 37(2) (2023) 519-530.
2. E. Pişkin, F. Ekinçi, Blow up, exponential growth of solution for a reaction-diffusion equation with multiple nonlinearities, *Tbilisi Mathematical Journal*, 12(4) 2019, 61-70.
3. L. X. Truong, Y. N. Van, Exponential growth with L^p -norm of solutions for nonlinear heat equations with viscoelastic term, *Applied Mathematics and Computation* 273, (2016) 656-663.

EXISTENCE OF SOLUTIONS FOR THE HYPERBOLIC TYPE EQUATION

Ayşe Demirhan¹ and Erhan Pişkin²

¹Institute of Natural and Applied Sciences, Dicle University, Diyarbakır, Turkey

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

aysedemirhan5@gmail.com, episkin@dicle.edu.tr

Abstract

In this presentation, we consider the hyperbolic type equation with initial and boundary conditions. We prove the existence of solutions under suitable conditions by using the potential well method.

Keywords: Existence, Hyperbolic type equation, Potential well method.

REFERENCES

1. Y. Liu, W. Li, A family of potential wells for a wave equation, Electronic Research Archive, 28(2) (2020) 807-820.
2. E. Pişkin, Sobolev Uzayları, Seçkin Yayıncılık, 2025
3. E. Pişkin, A. Demirhan, On potential wells for a nonlinear higher-order hyperbolic-type equation, International Journal of Open Problems in Computer Science and Mathematics, 17 (2) (2024) 24-37.
4. D.H. Sattinger, On global solution of nonlinear hyperbolic equations, Arch. Rational Mech. Anal., 30 (1968) 148-172.

LOCAL EXISTENCE OF SOLUTIONS FOR A DELAYED LOGARITHMIC WAVE EQUATION

Güldar Efe¹ and Erhan Pişkin²

¹Institute of Natural and Applied Sciences, Dicle University, Diyarbakır, Turkey

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

guldarefe@gmail.com, episkin@dicle.edu.tr

Abstract

In this work, we study a logarithmic viscoelastic wave equation with a time-dependent delay term in a bounded domain. We establish the local existence of solutions using the Faedo-Galerkin method.

Keywords: Existence, Delay, Faedo-Galerkin.

REFERENCES

1. P. Gorka, Logarithmic Klein--Gordon equation, Acta Phys. Pol. B, 40(1) (2009), 59-66.
2. M. Kafini, S. Messaoudi, Local existence and blow up of solutions to a logarithmic nonlinear wave equation with delay, Applicable Analysis, 99(3) (2020) 530-547.
3. E. Pişkin, H. Yüksekaya, Local existence and blow up of solutions for a logarithmic nonlinear viscoelastic wave equation with delay, Computational Methods for Differential Equations, 9(2) (2021) 623-636.

NONEXISTENCE OF GLOBAL SOLUTIONS FOR THE LOGARITHMIC KIRCHHOFF-TYPE HYPERBOLIC EQUATION WITH VARIABLE EXPONENT

Nebi Yılmaz

Sur Anadolu İmam Hatip Lisesi, Diyarbakır, Turkey

nebiyilmaz1981@gmail.com

Abstract

In this work, we investigate the nonexistence of global solutions to a logarithmic Kirchhoff-type hyperbolic equation with variable exponents. We prove that, under appropriate conditions on the initial data, there is nonexistence of global solution.

Keywords: Hyperbolic type equation, Nonexistence, Variable exponent.

REFERENCES

1. E. Pişkin, Sobolev Uzayları, Seçkin Yayıncılık, 2025
2. E. Pişkin, Evolüsyon Denklemlerin Çözümlerinin Patlaması (Blow up of Solutions of Evolution Equations), Pegem Akademi Yayıncılık, 2022.
3. N. Yılmaz, E. Pişkin (2024). Blow up of Solutions for the Logarithmic Higher-Order Kirchhoff-Type Equation with Variable Exponent. Journal of Nepal Mathematical Society, 7(2), 50-57.

Error and Erasure Correction Using Parikh Matrices of Uniformly Parikh-Friendly Words

Hakan Kutlu ¹, Aynur Yalçiner ²

^{1,2}Department of Mathematics, University of Selçuk, Konya, Türkiye

¹ hakan_282_42@hotmail.com ² ayalciner@selcuk.edu.tr

Abstract

The Parikh matrix mapping, a generalization of the Parikh mapping, was introduced by Mateescu et al. [2]. A word's Parikh matrix is an upper triangular matrix that represents the number of occurrences of specific subwords within that word. It's important to note that the Parikh matrix of a word doesn't usually identify that word uniquely. Words that share a common Parikh matrix M are classified as M -equivalent. In [3], the concept of uniformly Parikh-friendly words was introduced. Atanasiu et al. [1] suggested the use of M -equivalent words as codewords. Their results indicated that the associated Parikh matrix allows for the correction of received words containing up to a certain level of erasures and errors.

In this talk, we will consider the uniformly Parikh-friendly words as codewords. We will examine the erasure and error correction capabilities of the relevant Parikh matrix.

Keywords: M -equivalent word; Parikh matrix; uniformly Parikh-friendly word.

REFERENCES

1. Adrian Atanasiu, Ghajendran Poovanandran, Abdalhadi Abu Zeyneh, Wen Chean Teh Erasure and error correcting ability of Parikh matrices, Information Processing Letters, Vol:175, No:106223, 2022.
2. Alexandru Mateescu, Arto Salomaa, Kai Salomaa, Sheng Yu, A sharpening of the Parikh mapping, Theoretical Informatics and Applications, Theoretical Informatics and Applications, Vol:35, 551–564, 2001.
3. Wen Chean Teh, Parikh-friendly permutations and uniformly Parikh-friendly words, Australasian Journal of Combinatorics, Vol:76, No:1, 208–219, 2020.

A NEW APPROACH TO THE l_p ($0 < p \leq 1$) MINIMIZATION PROBLEM WITH TIGHT FRAME AND ITS IMAGE PROCESSING APPLICATION

Gülden Kapusuz¹, Ahmet Şahiner¹, Nurullah Yılmaz¹

¹Department of Mathematics, University of Suleyman Demirel, Isparta, Turkey

guldenkapusuz92@gmail.com

Amidst advancements in signal and image processing, achieving high-quality restorations from limited measurement data has become a significant research topic. Particularly, the need for irregular and non-convex optimization techniques is increasing, highlighting the demand for new and effective algorithms. In this context, studies focusing on sparsification and advanced enhancement methods contribute substantially to the development of signal processing technologies. In this study, the goal is to sparsify a given signal using the concepts of Tight Frame (TF) and Discrete Fourier Transform (DFT), and subsequently restore it through a novel approach developed for the l_p ($0 < p \leq 1$) norm minimization problem. The work specifically addresses the restoration of sparsified black-and-white images. In image restoration, irregular and non-convex regularization techniques offer significant advantages, often transforming restoration problems into large-scale, irregular, and non-convex optimization problems. However, existing minimization approaches have been observed to be ineffective in handling such challenges. To overcome this issue, a Smoothing Conjugate Gradient Method (SCGM) is proposed, combining the classical nonlinear conjugate gradient method with two new-generation smoothing techniques. The proposed method has been formulated as a numerical algorithm and applied to test images with different types of noise. Experimental results demonstrate that the proposed SCGM algorithm outperforms existing methods in terms of restoration performance.

Keywords: Tight Frame, Discrete Fourier Transform, Non-Smooth Optimization, Image Restoration, Conjugate Gradient Method.

REFERENCES

1. Sahiner, A., Yilmaz, N. Double stranded smoothing technique in energy minimization problems. Numerical Algebra, Control and Optimization, 14(1), 80-88, 2024.
2. Sahiner, A., Ibrahem, S. A., Mustafa, A. M. A novel smoothing technique and its application to image processing. 6th International New York Congress on Life and Applied Sciences, 2022.
3. Yilmaz, N., Sahiner, A., Kapusuz, G. Smoothing Conjugate Gradient Method Associated with New Generation Techniques to Solve Image Restoration Problems, ICMME 2024, Nevşehir-Turkey, 226-227, 2024.

On Pell Calculus

Eyüp Buğra YALÇIN¹, Mehmet Çağrı YILMAZER¹, Mikail ET¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

eyupbugra13@gmail.com, m.cagri.yilmazer@gmail.com, mikailet68@gmail.com

Abstract

In this paper, we introduce a new mathematical structure called “Pell calculus”. This original calculus contains establishment of Pell calculus and presentation of fundamental theorems related to derivatives, exponential and some trigonometric functions, binomial, Taylor expansion, integrals and their operational features within this mathematical structure. Our work make contribution to scientific discourse by providing extensive study of presented Pell calculus, introducing strong foundation for further examination in this specific mathematical framework.

Keywords: Pell number; Pell-Lucas Number.

REFERENCES

1. Koshy, T. (2019). *Fibonacci and Lucas Numbers with Applications, Volume 2* (Vol. 2). John Wiley & Sons.
2. Lupas, A. (1999). A guide of Fibonacci and Lucas polynomials. *Octagon mathematics magazine*, 7(1), 2-12.
3. Koshy, T., & Koshy, T. (2014). *Pell and Pell–Lucas Numbers* (pp. 115-149). Springer New York.
4. Yağmur, T. (2024). Non-Newtonian Pell and Pell-Lucas numbers. *Journal of New Results in Science*, 13(1), 22-35.
5. Spreafico, E. V. P., Catarino, P., & Vasco, P. (2023). On hybrid hyper k-pell, k-pell–lucas, and modified k-pell numbers. *Axioms*, 12(11), 1047.
6. Özvatan, M. (2018). *Generalized golden-Fibonacci calculus and applications* (Master's thesis, Izmir Institute of Technology (Turkey)).
7. Kac, V. G., & Cheung, P. (2002). *Quantum calculus* (Vol. 113). New York: Springer.

The Element-Free Galerkin Method for Solving the Fractional Schrödinger Equation

Süleyman Şengül¹

¹Department of Mathematics, University of Recep Tayyip Erdoğan, Rize, Turkey

suleyman.sengul@erdogan.edu.tr,

Abstract

In this paper, the fractional Schrödinger equation with Caputo-type derivatives was numerically solved using the Element-Free Galerkin method. The L_1 method was used to discretize the Caputo fractional derivative, and the Rubin-Graves approach was used to linearize the nonlinear terms. It has been observed that the approximate solutions are consistent with the exact solutions.

Keywords: Caputo fractional derivative, Schrödinger equation, Element free Galerkin method.

REFERENCES

1. Lancaster P.; Sakauskas K. Surfaces generated by moving least squares methods. Math Comput., 37, 141-58, 1981.
2. Belytschko, T.; Lu, Y.Y.; Gu, L. Element-free Galerkin methods.” International Journal for Numerical Methods in Engineering, 37.2, 229–256, 1994.
3. Caputo, M. Linear models of dissipation whose Q is almost frequency independent—II. Geophysical journal international, 13.5, 529-539, 1967.
4. Podlubny, Fractional Differential Equations. An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of their Solution and Some of their Applications. Academic Press, New York, 1999.

SOME NEW SEQUENCE SPACES DERIVED BY THE COMPOSITION OF BINOMIAL MATRIX AND QUINTET BAND MATRIX

Mustafa Cemil Bişgin¹

¹Department of Mathematics, University of Recep Tayyip Erdoğan, Rize, Turkey

mustafa.bisgin@erdogan.edu.tr

Abstract

In this paper, we define three new sequence spaces by using the matrix domain of composition matrix of the Binomial and Quintet band matrices. Also, we examine some inclusion relations between existing sequence spaces and newly defined ones. Moreover, we determine Schauder bases and α -, β - and γ -duals of those sequence spaces. Lastly, we characterize some matrix classes related to new sequence spaces.

Keywords: Matrix Transformations; Matrix Domain; Schauder Basis; Sequence Spaces.

REFERENCES

1. Bişgin, Mustafa Cemil, The binomial sequence spaces of nonabsolute type, Journal of Inequalities and Applications, Vol:2016, No:309, 1–16, 2016.
2. Bişgin, Mustafa Cemil, Four new sequence spaces obtained from the domain of quadruple band matrix operator, Turkish Journal of Mathematics and Computer Science, Vol:15, No:2, 294–311, 2023.
3. Topal, Kübra, New sequence spaces derived by the composition of binomial and quadruple band matrices, Recep Tayyip Erdogan University Journal of Science and Engineering, Vol:5, No:2, 76–92, 2024.

NEW SEQUENCE SPACES DEFINED BY THE DOMAIN OF SEXTUPLE BAND MATRIX

Mustafa Cemil Bişgin¹

¹Department of Mathematics, University of Recep Tayyip Erdoğan, Rize, Turkey

mustafa.bisgin@erdogan.edu.tr,

Abstract

In this work, we construct new sequence spaces by using the domain of sextuple band matrix $S(r, s, t, u, v, y)$ which is the general form of difference, double band, triple band, quadruple band and quintet band matrices. Furthermore, we study some inclusion relations and topological properties related to those spaces. Also, we define Schauder bases and α -, β - and γ -duals some of those spaces. Finally, we determine some matrix classes belonging to new sequence spaces.

Keywords: Matrix Domain; Matrix Transformations; Schauder Basis; Sequence Spaces.

REFERENCES

1. Kızmaz, Hüsnü, On certain sequence spaces, Canadian Mathematical Bulletin, Vol:24, No:2, 169–176, 1981.
2. Kirişçi, Murat, Başar, Feyzi, Some new sequence spaces derived by the domain of generalized difference matrix, Computers and Mathematics with Applications, Vol:60, No:5, 1299–1309, 2010.
3. Sönmez, Abdulcabbar, Some new sequence spaces derived by the domain of the triple band matrix, Computers and Mathematics with Applications, Vol:62, No:2, 641–650, 2011.
4. Bişgin, Mustafa Cemil, Four new sequence spaces obtained from the domain of quadruple band matrix operator, Turkish Journal of Mathematics and Computer Science, Vol:15, No:2, 294–311, 2023.

SOME SPECIAL CONDITION FOR PSEUDOSYMMETRIC KENMOTSU MANIFOLDS

Pakize Uygun¹, Mehmet Atçeken², Tuğba Mert³

¹Department of Mathematics, University of Aksaray, Aksaray, Turkey

pakizeuygun@hotmail.com,

²Department of Mathematics, University of Aksaray, Aksaray, Turkey

mehmetatceken@aksaray.edu.tr,

³Department of Mathematics, University of Sivas Cumhuriyet, Sivas, Turkey

tmert@cumhuriyet.edu.tr,

Abstract

In this paper pseudosymmetric and Ricci pseudosymmetric of a Kenmotsu manifolds are researched. We have achieved the necessary and sufficient conditions for a Kenmotsu manifold, W_8 -pseudosymmetric, W_8 -Ricci pseudosymmetric, W_9 -pseudosymmetric and W_9 -Ricci pseudosymmetric. Additionally, some interesting results on Kenmotsu manifolds are obtained.

Keywords: Kenmotsu Manifold; Pseudosymmetric Manifold; Ricci pseudosymmetric Manifold.

REFERENCES

1. M. Atçeken, On generalized Sasakian space forms satisfying certain conditions on the concircular curvature tensor, Bulletin of Mathematical Analysis and Applications, 6(1)2014, 1-8.
2. D. E. Blair, Contact manifolds in Riemannian geometry, Lecture Note in Mathematics, 509, Springer-Verlag, Berlin-New York, (1976).
3. W. M. Boothby, An Introduction to Differentiable Manifolds and Riemannian Geometry, Academic Press, Inc, London, UK, 1986.
4. E. Boeckx, P. Buecken and L. Vanhecke, ϕ -symmetric contact metric spaces, Glasgow Mathematical Journal, 41 (1999), 409-416.
5. E. Cartan, Sur one classe remarquable despaces de Riemannian, Bulletin de la Societe Mathematique de France, 54 (1926), 214-264.

Increasing Fruit Productivity with Modern Modeling Methods

Ahmet Şahiner¹, Sena Ülgen Çetinkaya¹

¹Department of Mathematics, University of Suleyman Demirel, Isparta, Turkey

senaulgen48@gmail.com

Agricultural productivity in fruit production varies depending on environmental factors such as temperature, precipitation, humidity, and wind, which often involve uncertainty. This uncertainty limits the accuracy of predictions made using classical mathematical methods. In this study, a fuzzy logic modeling approach, one of the modern optimization tools, was used. The model was developed using historical data on temperature, precipitation, and olive yield published by the Turkish Statistical Institute (TURKSTAT). It can produce successful predictions not only based on this data but also on untested datasets. As a result of regression analysis, a prediction accuracy of over 80% was observed with untested data. Due to its ability to represent nonlinear relationships and offer flexible interpretations, the model demonstrated strong performance in terms of generalizability. The results indicate that fuzzy logic can be used as an effective decision support tool in agricultural productivity analysis.

Keywords: Olive Productivity, Fuzzy Logic, Temperature, Precipitation, Agricultural Modeling.

REFERENCES

1. Şahiner A., Uçun F., Kapusuz G., and Yılmaz N. Completed Optimised Structure of Threonine Molecule by Fuzzy Logic Modelling. Zeitschrift für Naturforschung A, 71(4), 381-386. 2016.
2. Şahiner A., Yılmaz N., and Kapusuz G. Descent Global Optimization Method Based on Smoothing Techniques via Bezier Curves. Carpathian Journal of Mathematics, 33(3), 373-380. 2017.
3. Şahiner A., İbrahim S. and İbrahim A. A. Fuzzy Logic Modeling for Prediction of the Nuclear Tracks. Journal of Multidisciplinary Modeling and Optimization, 1(1), 33-40. 2018.
4. Uçar N., Ekinçi C., Şahiner A., Çalık A. and Kapusuz G. Use of Fuzzy Logic Modelling for Radiation Shielding Properties of Borided Fe-Ni Binary Alloys. Süleyman Demirel Üniversitesi Fen Edebiyat Fakültesi Fen Dergisi, 14(2), 276-284., 2019.

Fractional Solutions to the k -Hypergeometric Differential Equation

Yunus Aydogdu¹, Resat Yilmazer¹

¹Department of Mathematics, University of Firat, Elazig, Türkiye

yaydogdu_23@hotmail.com, ryilmazer@firat.edu.tr

Abstract

In this study, the fractional calculus operator was used to examine both homogeneous and non-homogeneous situations of the k -hypergeometric differential equation. We apply certain classical transformations to solve the equation and at the same time constrain the parameters required to determine their value. The major benefit of the fractional operator is that it implements singular differential equations and transforms them into fractional order equations. As a result, several new exact fractional solutions of the given equation were created.

Keywords: k -hypergeometric differential equation; Fractional calculus; Differential equations.

REFERENCES

1. Podlubny I. Fractional Differential equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of their Solution and Some of their Applications. San Diego, California: Elsevier; 1998.
2. Yilmazer, R. N-fractional calculus operator N_m method to a modified hydrogen atom equation. Math. Commun. 15, 489–501, 2010.
3. Li S, Dong Y. k -Hypergeometric series solutions to one type of non-homogeneous k -Hypergeometric Equations. Symmetry, 11(2), 262, 2019.
4. Mubeen S, Naz M, Rehman A, Rahman G. Solutions of k -hypergeometric differential equations. Journal of Applied Mathematics. 2014, 13, 2014.
5. Yilmazer, R., Ali, K.K., Discrete fractional solutions to the k -hypergeometric differential equation, Math Meth Appl Sci, 44:7614–7621, 2021.

THE GAUSSIAN PELL NUMBERS VIA PERMANENTS AND DETERMINANT OF TRIDIAGONAL MATRICES

İnci Gültekin¹ Yasemin Taşyurdu² and Hiranur Karaca³

¹ Department of Mathematics, Atatürk University, Erzurum, Turkey

igultekin@atauni.edu.tr,

² Department of Mathematics, Erzincan University, Erzincan, Turkey

ytasyurdu@erzincan.edu.tr,

³ Graduate School of Natural and Applied Sciences, Atatürk University, Erzurum, Turkey

hiranur.karaca17@ogr.atauni.edu.tr

Abstract

Gaussian number means representation as Complex numbers. Our goal in this work is to give the Gaussian Pell numbers by using permanent and determinant of some tridiagonal matrices.

Keywords: Permanent, Gaussian Pell number, Tridiagonal matrix.

REFERENCES

1. Horadam, A.F. Complex Fibonacci numbers and Fibonacci quaternions. The American Mathematical Monthly, Vol:70, No:3, 289-291, 1963.
2. Jordan, J.H. Gaussian Fibonacci and Lucas numbers. The Fibonacci Quarterly, Vol:3, No:4, 315-318, 1965.
3. Horadam, A.F. Generalized Fibonacci Sequence, The American Mathematical Monthly, Vol:68, No:5, 455-459, 1961.
4. Halıcı, S., Öz, S. On Some Gaussian Pell and Pell –Lucas numbers. Ordu University journal of sciences and technology, Vol:6, No:1, 8-18, 2016.
5. Minc, H., Marcus, M., Permanents, Encyclopedia Math. Appl., Addison–Wesley Publishing Company, London, Vol:6, 1978.

THE APPROXIMATION OF FIXED POINT OF ENRICHED ϕ -CONTRACTION MAPPINGS

Ümmügülsüm Günay¹

¹ Department of Mathematics, University of Ataturk, Erzurum, Turkey

ummugulsum.gunay18@ogr.atauni.edu.tr

Abstract

We introduce the class of enriched ϕ -contractions in Banach spaces as a natural generalization of ϕ -contractions and study the existence and approximation of the fixed points of mappings in this new class, which is shown to be an unsaturated class of mappings in the setting of a Banach space. We illustrated the usefulness of our fixed point results by studying the existence and uniqueness of the solutions of some second order (p,q) -difference equations with integral boundary value conditions.

Keywords: Banach space; enriched ϕ -contraction; enriched cyclic ϕ -contraction; fixed point; Maia type fixed point theorem; (p,q) -difference equation; integral boundary value condition

REFERENCES

1. Banach, S. Sur les opérations dans les ensembles abstraits et leurs applications aux équations intégrales. Fundam. Math. 1922, 3, 133–181. [CrossRef]
2. Agarwal, R.P.; O'Regan, D.; Sahu, D.R. Fixed Point Theory for Lipschitzian-Type Mappings with Application; Topological Fixed Point Theory and Its Applications; Springer: New York, NY, USA, 2009.
3. Agarwal, R.P.; Karapinar, E.; O'Regan, D.; Roldán-López-de-Hierro, A.F. Fixed Point Theory in Metric Type Spaces; Springer: Cham, Switzerland, 2015.
4. Berinde, V. Iterative Approximation of Fixed Points, 2nd ed.; Lecture Notes in Mathematics, 1912; Springer: Berlin, Germany, 2007.
5. Granas, A.; Dugundji, J. Fixed Point Theory; Springer: New York, NY, USA, 2003.
6. Kirk, W.A.; Sims, B. (Eds.) Handbook of Metric Fixed Point Theory; Kluwer Academic Publishers: Dordrecht, The Netherlands, 2001.
7. Kirk, W.; Shahzad, N. Fixed Point Theory in Distance Spaces; Springer: Cham, Switzerland, 2014.
8. Rus, I.A.; Petrusel, A.; Petrusel, G. Fixed Point Theory; Cluj University Press: Cluj-Napoca, Romania, 2008.

ESTIMATING THE AIR POLLUTION VALUE OF BISHKEK CITY WITH ARIMA USING TIME SERIES METHOD

Canıl TALANTBEKOVA and Ercan ÇELİK

Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University,
Bishkek, Kyrgyzstan

2251y03003@manas.edu.kg, ercan.celik@manas.edu.kg

Abstract

This study aims to estimate the air pollution levels in Bishkek, the capital of Kyrgyzstan, using the ARIMA (AutoRegressive Integrated Moving Average) time series model. Air pollution poses a significant threat to public health and environmental sustainability, especially in rapidly developing urban areas like Bishkek. The research utilizes daily PM10 data for the year 2021, along with meteorological variables such as temperature, humidity, wind speed, and precipitation. After conducting data cleaning, outlier detection, and seasonal decomposition, the ARIMA(1,0,1) model was selected as the best-fitting model based on AIC and residual analysis. The results indicate that the model provides reliable short-term forecasts for PM10 concentration levels and effectively captures seasonal pollution trends. This research offers a valuable tool for policymakers and urban planners to implement timely and data-driven interventions to improve air quality in Bishkek.

Keywords: Air Pollution, Time Series Analysis, ARIMA Model, PM10 Forecasting, Bishkek, Environmental Monitoring

REFERENCES

1. Akbostancı, E., Türüt-Aşık, S., & Tunç, G. İ. (2009). The relationship between income and environment in Turkey: Is there an environmental Kuznets curve? *Energy Policy*, 37(3), 861–867. <https://doi.org/10.1016/j.enpol.2008.09.088>
2. Aladağ, E. (2021). Forecasting of particulate matter with a hybrid ARIMA model based on wavelet transformation and seasonal adjustment. *Urban Climate*, 39, 100930. <https://doi.org/10.1016/j.uclim.2021.100930>
3. Aladağ, E. (2023). The influence of meteorological factors on air quality in the province of Van, Turkey. *Water, Air, & Soil Pollution*, 234(4), 259. <https://doi.org/10.1007/s11270-023-06265-0>

Numerical Investigations of the Sawada-Kotera Equation with Septic B-spline Functions

Oktay ŞAHİN¹, Prof. Dr. Seydi Battal Gazi KARAKOÇ²

1 MsC Student, Hacı Bektaş Veli University, Mathematics Department, Nevşehir, 44280, TÜRKİYE

<https://orcid.org/0000-0003-0579-8712>

36203614003@nevsehir.edu.tr, 0.505. 910 24 42

2 Professor, Hacı Bektaş Veli University, Mathematics Department, Nevşehir, 44280, TÜRKİYE

ORCID ID: <https://orcid.org/0000-0002-2348-4170>

sbgkarakoc@nevsehir.edu.tr, 0.505 391 92 00

Abstract

This study uses the B-spline functions as approximate functions to offer a finite element scheme for numerical solutions of the Sawada Kotera (S-K) equation. Additionally, an investigation of the algorithm's Von-Neumann stability has been done [1]. Furthermore, the method's practicality and dependability are illustrated through an analysis of a single soliton's behavior. To regulate the efficiency and conservation features of the proposed algorithm, the I_2 , L_2 and L_∞ error norms as well as the two lowest invariants, I_1 and I_2 , of the equation have been calculated [2]. The aspects of the problem modelled are easily visualized thanks to the tables and images that accompany the obtained numerical results. The outcomes also show that our approach is effective .

Keywords: Sawada Kotera Equation; Finite Element Method; Collocation; Septic B-Spline;

REFERENCES

- [1] Karakoc SBG, Zeybek H. A numerical investigation of the GRLW equation using lumped Galerkin approach with cubic B spline. SpringerPlus 2016;5:1–17.
- [2] Karakoc SBG, Saha A, Sucu D. A novel implementation of Petrov-Galerkin method to shallow water solitary wave pattern and superperiodic traveling wave and its multistability generalized Korteweg-de Vries equation. Chin J Phys 2021;68:605–617

New soliton solutions with modified generalized exponential rational function method

Ugur BAYRAKCI^{1,*}, Seyma TULUCE DEMIRAY¹, Huseyin YILDIRIM²

¹ Department of Mathematics, University of Osmaniye Korkut Ata, Osmaniye, Turkey

² Department of Mathematics, University of Kahramanmaraş Sütçü İmam, Kahramanmaraş, Turkey

*corresponding author

ubayrakci42@gmail.com, seymatuluce@gmail.com, hyildir.ksu.edu.tr

Abstract

In this study, the fractional Kairat-II equation is studied. The modified generalized exponential rational function method is applied to this equation. Thus, various soliton solutions of this equation are obtained. Various values are given to the obtained solutions and graphical drawings are made.

Keywords: Modified generalized exponential rational function method; Fractional Kairat-II equation; Soliton solutions.

REFERENCES

1. W. A. Faridi, G. H. Tipu, Z. Myrzakulova, R. Myrzakulova, S. A. AlQah tani, P. Pathak, The sensitivity demonstration and propagation of hyper geometric soliton waves in plasma physics of Kairat-II equation, *Physica Scripta* 99, 045209, 2024.
2. M. Iqbal, D. Lu, M. Alammari, A. R. Seadawy, N. E. Alsubaie, Z. Umurza khova, R. Myrzakulov, A construction of novel soliton solutions to the nonlinear fractional Kairat-II equation through computational simulation, *Optical and Quantum Electronics* 56, 845, 2024.
3. K. K. Ali, A. Yokus, A. R. Seadawy and R. Yilmazer, The ion sound and Langmuir waves dynamical system via computational modified generalized exponential rational function, *Chaos Solitons Fractals* 161 112381, 2022.
4. S. Duran, H. Durur and A. Yokuş, Traveling wave and general form solutions for the coupled Higgs system, *Mathematical Methods in the Applied Sciences*, 46(8), 8915-8933, 2023.

APPLICATION OF EXTENDED WEIERSTRASS TRANSFORM METHOD TO BETA TIME DIFFERENTIAL EQUATIONS

Aslı Öner¹, Sertac Goktas¹, Yusuf Gurefe¹

¹Department of Mathematics, Mersin University, Mersin, Turkey

aslioner@mersin.edu.tr, sertacgoktas@mersin.edu.tr, yusufgurefe@mersin.edu.tr

Abstract

In this paper, the exact solutions of beta-conformable fractional equations are constructed by using the extended Weierstrass transformation method. This method is applied to the equations widely used in engineering and mathematical physics, providing exact solutions involving trigonometric, Weierstrass elliptic, and rational functions. By applying a traveling wave transformation, the beta-conformable fractional equations are reduced to a nonlinear ordinary differential equation, and the solution function is determined by expanding it into a finite series using the Weierstrass function solutions of the corresponding elliptic differential equation.

Keywords: Beta derivative; The extended Weierstrass transformation method; Weierstrass elliptic function.

REFERENCES

1. Roshdi Khalil, Mohammed Al Horani, Abdelrahman Yousef, Mohammad Sababheh, A New Definition of Fractional Derivative, Journal of Computational and Applied Mathematics, Vol:264, 65–70, 2014.
2. Abdon Atangana, Dumitru Baleanu, Ahmed Alsaedi, New Properties of Conformable Derivative, Open Mathematics, Vol:13, No:1, 889–898, 2015.
3. Alfred Huber, A Novel Class of Solutions for a Non-linear Third Order Wave Equation Generated by the Weierstrass Transformation, Chaos, Solitons and Fractals, Vol:28, No:4, 972–978, 2006.
4. El-Said A. El-Wakil, Essam M. Abulwafa, Mohammed A. Abdou, Extended Weierstrass Transformation Method for Nonlinear Evolution Equations, Nonlinear Science Letters A: Mathematics, Physics and Mechanics, Vol:1, No:3, 253–262, 2010.

Dynamical behavior of analytical soliton solutions for nonlinear partial differential equation

Ajay Kumar¹ Haci Mehmet Baskonus²

¹ Department of Basic Engineering and Applied Science, College of Technology, Sardar Vallabhbhai Patel University of Agriculture & Technology, Meerut 250 110, Uttar Pradesh, India

²Department of Mathematics and Science Education, Faculty of Education, Harran University, Sanliurfa, Turkey

kajay9249@gmail.com, ajay.cot@svpuat.edu.in

Abstract

In this paper, we obtain some new analytical soliton solutions to the nonlinear partial differential equation (PDF) arising in nonlinear sciences ocean physics, fluid dynamics, plasma physics, scientific applications, and marine engineering. By employing powerful analytical methods such as the sine-Gordon Expansion method, we derive exact soliton solutions, including bright, dark, and singular solitons. There is no prior publication of these determined solutions. By giving appropriate values to free parameters, three-dimensional images are used to illustrate the dynamical wave structures of specific analytical solutions. The soliton solutions of other well-known equations in fluid dynamics, engineering physics, and other nonlinear science domains can also be obtained using this method.

Keywords: sine-Gordon Expansion method; nonlinear partial differential equation; analytical soliton solutions.

REFERENCES

- [1] Yamgoué, S.B., Deffo, G.R. & Pelap, F.B. A new rational sine-Gordon expansion method and its application to nonlinear wave equations arising in mathematical physics. *Eur. Phys. J. Plus* **134**, 380 (2019).
- [2] Ismael, H.F., Bulut, H. & Baskonus, H.M. Optical soliton solutions to the Fokas–Lenells equation via sine-Gordon expansion method and $(m+(G'/G))$ -expansion method. *Pramana - J Phys* **94**, 35 (2020).

LORENTZ MATRIX MULTIPLICATION: EXPLORING ITS RELATIONSHIP WITH FIBONACCI AND LUCAS NUMBERS

Ümmügülsün Çağlayan¹, İbrahim Gökcan², Ali Hikmet Değer³

¹Department of Mathematics, Karadeniz Technical University, Türkiye

²Faculty of Art and Science, Artvin Çoruh University, Türkiye

³Department of Mathematics, Karadeniz Technical University, Türkiye

ummugulsun.akbaba@gmail.com

gokcan@artvin.edu.tr

ahikmetd@ktu.edu.tr

Abstract

This paper introduces new matrix representations of Fibonacci and Lucas numbers by using Lorentz matrix multiplication. We present novel identities, characteristic equations, and quadratic properties derived from Lorentzian transformations applied to traditional matrix formulations of these sequences.

Keywords: Matrix multiplication, Fibonacci numbers, Lucas numbers.

REFERENCES

1. Ü. Akbaba and A.H.Değer, "On applications of Pell and Pell-Lucas numbers with matrix method," Journal of Intelligent Fuzzy Systems, vol.44, no.6, pp.10703-10707, 2023.
2. J.L. Brenner, "Lucas' matrix", The American Mathematical Monthly, vol.58, 220-221, 1951.
3. J. Ercolano, "Matrix generators of Pell sequences," The Fibonacci Quarterly, 17(1), 71-77, 1979.
4. S. Moore, "Fibonacci Matrices," Mathematical Gazette, 67, 56-57, 1983
5. H.W., Gould, "A history of the Fibonacci Q-matrix and a higher-dimensional problem," Fibonacci Quarterly, 9, 250-250, 1981.
6. H. Gündoğan, and O. Keçilioğlu, "Lorentzian matrix multiplication and the motions on Lorentzian plane," GlasnikMatematicki, 41, 329-334, 2006.
7. T. Koshy, Fibonacci and Lucas Numbers with Applications. New York: A Wiley-Interscience Publication (2001).

DECAY AND BLOW UP OF SOLUTIONS FOR A WEIGHTED BIHARMONIC EQUATION

Ayşe Fidan¹ and Erhan Pişkin²

¹Institute of Natural and Applied Sciences, Dicle University, Diyarbakır, Turkey

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

afidanmat@gmail.com, episkin@dicle.edu.tr

Abstract

In this presentation, we investigate the weighted biharmonic equation with nonlinear damping and source terms. The decay of the energy is established using Nakao's inequality. Furthermore, we prove that solutions blow up in finite time.

Keywords: Blowup, Decay, weighted biharmonic equation.

REFERENCES

1. M.R. Li, L.Y. Tsai, Existence and nonexistence of global solutions of same system of semilinear wave equations, *Nonlinear Anal.* 54(8), 1397--1415 (2003).
2. E. Pişkin, On the decay and blowup of solutions for a quasilinear hyperbolic equations with nonlinear damping and source terms, *Bound. Value Probl.*, 2015:127 (2015).
3. E. Pişkin, *Evolüsyon Denklemlerin Çözümlerinin Azalması (Decay of Solutions of Evolution Equations)*, Pegem Akademi Yayıncılık, 2025.
4. E. Pişkin, *Evolüsyon Denklemlerin Çözümlerinin Patlaması (Blowup of Solutions of Evolution Equations)*, Pegem Akademi Yayıncılık, 2022.
5. E. Pişkin, A. Fidan, Nonexistence of global solutions for the strongly damped wave equation with variable coefficients, *Universal Journal of Mathematics and Applications*, 5 (2) 51-56 (2022).

MODIFIED EXPONENTIAL FUNCTION METHOD FOR TRAVELING WAVE SOLUTIONS OF THE NONLINEAR MATHEMATICAL MODEL

Nursemin KAŞ¹, Tolga AKTÜRK²

¹Department of Mathematics, Institute of Science, Ordu University, Turkey

²Department of Mathematics and Science Education, Faculty of Education, Ordu University, Turkey

¹nurseminkas52@gmail.com; ²tolgaakturkk@gmail.com;

ABSTRACT

In this paper, the traveling wave solutions of the Oskolkov equation, a model describing the dynamics of incompressible viscoelastic Kelvin-Voigt fluid, are investigated using the modified exponential function method. This method is used to find analytical traveling wave solutions of the Oskolkov equation. Different traveling wave solutions are obtained by giving appropriate values to the parameters of the nonlinear mathematical model. In this way, two and three-dimensional graphs of the different wave solutions found are drawn according to the appropriate parameters..

Keywords: Modified Exponential Function method; Wave solutions: Oskolkov equation; Nonlinear partial differential equation;

REFERENCES

1. Durur , Hülya (2023). Exact Solutions of the Oskolkov Equation in Fluid Dynamics. Department of Computer Engineering, Faculty of Engineering.
2. Ghanbari , Behzad . New analytical solutions for the Oskolkov-type equations in fluid dynamics via a modified methodology. Department of Basic Science.2021;28 (104610).
3. Ak Turgut, Aydemir Tuğba , Saha Asit ,Kara Abdul Hamid,2018.Propagation of nonlinear shock waves for the generalised Oskolkov equation and its dynamic motions in the presence of an external periodic perturbation,78(2),
4. Aktürk Tolga , Kubal Çağlar. Analysis of wave solutions of (2+1)- dimensional Nizhnik – Novikov - Veselov equation . Department of Mathematics and Science Education, Faculty of Education. Ordu Univ. J. Sci. Tech., 2021, 11(1), 13-24.
5. Alkan Aslı,Aktürk Tolga,Bulut Hasan. The Traveling Wave Solutions Of The Conformable Time-Fractional Zoomeron Equation By Using The Modified Exponential Function Method. 2024, 25(1),108-114, 1370631.
6. Roshid-Or-Harun, Roshid Mamunur , Abdeljabbar Alrazi, Begum Momtaz,Basher Habibul. Abundant dynamical solitary waves through Kelvin-Voigt fluid via the truncated M-fractional Oskolkov model.Results in Physics 55 (2023),107128.

SOLUTIONS OF TWO EXPONENTIAL TYPE POTENTIALS FOR DIATOMIC MOLECULES

Haydar ALICI¹

¹Department of Mathematics, Harran University, Şanlıurfa, Türkiye

haydara@harran.edu.tr,

Abstract

In this study, we consider the Schrödinger equation with the Manning-Rosen and Deng-Fan molecular potentials over the positive half line. First, we obtain their closed form solutions when the angular quantum number $\ell = 0$. Then, we show that the two potentials are the same. Hence, by taking one of them, we obtain numerical solutions by using the Laguerre pseudospectral method, in the presence of the centrifugal term ($\ell \neq 0$) for which the solutions can not be expressed in closed form. Comparison with the exact solution for $\ell = 0$ and numerical literature results when $\ell \neq 0$ reveals that the Laguerre pseudospectral method produces highly accurate numerical results for the Schrödinger equation with the Manning-Rosen and Deng-Fan molecular potentials.

Keywords: Schrödinger equation, Manning-Rosen potential, Deng-Fan potential, Laguerre pseudospectral methods.

REFERENCES

1. Wang, P. Q., Zhang, L. H., Jia, C. S., & Liu, J. Y. (2012). Equivalence of the three empirical potential energy models for diatomic molecules. *Journal of Molecular Spectroscopy*, 274, 5-8.
2. Alıcı, H., & Ulusoy, S. (2025). On vibrational levels of the Deng-Fan molecular potential. *Physica Scripta*, 100(4), 045239.

EXPLICIT REPRESENTATIONS FOR A CLASS OF GAUSS HYPERGEOMETRIC FUNCTIONS

Haydar ALICI¹

¹Department of Mathematics, Harran University, Şanlıurfa, Türkiye

haydara@harran.edu.tr,

Abstract

In this study, we derive explicit elementary formulas for the Gauss hypergeometric functions which are the solutions of the Gauss hypergeometric equation with characteristic exponent differences $\{\frac{1}{2}, n + \frac{1}{2}, \alpha\}$ and $\{n + \frac{1}{2}, n + \frac{1}{2}, \alpha\}$ up to permutations and sign changes where n is a non negative integer. To this end, we obtain two different sets of linearly independent solutions of the specific second order ordinary differential equation where the first set contains the aforementioned Gauss hypergeometric functions and the second one consists of elementary functions. Then the identities will be obtained by expressing the hypergeometric solutions as the linear combinations of the elementary ones.

Keywords: Gauss hypergeometric equation, Gauss hypergeometric functions.

REFERENCES

1. Alıcı, H. (2023). Explicit general solution of the squared secant potential and some consequences. The Ramanujan Journal, 62(1), 111-140.

TOOL FRAME OF ROBOT END-EFFECTOR USING TWO RULED SURFACES CORRESPONDING TO THE HYPER-DUAL CURVE

Selahattin Aslan

Department of Mathematics, University of Ankara, Ankara, Turkey

selahattinnaslan@gmail.com

Abstract

Hyper-dual numbers are a new number systems developed at the beginning of the twenty-first century as extension of dual numbers. A hyper-dual number is constituted of two dual numbers, similarly, a hyper-dual curve constituted of two dual curves. If the dual curves of the hyper-dual curve are on the unit dual sphere, then this hyper-dual curve represents two ruled surfaces in Euclidean 3-space which intersect perpendicular along a common base curve. In this paper, we show that these ruled surfaces can be expressed by a same tool frame of the trajectory of the motion of a robot end-effector. Moreover, we give examples of tool frame of robot end-effector of two ruled surfaces corresponding to the hyper-dual curve.

Keywords: Hyper-dual numbers; Unit hyper-dual sphere; Hyper-dual curves; Ruled surfaces; Robot end-effector; Tool frame.

REFERENCES

1. Jeffrey Fike, Numerically exact derivative calculations using hyper-dual numbers}, 3rd Annual Student Joint Workshop in Simulation-Based Engineering and Design, 2009.
2. Jeffrey Fike, Juan Alonso, The development of hyper-dual numbers for exact second-derivative calculations, 49th AIAA Aerospace Sciences Meeting including the New Horizons Forum and Aerospace Exposition, 4-7, 2011.
3. Avraham Cohen, Moshe Shoham, Application of hyper-dual numbers to multi-body kinematics, J. Mech. Rob., 8, 2015.
4. Aleksandr Petrovich Kotelnikov, Screw calculus and some applications to geometry and mechanics, Annal Imp. Univ., Kazan, Russia, 1895.
5. Selahattin Aslan, Murat Bekar, Yusuf Yaylı, Ruled surfaces corresponding to hyper-dual curves, Hacettepe Journal of Mathematics and Statistics, 51(1), 187-198, 2022.
6. Beom-Sahng Ryuh, Gordon R. Pennock, Accurate motion of a robot end-effector using the curvature theory of ruled surfaces, Journal of Mechanisms, Transmissions, and Automation in Design, 110 (4) 383 – 388, 1988.

Generalized Lacunary Statistical Convergence of Modulus Function Sequences

Muhammet Emin Aydın¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

Aydinemin607@gmail.com

Abstract

In this paper, we propose a generalized structure for statistical convergence by adapting the concept of density functions using sequences of modulus functions $(f_{\{t\}})$ and an order parameter $a \in (0,1]$. This method introduces a new density framework designed specifically for lacunary sequences, facilitating the definitions of strong $(f_{\{t\}})$ -lacunary summability of order a and $(f_{\{t\}})$ -lacunary statistical convergence of order a . These newly defined concepts serve as a bridge between traditional convergence and statistical convergence in the context of lacunary sequences, providing a middle ground that enhances analytical versatility. Furthermore, we establish inclusion results and investigate the interconnections between these notions, offering a thorough exploration of their implications. This study extends the reach of statistical convergence theory and advances the field by presenting a more generalized analytical approach.

Keywords: Lacunary statistical convergence, Lacunary summability, Modulus function, Difference sequence space.

REFERENCES

- [1] Zygmund, A.: Trigonometric Series, Cambridge University Press, Cambridge, (1979).
- [2] Steinhaus, H.: Sur la convergence ordinaire et la convergence asymptotique. Colloquium Mathematicum. 2, 73-74 (1951).
- [3] Fast, H.: Sur la convergence statistique. Colloquium Mathematicum. 2, 241-244 (1951).
- [4] Schoenberg, I.J.: The integrability of certain functions and related summability methods. The American Mathematical Monthly. 66, 361-375 (1959).
- [5] Salat, T.: On statistically convergent sequences of real numbers. Mathematica Slovaca. 30, 139-150 (1980).

MULTIPLICATIVE EQUIAFFINE CURVES AND THEIR PROPERTIES

Meltem Ogrenmis¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

mogrenmis@firat.edu.tr

Abstract

This study introduces a multiplicative calculus framework for analyzing curves in equiaffine space. It defines multiplicative arc length, curvature, and torsion, and derives the corresponding Frenet equations. Curves with constant multiplicative curvature and torsion are characterized, accompanied by illustrative examples.

Keywords: Equiaffine space; Multiplicative calculus; Curves; Curvatures; Frenet Equations

REFERENCES

1. H.W. Guggenheimer, Differential Geometry, McGraw-Hill, New York, 1963.
2. Georgiev, S.G.; Zennir, K. Multiplicative Differential Calculus, 1st ed.; Chapman and Hall/CRC: New York, NY, USA, 2022.
3. 10. Georgiev, S.G. Multiplicative Differential Geometry, 1st ed.; Chapman and Hall/CRC: New York, NY, USA, 2022.
4. 11. Georgiev, S.G.; Zennir, K.; Boukarou, A. Multiplicative Analytic Geometry, 1st ed.; Chapman and Hall/CRC: New York, NY, 2022.
5. Has, A.; Yilmaz, B. On non-Newtonian helices in multiplicative Euclidean space E^3 , arXiv 2024, arXiv.2403.11282.
6. Aydin, M.E.; Has, A.; Yilmaz, B. A non-Newtonian approach in differential geometry of curves: Multiplicative rectifying curves. Bull. Korean Math. Soc. 61, 849–866, 2024.

CHARACTERIZATION OF PLANE CURVES WITH CONSTANT MULTIPLICATIVE EQUIAFFINE CURVATURE

Meltem Ogrenmis¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

mogrenmis@firat.edu.tr

Abstract

This study presents a multiplicative calculus approach to the differential geometry of plane curves within equiaffine geometry. By introducing multiplicative arc length and curvature, the research derives the corresponding Frenet formulas and establishes an analog of the fundamental theorem. The classification of plane curves with constant multiplicative curvature is also provided, supported by illustrative examples.

Keywords: Equiaffine plane; Plane curves; Multiplicative Equiaffine curves

REFERENCES

1. D. Davis, Generic affine differential geometry of curves in R^n , Proceedings of the Royal Society of Edinburgh A, 136, 1195–1205, 2006.
2. Aydin, M.E.; Has, A.; Yilmaz, B. A non-Newtonian approach in differential geometry of curves: Multiplicative rectifying curves. Bull. Korean Math. Soc. 61, 849–866, 2024.
3. H.W. Guggenheimer, Differential Geometry, McGraw-Hill, New York, 1963.
4. Georgiev, S.G.; Zennir, K. Multiplicative Differential Calculus, 1st ed.; Chapman and Hall/CRC: New York, NY, USA, 2022.
5. 10. Georgiev, S.G. Multiplicative Differential Geometry, 1st ed.; Chapman and Hall/CRC: New York, NY, USA, 2022.
6. 11. Georgiev, S.G.; Zennir, K.; Boukarou, A. Multiplicative Analytic Geometry, 1st ed.; Chapman and Hall/CRC: New York, NY, 2022.
7. M. E. Aydin, A. Has, B. Yilmaz, Multiplicative rectifying submanifolds of multiplicative Euclidean space, Mathematical Methods in the Applied Sciences, 2024, <https://doi.org/10.1002/mma.10329>, 1–11.
8. A. Has, B. Yilmaz, A non-Newtonian conics in multiplicative analytic geometry, Turkish Journal of Mathematics, 48(5), 976-994, 2024.

FRACTIONAL DERIVATIVES: DEFINITIONS AND PROPERTIES

Çiğdem Süsoy Atalar¹, Tefik Şahin¹

¹ Department of Mathematics, Institute of Science, University of Amasya, Amasya, Turkey

cigdemsusoyatalar@gmail.com,

Abstract

Fractional derivatives are the extension of the concept of derivative to fractional (rational or irrational) degrees and have an important place in mathematical analysis. It extends the process of differentiation to rational or irrational degrees. In this presentation, different fractional derivative definitions and properties will be expressed.

Keywords: Fractional derivatives, Fractional degrees, Mathematical analysis

REFERENCES

1. I.Podlubny (1999). Fractional Differential Equations. Academic Press, USA. Konstantinos A. Lazopoulos and Anastasios
2. K. Lazopoulos (2021). On Fractional Geometry of Curves fractal and fractional , 5, 161. <https://doi.org/10.3390/fractalfract5040161>

ON BRIEF NOTE REGARDING F-CONTRACTIONS IN COMPLETE S-METRIC SPACES

Süheyla Elmas¹

Ferdi Çınar²

¹ Faculty of Education, Ataturk Universiy, Erzurum, Turkey

² Graduate School of Fine Arts, Ataturk Universiy, Erzurum, Turkey

suheylaelmas@atauni.edu.tr,

ferdi.cinar18@ogr.atauni.edu.tr,

Abstract

In this writing, we investigate fixed points of F-contraction in an abstract space. Additionally, the effectiveness of our work is confirmed through appropriate examples

Keywords: Fixed point, S-metric, F-contraction

REFERENCES

- [1] Banach, B: Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. Fundam. Math. 3, 133-181 (1922)
- [2] Suzuki, T: A new type of fixed point theorem in metric spaces. Nonlinear Anal. 71, 5313-5317 (2009)
- [3] Suzuki, T: Generalized distance and existence theorems in complete metric spaces. J. Math. Anal. Appl. 253, 440-458 (2001)
- [4] Suzuki, T: Several fixed point theorems concerning τ -distance. Fixed Point Theory Appl. 2004, 195-209 (2004)
- [5] Tataru, D: Viscosity solutions of Hamilton-Jacobi equations with unbounded nonlinear terms. J. Math. Anal. Appl. 163, 345-392 (1992)
- [6] Vályi, I: A general maximality principle and a fixed point theorem in uniform space. Period. Math. Hung. 16, 127-134 (1985)
- [7] Włodarczyk, K, Plebaniak, R: Quasigauge spaces with generalized quasipseudodistances and periodic points of dissipative set-valued dynamic systems. Fixed Point Theory Appl. 2011, Article ID 712706 (2011)
- [8] Włodarczyk, K, Plebaniak, P: Kannan-type contractions and fixed points in uniform spaces. Fixed Point Theory Appl. 2011, Article ID 90 (2011)

A STUDY ON WARDOWSKI CONTRACTION IN A-METRIC SPACES

Süheyla Elmas¹

Aleyna Özel²

^{1 1} Faculty of Education, Ataturk Universiy, Erzurum, Turkey

² Graduate School of Fine Arts, Ataturk Universiy, Erzurum, Turkey

suheylaelmas@atauni.edu.tr,

aleyna.kamber16@ogr.atauni.edu.tr,

Abstract

In this writing, we explore Wardowski's contraction principle for F-contraction mappings and demonstrate the existence and uniqueness of fixed points in A-Metric Spaces.

Keywords: Fixed point, A-metric, F-contraction

REFERENCES

- [1] Banach, B: Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. Fundam. Math. 3, 133-181 (1922)
- [2] D. Wardowski, Fixed Points of a New Type of Contractive Mappings in Complete Metric Spaces, Fixed Point Theory Appl. 2012 (2012), 94
- [3] E. Karapınar, A. Fulga, R.P. Agarwal, A Survey: F-Contractions With Related Fixed Point Results, J. Fixed Point Theory Appl. 22 (2020), 69. <https://doi.org/10.1007/s11784-020-00803-7>.
- [4] Fabiano, Nicola, et al. "On F-contractions: A survey." Contemporary Mathematics (2022): 327-342.
- [5] Acar, Özlem, Aybala Sevd e Özkapu, and Mahpeyker Öztürk. "Some fixed point results on ultrametric spaces endowed with a graph." Demonstratio Mathematica 57.1 (2024): 20230132.
- [6] Bajović, Dusan, et al. "Remarks on Some Results of Extended Interpolative Hardy-Rogers-Geraghty-Wardowski Contractions and Ćirić-Reich-Rus Type F-Contractions." Sahand Communications in Mathematical Analysis 22.1 (2025): 25-38.
- [7] Abbas, Mujahid, Bashir Ali, and Yusuf I. Suleiman. "Generalized coupled common fixed point results in partially ordered A-metric spaces." Fixed point theory and applications 2015 (2015): 1-24.
- [8] Gähler, S: 2-metrische raume und ihre topologische strukture. Math. Nachr. 26, 115-148 (1963)

INVESTIGATION OF APPLICATIONS OF FIXED POINT ITERATION METHODS WITH THE AID OF GREEN'S FUNCTION

Sezgin Akbulut, İsa Yıldırım, Büşra Melike Kör

Department of Mathematics, University of Atatürk, Erzurum, Turkey

sezginakbulut@atauni.edu.tr, isayildirim@atauni.edu.tr, busramelikeakman@gmail.com

Abstract

In this paper, a new fixed point iteration scheme is introduced. The strong convergence of the scheme to the fixed point of a contractive mapping is demonstrated. It is also proved that the defined scheme converges faster to the fixed point compared to the Modified-JK iteration scheme existing in the literature. Moreover, the iteration scheme is shown to be faster than some existing methods by means of examples and illustrated with tables and graphs. Then data dependency is proved for the scheme in question. Then the defined iteration scheme is extended by embedding it in the Green's function. It is shown that the extended iteration scheme converges strongly to the fixed point of the transformation and is weakly- ω^2 stable. Finally, it is shown on an example that the extended iteration scheme converges to the fixed point faster than the Modified-JK-Green iteration scheme.

Keywords: Fixed Point Theory; Iteration Scheme; Convergence Analysis, Strong Convergence; Speed Comparison; Efficiency; Green's Function; Weak- ω^2 Stability; Boundary Value Problem.

REFERENCES

1. Abushammala, Mariam and Khuri, Suheil A and Sayfy, Ali. (2015). A novel fixed point iteration method for the solution of third order boundary value problems. *Applied Mathematics and Computation*, 131-141. Elsevier.
2. Ahmad, Junaid and Arshad, Muhammad and George, Reny. (2023). A fixed point iterative scheme based on Green's function for numerical solutions of singular BVPs. *AIMS Mathematics*, 29517-29534.
3. Ahmad, Junaid and Arshad, Muhammad and Hammad, Hasanen A and Kattan, Doha A. (2023). A Three-Step Iterative Scheme Based on Green's Function for the Solution of Boundary Value Problems. *International Journal of Analysis and Applications*, 116.

EXISTENCE RESULT OF WEAK SOLUTION FOR A SINGULAR KIRCHHOFF TYPE PROBLEM WITH STEKLOV BOUNDARY CONDITION

Berat Süer¹ and Zehra Yücedag²

¹Dicle University, Institute of Natural and Applied Sciences, Diyarbakir, Turkey

²Dicle University, Vocational School of Social Sciences, 21280-Diyarbakir, Turkey

E-mail: beratsuer72@gmail.com and zyucedag@dicle.edu.tr

Abstract.

In this paper, we study the existence of solutions for a singular $p(x)$ -Kirchhoff type problem with Steklov boundary value in variable exponent Sobolev spaces. Using the variational approach, we prove the existence a nontrivial weak solution on variable exponent Sobolev spaces under appropriate conditions.

We study the following problem involving singular $p(x)$ -Kirchhoff type equation with Hardy term

$$\begin{cases} -M \left(\int_{\Omega} \frac{|\nabla u|^{p(x)}}{p(x)} dx \right) \Delta_{p(x)} u = \frac{|u|^{q(x)-2}}{|x|^{q(x)}}, & x \in \Omega \\ |\nabla u|^{p(x)} \frac{\partial u}{\partial \nu} = \lambda f(x, u), & x \in \partial\Omega \end{cases} \quad (E)$$

where $\Omega \subset \mathbb{R}^N$ ($N \geq 2$) is a bounded with smooth boundary, $p(x)$ is continuous functions on $\bar{\Omega}$ such that $p^- := \inf p(x)$, $f: \partial\Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is a Carathéodory condition, λ is a positive parameter,

$\frac{\partial u}{\partial \nu}$ is the outer unit normal derivative on $\partial\Omega$, $M: (0, \infty) \rightarrow (0, \infty)$ is a continuous and

$\Delta_{p(x)} u = \operatorname{div} (|\nabla u|^{p(x)-2} \nabla u)$ is $p(x)$ -Laplacian type operator.

Keywords: $p(x)$ -Laplacian operator, Variational methods, Weak solutions. Singular problem.

Theorem 1.1. The problem (E) has a nontrivial weak solution for all $\lambda > 0$.

References.

- [1] X. L. Fan, D. Zhao D, On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$, J. Math. Anal. Appl. 263 (2001) 424-446
- [2] O. Kovacik, J. Rakosnik, On spaces $L^{p(x)}(\Omega)$ and $W^{k,p(x)}(\Omega)$, Czechoslovak Mathematical Journal. 41 (1991) 592-618.
- [3] M. Laghzal, A. Touzani, On a singular Kirchhoff type problems driven by $p(x)$ -Laplacian operator, Appl. Anal. 101 (2022), 5932-5947.

R. Mahdavi Khanghahi, A. Razani, Solutions for a Singular Elliptic Problem Involving the $p(x)$ -Laplacian, Filomat. 32 (2018) 4841-4850.

USE OF ROUGH SET THEORY IN DATA MINING

M. Mustafa Beydağı¹, İlhan İcen¹, A. Fatih Özcan¹

¹Department of Mathematics, Inonu University, Malatya, Türkiye

mbeydagi23@gmail.com , ilhan.icen@inonu.edu.tr , abdullah.ozcan@inonu.edu.tr

Abstract

Rough set theory, presented by Pawlak, has been studied in many disciplines along with mathematics. Especially in data mining, rough set theory makes a great contribution as it enables the storage of large data and the reduction of unimportant data.

In this study, general information about rough set theory, its usage areas and its importance in data mining are emphasized.

Keywords: Rough set theory; Data mining; Data reduction.

REFERENCES

1. Pawlak, Z., (1982). Rough sets, Int. J. Comput. Inform. Sci. 11 341-356.
2. J. Komorowski, L. Polkowski ve A. Skowron, Rough Sets: A Tutorial, 9 dü.,Singapur: Springer-Verlag, 1998.
3. J. W. Grzymala-Busse, “Managing Uncertainty in Expert Systems”, Boston, London, 1991.
4. B. Walczak ve D. Massart, “Rough sets theory”, Chemometrics and Intelligent Laboratory Systems, cilt 47, pp. 1-16, 1999.
5. P. Jacobs, “Data Mining: What General Managers Need to Know”, Harvard Management Update, cilt 4, no. 10, p. 8, 1999.
6. Ş. Doğan ve İ. Türkoğlu, “Iron-Deficiency Anemia Detection From Hematology Parameters By Using Decision Trees”, International Journal of Science & Technology, cilt 3, no. 1, pp. 85-92, 2008.
7. S. Albayrak ve Ş. K. Yılmaz, “Veri Madenciliği: Karar Ağacı Algoritmaları ve IMKB Verileri Üzerine Bir Uygulama”, Süleyman Demirel University The Journal of Faculty of Economics and Administrative Sciences, pp. 1-52, 2009.
8. C. Erden, “Kaba Kümeler Teorisi ve Trafik Kazaları Üzerine Uygulaması”, İstanbul Üniversitesi Endüstri Mühendisliği Anabilim Dalı Yüksek Lisans Tezi, Haziran 2014.

SOLUTION OF AN OPTIMIZATION PROBLEM USING GROVER'S ALGORITHM FOR QUTRIT SYSTEMS

Orhan Amirov¹, Mesut Karabacak¹

¹Department of Mathematics, Atatürk University, Erzurum, Türkiye

amirov.orhan@atauni.edu.tr, mkarabacak@atauni.edu.tr,

Abstract

In this study, we investigate the acceleration of solving an optimization problem using Grover's algorithm implemented on qutrit systems. Unlike conventional qubit-based approaches, qutrit-three level quantum systems offer a richer Hilbert space, potentially enabling more efficient quantum search processes. The proposed approach aims to lay the groundwork for addressing more complex and larger-scale problems by leveraging the advantages of higher-dimensional quantum systems. This work serves a preliminary study toward extending quantum algorithm applications beyond binary quantum architectures.

Keywords: Quantum computing , Qubit systems, Qutrit systems, Grover's Algorithm , Superposition, Quantum Fourier Transform(QFT).

REFERENCES

1. Khan, F. S., & Pervez, K. M. (2023). Realisation of Grover's search algorithm with photonic qutrits. *Quantum Information Processing*, 22(10), 353.
2. Pawela, Ł., Gawron, P., & Miszczak, J. A. (2021). Grover search on qudit-based quantum architectures. *Physical Review A*, 104(4), 042611.
3. Li, T., Li, L., & Wu, Z. (2022). High-fidelity implementation of Grover's search on a superconducting qutrit processor. *Physical Review Applied*, 18(6), 064001.
4. Long, G.-L., & Liu, Y. (2020). Generalised Grover algorithm with multi-valued logic for optimisation problems. *New Journal of Physics*, 22(11), 113015.
5. Gokhale, P., Angiuli, O., Martonosi, M., & Chong, F. T. (2022). Hybrid quantum-classical Grover search for combinatorial optimisation. *IEEE Transactions on Quantum Engineering*, 3, 3101611.
6. Shukla, V., Misra, A., & Verma, H. (2021). Efficient amplitude amplification on qutrit systems. *Quantum Information Processing*, 20(9), 275.
7. Li, Y., Ding, Y., & You, J. Q. (2013). Realisation of quantum algorithms on superconducting qutrits. *Scientific Reports*, 3, 1624.
8. Yan, F., Sung, Y., Place, A. P., et al. (2019). Tunable coupling and high-fidelity entanglement of superconducting qutrits for quantum search. *Nature Communications*, 10, 5618.

Oscillation results of second-order noncanonical neutral differential equations with distributed deviating arguments

Ercan Tunç¹

¹Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Türkiye

ercantunc72@yahoo.com

Abstract

Sufficient conditions for the oscillation of solutions to a class of second-order noncanonical differential equations with bounded and unbounded neutral coefficients are established. The results obtained improve and simplify existing ones in the literature.

Keywords: Oscillation; neutral; second-order; noncanonical.

REFERENCES

1. S. R. Grace, J. R. Graef, T. Li, E. Tunç, Oscillatory behavior of second-order nonlinear noncanonical neutral differential equations, *Acta Univ. Sapientiae-Mathematica*, Vol: 15, 259–271, 2023.
2. R. Koplatadze, G. Kvinikadze, I. P. Stavroulakis, Properties A and B of n th order linear differential equations with deviating argument, *Georgian Mathematical Journal*, Vol: 6, 553–566, 1999.
3. E. Tunç, J. R. Graef, Oscillation results for second order neutral dynamic equations with distributed deviating arguments, *Dynamic Systems and Applications*, Vol: 23, 289-303, 2014.

CURVES AND THEIR PROPERTIES IN DYNAMIC GEOMETRY

Kübra Tunc¹ and Alper Osman Ogrenmis¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

tunckubra600@gmail.com , aogrenmis@firat.edu.tr

Abstract

In this study, Frenet curves on time scales are considered within a unified framework that connects discrete and continuous analysis. Fundamental geometric concepts such as tangent lines, normal planes, osculating planes, and curvature are presented for plane and space curves in $\mathbb{R}^n$. The Frenet frame and its corresponding equations are given in the context of time scale calculus, offering a generalized geometric approach over varying time domains.

Keywords: Time Scale; Curves; Curvatures; Frenet Equations

REFERENCES

1. H.W. Guggenheimer, Differential Geometry, McGraw-Hill, New York, 1963.
2. D. Davis, Generic affine differential geometry of curves in $\mathbb{R}^n$, Proceedings of the Royal Society of Edinburgh A, 136, 1195–1205, 2006.
3. Georgiev, Svetlin G. Dynamic Geometry on Time Scales (1st ed.). Chapman and Hall/CRC: New York, NY, USA, 2021.
4. Bohner, M., & Georgiev, S. G. Multivariable dynamic calculus on time scales, 449-515. Cham: Springer, 2016.
5. Georgiev, Svetlin G. Functional dynamic equations on time scales. Springer International Publishing, 2019.
6. M. Bohner and G. S. Guseinov, Multiple integration on time scales, Dynam. Systems Appl. 14 , no. 3–4, 579–606, 2005.

ON THE MANGENETIC CURVES ACCORDING TO THE RIBBON FRAME IN MINKOWSKI SPACE

Yusuf ÖZDEMİR¹, Mustafa YENEROĞLU¹

¹ Department of Mathematics, University of Firat, Elazig, Turkey

yusuf-islam72@gmail.com, mustafayeneroglu@gmail.com

Abstract

Curves in Minkowski space have been examined in many studies according to different frame works. While examining curves, researchers studied frameworks such as the Frenet frame, Bishop frame and Ribbon frame. These frame help us with characterizations of curves.

The aim of this paper is to investigate magnetic curves according to ribbon frame in 3-dimensional Minkowski space. For this purpose, an ribbon frame was first defined. Then, magnetic curves were given depending on this frame and various characterizations were obtained.

Keywords: Frenet frame, magnetic curves, ribbon frame, Minkowski space

REFERENCES

- Körpınar, T. and Demirkol, R. Gravitational magnetic curves on 3D Riemannian manifolds. [International Journal of Geometric Methods in Modern Physics](#), 15(11), 2018.
- Kazan, A., Karadağ, H.B. Magnetic Curves According to Bishop Frame and Type-2 Bishop Frame in Euclidean 3-Space. *British J. Math.*, 22(4), 1-18, 2017.
- Barros, M., Cabrerizo, L., Fernandez, M., Romeo, A. Magnetic Vortex Filament Flows. *J. Math Phys*, 48, 1-27, 2007.
- Barros, M., Romero, A. Magnetic vortices. *EPL*, 77, 1-5, 2007.
- Bohr, J. and Markvorsen, S. Ribbon Crystals, *Plos one*, Vol:8, No:10, 2013.
- Ayyıldız, N., Coken, A.C., Yucesan, A. Differential-geometrical conditions between geodesic curves and ruled surfaces in the Lorentz space, *Balkan Journal of Geometry and Its Applications*, 7(1), 1-6, 2002.

FIXED POINT THEOREMS FOR FOUR-POINT CONTRACTIONS

Merve Özkan¹

¹ Department of Mathematics, University of Atatürk, Erzurum, Turkey

merve.ozkan@atauni.edu.tr,

Abstract

In this paper, we give a novel type of mappings that are four-point instead of two or three-point. Some fixed point theorems in metric spaces are proved using four-point contractions, and an example is also given related to four-point contractions.

Keywords: metric space; Fixed point theorems; four-point contractions; mappings contracting perimeters of triangles.

REFERENCES

1. Banach S., Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, Fundam. Math., Vol:3, 133-181, 1922.
2. Ćirić Lj B., Generalized contractions and fixed-point theorems, Publ. Inst. Math., Vol:12, 19-26, 1971.
3. Petrov E., Fixed point theorem for mappings contracting perimeters of triangles, J. Fixed Point Theory Appl., Vol:25, 74-85, 2023.
4. Petrov E., Bisht R.K., Fixed point theorem for generalized Kannan type mappings, arXiv:2308.05419 2023. (Submitted)
5. Bey C., Petrov E. Salimov, R. On three-point generalizations of Banach and Edelstein fixed point theorems, Filomat, Vol:39, No:1, 185-195, 2025.
6. Pacurar C. M., Popescu O., Fixed point theorem for generalized Chatterjea type mappings, Acta Math. Hungar., Vol:173173, No:2, 500-509, 2024.
7. Popescu O., Stan G., Some remarks on Reich and Chatterjea type nonexpansive mappings, Mathematics, Vol:8, 1270, 2020.
8. Bisht R., Petrov E., A three point extension of Chatterjea's fixed point theorem with at most two fixed points, arXiv:2403.07906 2024.
9. Chatterjea S. K., Fixed-point theorems, C. R. Acad. Bulgare Sci., Vol:25, 727-730, 1972.1010.
10. Dung N. Van, Petrusel A., On iterated function systems consisting of Kannan maps, Reich maps, Chatterjea type maps, and related results, J. Fixed Point Theory Appl., Vol:19, 2271-2285, 2017.
11. Faraji H., Nourouzi K., A generalization of Kannan and Chatterjea fixed point theorems on complete-metric spaces, Sahand Commun. Math. Anal., Vol:6, 77-86, 2017.
12. Kannan R., Some results on fixed point-II, Amer. Math. Monthly, Vol:76, 405-408, 1969.
13. Subrahmanyam P. V. Completeness and fixed points, Monatsh. Math., Vol:80, 325-330, 1975.

SOFT SET THEORY

AND VARIOUS APPLICATION AREAS

Semih GEÇEN¹, A. Fatih ÖZCAN², İlhan İÇEN³

¹Department of Mathematics, University of Inonu, Malatya, Türkiye, semihgecen44@gmail.com

²Department of Mathematics, University of Inonu, Malatya,Türkiye, abdullah.ozcan@inonu.edu.tr

³Department of Mathematics, University of Inonu, Malatya, Türkiye, ilhan.icen@inonu.edu.tr

Abstract

Soft set theory, defined by Molodtsov, has found a wide place in contemporary mathematics [1]. After the soft set theory, which gave a new perspective to the concepts of completeness and precision in mathematics, was introduced, it was studied by many mathematicians from topological, categorical and algebraic perspectives [2-22]. In this study, the development process and various application areas of soft set theory, which has been studied by many mathematicians from topological, categorical and algebraic perspectives, are discussed.

Keywords: Soft set, Development process, Application areas.

REFERENCES

1. D.A. Molodtsov, *Soft set theory-First results*, Comput. Math. Appl., 37 (4-5), 19-31, 1999.
2. P. K. Maji, R. Biswas and R. Roy, *Soft set theory*, Comput. Math. Appl., 45(4-5), 555-562, 2003.
3. M. I. Ali, *A note on soft sets, rough soft sets and fuzzy soft sets*, Appl.Soft Comput., 11, 3329-3332, 2011.
4. H. Aktas and N. Cagman, *Soft sets and soft groups*, Inform.Sci., 77(13), 2726-2735, 2007.
5. A. O. Atagun, A. Sezgin, *Soft substructures of rings, fields and modules*, Comput. Math. Apply., 61, 592-601, 2011.
6. M. Shabir, M. Naz, *On Soft topological spaces*, Comput. Math. Apply., 61(7), 1786-1799, 2011.

ELASTIC ANALYSIS OF ROTATING FG PRESSURE VESSELS USING THE POWER SERIES METHOD

Özlem Öcal¹, Durmuş Yarımabaç¹

¹ Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

ozlemocal880@gmail.com, durmusyayarimpabuc@osmaniye.edu.tr

Abstract

In this paper, stresses in rotating thick walled functionally graded cylindrical and spherical pressure vessels are investigated using elasticity theory. It is assumed that the material properties except Poisson's ratio vary in the radial direction with the Voigt model. Under these conditions, the analytical solution of the Navier equation is obtained by the power series solution method. The solutions available in the literature are used to verify the results obtained in this study. The effect of the rotation and pressure on the radial displacement, radial and circumferential stress are discussed for a randomly selected material pair consisting of metal and light metal.

Keywords: Elastic analysis, Cylindrical and spherical pressure vessels, Rotation, Functionally graded materials.

REFERENCES

1. Jabbari, M., Nejad, M.Z., Ghannad, M., Thermo-elastic analysis of axially functionally graded rotating thick cylindrical pressure vessels with variable thickness under mechanical loading, International Journal of Engineering Science, Vol. 96, 1–18, 2015.
2. Mazarei, Z., Nejad, M.Z., Hadi, A., Thermo-Elasto-Plastic Analysis of Thick-Walled Spherical Pressure Vessels Made of Functionally Graded Materials, International Journal of Applied Mechanics, Vol:8, No:4, 1650054, 2016.
3. Kacar, I., Exact elasticity solutions to rotating pressurized axisymmetric vessels made of functionally graded materials, Materialwiss und Werkstofftech, Vol:51, 1481-1492, 2020.
4. Yarımabaç, D., Çalışkan, A., Elastic Analysis Of Functionally Graded Rotating Pressurized Cylinders, Composites: Mechanics, Computations, Applications: An International Journal, Vol:15, No:1, 53-64, 2024.

OSTROWSKI-GRÜSS TYPE INEQUALITIES VIA KATUGAMPULA FRACTIONAL INTEGRALS

CANAN ÖDEMiŞ*¹ and ABDULLAH AKKURT¹

¹Department of Mathematics, Faculty of Science, Kahramanmaraş Sütçü İmam University, Turkey

cananodemis200683@gmail.com, abdullahmat@gmail.com

Abstract

In this study, the authors obtained new integral inequalities of Ostrowski and Ostrowski-Grüss type by using Katugampola fractional integral definition. It was shown that the obtained results support the results in the literature with special selection of parameters.

Let the space $X_c^p(a, b)$, $(c \in \mathbb{R}, 1 \leq p \leq \infty)$ of those complex-valued Lebesgue measurable functions on for which $\|f\|_{X_c^p} < \infty$, where the norm is defined by (Kilbas, 2006):

$$\|f\|_{X_c^p} = \left(\int_a^b |t^c f(t)|^p \frac{dt}{t} \right)^{\frac{1}{p}} < \infty,$$

for $1 \leq p \leq \infty$, $c \in \mathbb{R}$ and for the case ,

$$\|f\|_{X_c^p} = \text{ess sup}_{a \leq t \leq b} [t^c |f(t)|].$$

Let $[a, b] \subset \mathbb{R}$ be a bounded interval. Then, for any function and any order , the Katugampola fractional integrals defined on the left and right are given by (Katugampola, 2011):

$${}^\rho I_{a+}^\alpha f(x) = \frac{\rho^{1-\alpha}}{\Gamma(\alpha)} \int_a^x \frac{t^{\rho-1}}{(x^\rho - t^\rho)^{1-\alpha}} f(t) dt,$$

and,

$${}^\rho I_{b-}^\alpha f(x) = \frac{\rho^{1-\alpha}}{\Gamma(\alpha)} \int_x^b \frac{t^{\rho-1}}{(t^\rho - x^\rho)^{1-\alpha}} f(t) dt.$$

with $a < t < b$ and if the integral exists.

Keywords: Katugampola Fractional Integral, Integral Inequalities, Ostrowski inequality.

REFERENCES

1. Farid, G., Mehmood, S., Younis, B. A., Ahamd, H. U. M., Egami, R. H., and Adam, A. M. I. (2024). Generalized Ostrowski and Ostrowski-Grüss type inequalities. Rendiconti del Circolo Matematico di Palermo Series 2, 1-11.

A STUDY ON OSCILLATION OF NEUTRAL DIFFERENTIAL EQUATIONS WITH DAMPING TERM

Ercan Tunç¹

Adil Kaymaz¹

¹Department of Mathematics, Gaziosmanpaşa University, Tokat, Turkey

ercantunc72@yahoo.com

adilkaymaz@gmail.com,

Abstract

In this work, it has been concerned that the oscillatory behavior of solutions to damped second-order linear functional differential equations with a mixed neutral term. It is presented a new oscillation criteria that improve and extend some existing ones in the literature. Example to illustrate the result is included.

Keywords: Oscillation, Neutral Differential Equations of Mixed Type, Second Order, Damping Term.

REFERENCES

1. Ercan Tunç, Adil Kaymaz. Oscillatory behavior of second-order half-linear neutral differential equations with damping. *Advances in Dynamical Systems and Applications*, Vol: 14, No: 2, 213-227, 2019.
2. Ercan Tunç, Adil Kaymaz. On oscillation of second-order linear neutral differential equations with damping term. *Dynamic Systems and Applications*, 28, No: 2, 289-301, 2019.
3. Ercan Tunç, John R. Graef. Oscillation results for second order neutral dynamic equations with distributed deviating arguments. *Dynamic Systems and Applications*, 23, 289-303, 2014.

Unital Designs Admitting Blocking Sets

Kevser Soytürk¹, Mustafa Gezek¹, Mahmut Ergüt¹

¹Department of Mathematics, University of Tekirdag Namık Kemal, Tekirdag, Turkey

ksoyturk@nku.edu.tr, mgezek@nku.edu.tr, mergut@nku.edu.tr

Abstract

A unital design of order q is a combinatorial structure consisting of q^3+1 points, in which each block contains exactly $q+1$ points, and every distinct pair of points is contained in precisely one block. A blocking set in such a design is defined as a subset of the point set that intersects every block in at least one point, yet does not fully contain any block. In this study, we address the classification of unital designs that contain a blocking set for small values of q , presenting both our methodology and computational results.

Keywords: Blocking Sets; Discrepancy; Unital design.

REFERENCES

1. Al-Azemi, Abdullah, Anton Betten, and Dieter Betten, Unital designs with blocking sets, Discrete Applied Mathematics, 163 (2014): 102-112.
2. A. E. Brouwer, Some unitals on 28 points and their embeddings in projective planes of order 9, in: Geometries and Groups (Berlin, 1981), in: Lecture Notes in Math., vol. 893, Springer, Berlin, 1981, pp. 183–188.
3. Vedran Krčadinac, Steiner 2-designs $S(2,4,28)$ with nontrivial automorphisms, Glas. Mat. Ser. III, 37(57)(2), 2002, pp. 259–268.
4. R. Mathon, A. Rosa, $2-(v,k,\lambda)$ designs of small order, in C. J. Colbourn, J. H. Dinitz (Eds.), The CRC Handbook of Combinatorial Designs, CRC Press, New York, 1996, pp. 3–41.
5. Anton Betten, Dieter Betten, Vladimir D. Tonchev, Unitals and codes, Discrete Math. 267 (1-3) (2003)23–33. Combinatorics 2000 (Gaeta).

ON THE OSCILLATION OF FOURTH-ORDER NEUTRAL DIFFERENTIAL EQUATIONS

Orhan Özdemir¹ and Ercan Tunç¹

¹Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Türkiye

orhanozdemir37@yahoo.com, ercantunc72@yahoo.com

Abstract

In this talk, we consider a class of fourth--order functional differential equations with a mixed neutral term and establish some new criteria related to the oscillation of all solutions of that class of equations. The applicability and significance of the results are demonstrated with several examples.

Keywords: Oscillation; Fourth--order; Neutral differential equations.

REFERENCES

1. Agarwal, R. P., Bazighifan, O. and Ragusa, M. A., Nonlinear neutral delay differential equations of fourth—order: oscillation of solutions. *Entropy*, 23, 129, 2021.
2. Dharuman, C., Prabaharan, N., Thandapani, E. and Tunç, E., Oscillatory behavior of even—order functional differential equations with a superlinear neutral term. *Palestine Journal of Mathematics*, 12(1), 722—731, 2023.
3. Özdemir, O. and Kılıç, A., Oscillation of second-order functional differential equations with superlinear neutral terms. *Bulletin of the Malaysian Mathematical Sciences Society*, 45, 83—99, 2022.

ON FINITE FAMILIES OF HEMICONTRACTIVE MAPPINGS

Hülya Altınok¹, İsa Yıldırım¹

¹ Atatürk University, Department of Mathematics, Faculty of Science, Erzurum, Türkiye

hulya.altinok18@ogr.atauni.edu.tr

isayildirim@atauni.edu.tr

Abstract

This work explores hemicontractive mappings, which are significant in fixed point theory, a core area of functional analysis. The research focuses on the definition and properties of these mappings in Hilbert spaces. In this study, firstly, we will introduce an implicit iteration process for two finite families of hemicontractive mappings in Hilbert spaces. After, we will give a main fixed point result for such mapping under suitable conditions. Finally, we will prove our second result using a projection operator.

Keywords: Fixed Point Theory; Hemicontractive Mappings; Hilbert Space; Convergence Analysis; Iteration Method.

REFERENCES

1. Berinde, V., & Takens, F. (2007). *Iterative approximation of fixed points* (Vol. 1912, pp. xvi+-322). Berlin: Springer.
2. Kohlenbach, U. (2005). Some logical metatheorems with applications in functional analysis. *Transactions of the American Mathematical Society*, 357(1), 89-128.
3. Jailoka, P., Berinde, V., & Suantai, S. (2020). Strong convergence of Picard and Mann iterations for strongly demicontractive multi-valued mappings. *Carpathian Journal of Mathematics*, 36(2), 269-276.
4. Mărușter, L., & Mărușter, Ș. (2011). Strong convergence of the Mann iteration for α -demicontractive mappings. *Mathematical and computer modelling*, 54(9-10), 2486-2492.

Some Applications of Fuzzy Logic Using Fuzzy Data

Amine Mahzum¹, Mehmet Çitil²

¹Graduate School of Natural and Applied Science, Kahramanmaraş Sütçü İmam University,
Kahramanmaraş, Turkey

²Department of Mathematics, Science Faculty of Kahramanmaraş Sütçü İmam University,
Kahramanmaraş, Turkey

aminemahzum@gmail.com, citil@ksu.edu.tr,

Abstract

In this study, applications using fuzzy and intuitionistic data will be introduced. In line with recent developments such as artificial intelligence, we will examine the problems of managing and making decisions using fuzzy, intuitionistic fuzzy data in some areas such as regulating traffic flow and improving production in greenhouses.

Keywords: Artificial intelligence, fuzzy logic, intuitionistic fuzzy, traffic control, greenhouse.

REFERENCES

1. Zadeh, L. A. Fuzzy Sets, Information and Control, 8: 338-353, (1965)
2. Kesler, S., Karakan, A., and Oğuz, Y., Real-Time Strawberry Plant Classification and Efficiency Increase with Hybrid System Deep Learning Microcontroller and Mobile Application, Applied Science, vol. 12, no. 17, pp. 1–18, Sep. 2022.
3. Özdemir, O., and Kalınkara, Y. (2020). Fuzzy Logic: A Content Analysis for Thesis and Article Studies Between 2000-2020. Acta Infologica, 4(2), 155-174.
4. K. Atanassov, Intuitionistic fuzzy sets, in: V. Sgurev, Bd., VII ITKR's Session, Sofia, June 1983 (Central Sci. and Techn. Library, Bulg. Academy of Sciences, 1984).

A COMPARATIVE ANALYSIS OF THE PROBLEM-SOLVING PROCESSES OF PROSPECTIVE ELEMENTARY MATHEMATICS AND SCIENCE TEACHERS IN THE CONTEXT OF A REAL-LIFE PROBLEM

Himmet Korkmaz^{1a}, Ayhan Çinici^{1b}, Ali İhsan Mut²

^{1a} Department of Mathematics & Science Education, Ordu University, Ordu, Turkey

himmetkorkmaz@odu.edu.tr,

^{1b} Department of Mathematics & Science Education, Ordu University, Ordu, Turkey

ayhancinici@odu.edu.tr,

² Department of Mathematics & Science Education, Dicle University, Diyarbakır, Turkey

aliihsan.mut@dicle.edu.tr,

Abstract

The aim of this study is to examine and compare the problem-solving processes of prospective elementary mathematics and science teachers when dealing with a context-based real-life problem. In recent years, there has been a growing interest in and emphasis on the use of real-life problem situations in teaching and learning processes. It has been stated that encouraging students to produce individual or group-based solutions to problems grounded in real-life numerical data significantly supports conceptual understanding and internalization of mathematical and scientific concepts (Tan & Nie, 2015; Mebert et al., 2020). From this perspective, identifying the approaches adopted by prospective mathematics and science teachers in solving real-life problems, as well as revealing potential similarities and differences, is expected to provide valuable insights into the cognitive processes of teacher candidates with a background in quantitative disciplines.

This study adopted a qualitative research design, specifically a comparative case study approach. The participants consisted of 27 third-year prospective elementary mathematics teachers and 22 prospective science teachers enrolled in the Faculty of Education at a public university located in the Black Sea Region of Turkey. Data were collected through solution papers, audio recordings of group work, and observations. The collected data were analyzed using descriptive and content analysis methods.

Keywords: Pre-service mathematics teachers; Pre-service science teachers; Real-life problem.

REFERENCES

1. Tan, Jennifer Pei-Ling; Nie, Youyan. The role of authentic tasks in promoting twenty-first century learning dispositions. *Authentic problem solving and learning in the 21st century: Perspectives from Singapore and beyond*, 19-39, 2015.

A COLLECTION OF SOLITARY WAVES FOR THE FRACTIONAL KADOMTSEV-PETVIASHVILI-MODIFIED-EQUAL-WIDTH EQUATION

Özlem Kırıcı¹, Arif Özkul², Hasan Bulut²

¹Department of Mathematics, Kırklareli University, Türkiye

²Department of Mathematics, Fırat University, Elazığ, Türkiye

ozlem.isik@klu.edu.tr, arifozkul@gmail.com, hbulut@firat.edu.tr

Abstract

In this research, diverse wave samples of (2+1)-dimensional fractional Kadomtsev-Petviashvili-modified-equal-width (KPmEW) equation have been examined. The new version of trial equation method (NVTEM) has been handled to establish the wave solutions of the governing nonlinear model in frame of M-truncated derivative. Plentiful rational, trigonometric, exponential, hyperbolic and Jacobi elliptic type functions have been attained and these new solutions have been observed in two and three dimensions to highlight the physical demeanors. The software system Mathematica has been utilized to validate the evaluated waves. This research is marked by the novel results considering the former studies.

Keywords: Kadomtsev-Petviashvili-modified-equal-width (KPmEW) equation; The new version trial equation method; Traveling wave solutions

REFERENCES

1. Qawaqneh, H., Alrashedi, Y., Ahmad, H., Bekir, A., Discovery of exact solitons to the fractional KP-MEW equation with stability analysis. *Eur. Phys. J. Plus.* 140(4), 1-18 (2025).
2. Das, A., Ghosh, N., Bifurcation of traveling waves and exact solutions of Kadomtsev–Petviashvili modified equal width equation with fractional temporal evolution. *Comput. Appl. Math.* 38(1), 9 (2019).
3. Alomair, M. A., Analysis of Truncated M-Fractional Mathematical and Physical (2+1)-Dimensional Nonlinear Kadomtsev–Petviashvili-Modified Equal-Width Model. *Fractal and Fractional*, 8(8), 442 (2024).
4. Islam, M.T., Akter, M.A., Gómez-Aguilar, J.F., Akbar, M.A., Torres-Jiménez, J., A novel study of the nonlinear Kadomtsev–Petviashvili-modified equal width equation describing the behavior of solitons. *Opt. Quant. Electron.* 54(11), 725 (2022).
5. Seadawy, A.R., Iqbal, M., Lu, D., Applications of propagation of long-wave with dissipation and dispersion in nonlinear media via solitary wave solutions of generalized Kadomtsev–Petviashvili modified equal width dynamical equation. *Comput. Math. Appl.* 78(11), 3620-3632 (2019).

Various Optical Soliton Solutions for Combined Model Using Analytical Technique

Derya DENİZ¹ and Ebru CAVLAK ASLAN¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

deryadeniz485@yandex.com , ebrucavlak@hotmail.com

Abstract

In this paper, various exact wave solutions of the concatenation model which combines the nonlinear Schrödinger equation, the Lakshmanan-Porsezian-Daniel model, and the Sasa-Satsuma model are investigated. Using the unified method, different types of wave structures such as optical soliton, elliptic wave, and rational soliton solutions are obtained. These solutions are visualized through two- and three-dimensional plots based on appropriate parameter selections to demonstrate their physical relevance. The soliton solutions are anticipated to yield effective results, particularly in long-distance optical communication systems. The employed method provides an effective, reliable, and systematic approach for computing soliton solutions in nonlinear models. The results are expected to contribute significantly to applications in optical fiber systems, plasma physics, telecommunications, engineering, and related fields.

Keywords: Soliton; Optical Soliton; The Nonlinear Schrödinger Equation; The Lakshmanan-Porsezian-Daniel Model; The Sasa-Satsuma Model; The Unified Method.

REFERENCES

1. Ali Rehman, Mustafa Inc, Dean Chou, Magnetohydrodynamic flow of carbon nanotubes blood based hybrid nanofluids with the impact of thermal radiation over a permeable surface, Journal of Radiation Research and Applied Sciences, Vol:18, No:101–408, 2025.
2. Tarek. A. Khalil, Niveen Badra, Hamdy M. Ahmed, Wafaa B. Rabie, Optical solitons and other solutions for coupled system of nonlinear Biswas–Milovic equation with Kudryashov’s law of refractive index by Jacobi elliptic function expansion method, Optik, Vol:253, No:168–540, 2022.
3. Fan Sun, Optical solutions of Sasa-Satsuma equation in optical fibers. Optik, Vol:228, No:166–127, 2021.

PSEUDO NULL AND NULL SMARANDACHE CURVES IN E_1^3 AND THEIR CHARACTERIZATIONS

Sultan CERİTLİ¹ & Münevver YILDIRIM YILMAZ²

^{1,2}Department of Mathematics, University of Firat, Elazığ, Turkey

crtlsln46@gmail.com , munyildirim@gmail.com

Abstract

In this paper, we define pseudo null and null smarandache curves E_1^3 and examine some curvature and torsion characterizations for these curves.

Keywords: Smarandache curve, Pseudo null curve , Null curve.

REFERENCES

1. Turgut M. , Yılmaz S. , (2008), Smarandache Curves in Minkowski Space-time, International Mathematical Combinatories, 2, 30-36.
2. Bayrak N. , Bektaş Ö. , Yüce S. , (2016). Special Smarandache Curves in E_1^3 , Communications Faculty of Sciences University of Ankara Series A1: Mathematics and Statistics, 65, 2, 143-160.
3. Almaz F. , Kulahcı M. A. , Yılmaz M.Y. , (2018). A Note on Special Smarandache Curves in the Null Cone Q^3 , Acta Universitatis Apulensis , 56 , 111-124.
4. Almaz F. , Kulahcı M. A. , Yılmaz M.Y. , (2020). Approximation to the Smarandache Curves in the Null Cone , Palestine Journal of Mathematics , 9, 1, 527-536.
5. Duggal, K. L. ,& Jin, D. H. (2007). Null Curves and Hypersurfaces of Semi-Riemannian Manifolds. World Scientific.

GENERALIZED V-MANNHEIM CURVES IN E^4 AND THEIR CHARACTERIZATIONS

Gamze YAPRAK¹ & Münevver YILDIRIM YILMAZ²

¹Department of Mathematics, University of Firat, Elazig, Turkey

gamzeyprk4460@gmail.com, munyildirim@gmail.com

Abstract

In this paper we define generalized V-Mannheim curves in E^4 and also obtain some characterizations for these curves

Keywords: Mannheim curve; Frenet frame; Curvature functions.

REFERENCES

1. Matsuda, H., & Yorozu, S. (2009). On generalized Mannheim curves in Euclidean 4-space.
2. Okuyucu, O. Z. (2013). Characterizations of the quaternionic Mannheim curves in Euclidean space E^4 . *Mathematical Combinatorics*, 2, 44-53.
3. Ersoy, S., Tosun, M., & Matsuda, H. (2012). Generalized Mannheim curves in Minkowski space-time $E^{1,3}$. *Hokkaido Mathematical Journal*, 41(3), 441-461.
4. Aslan, N. K., & İlarslan, K. (2020). Some Characterizations of Generalized Null Mannheim Curves in Semi-Euclidean Space.
5. Avcı, E., & Camcı, Ç. (2022). TIMELIKE V-MANNHEIM CURVES IN MINKOWSKI 3-SPACE $E^{1,3}$. *Lapseki Meslek Yüksekokulu Uygulamalı Araştırmalar Dergisi*, 3(5), 58-71.
6. Aslan, N. K., & İlarslan, K. (2020). Some Characterizations of Generalized Null Mannheim Curves in Semi-Euclidean Space.
7. Li, Y., Uçum, A., İlarslan, K., & Camcı, Ç. (2022). A new class of Bertrand curves in Euclidean 4-space. *Symmetry*, 14(6), 1191.

ANALYSIS OF THE TRIANGLES IN THE CONTEXT OF PROTOTYPICALITY IN THE TÜRKİYE CENTURY MAARİF MODEL MATHEMATICS TEXTBOOKS

Ali İhsan Mut¹, Muhammed Ahmet Pekyen², Himmet Korkmaz³

¹Department of Mathematics & Science Education, Dicle University, Diyarbakır, Turkey

aliihsan.mut@dicle.edu.tr,

²Ministry of Education, Alitaş İmam Hatip Middle School, Ergani, Diyarbakır, Turkey

muhammedpekyen@gmail.com,

³Department of Mathematics & Science Education, Ordu University, Ordu, Turkey

himmetkorkmaz@odu.edu.tr

Abstract

Utilizing real-life contexts or concrete representations is regarded as an effective strategy for enhancing the understanding and teaching of mathematical concepts. Prototypical examples that are easily recalled and support cognitive processing can be especially effective in facilitating learning during the early phases of instruction (Tapan Broutin, 2014). However, the unconscious or uncritical use of such examples may lead to misconceptions about mathematical concepts (Akkoç, 2006; Ayaz, 2016; Aydın Karaca, 2014; Özkan, 2019; Sarı Arıkan, 2019; Zembat, 2014). The aim of this study is to analyze the triangle examples in the mathematics textbooks prepared in accordance with the Türkiye Century Maarif Model, which came into effect in 2024 and has been implemented for the first time in grades 1, 5, preparatory, and 9, in terms of whether they are prototypical or not. The scope of the study includes seven mathematics textbooks used in the 2024–2025 academic year. In this research, the document analysis method was utilized, and data were gathered through the use of a “Mathematics Textbook Analysis Form” specifically designed for this study. The collected data were analyzed using the descriptive analysis method. The findings indicate that the highest proportion of non-prototypical triangle usage (21.1%) was observed at the 5th grade level in the 2024-2025 mathematics textbooks, all of which were published by the Ministry of National Education (MoNE). In the preparatory grade mathematics textbooks, it was found that only 1 out of the 44 triangle examples used was non-prototypical.

Keywords: Prototypical examples; Prototype; Mathematics textbook; Triangles; Türkiye Century Maarif Model.

REFERENCES

1. Akkoç, H. (2006). Fonksiyon kavramının çoklu temsillerinin çağrıştırdığı kavram görüntüleri. Hacettepe Üniversitesi Eğitim Fakültesi Dergisi, 30(30), 1-10.

A NOTE ON DEMI WEAK ALMOST LIMITED OPERATORS

Gül Sinem Keleş¹, Birol Altın²,

Department of Mathematics, University of Gazi, Ankara, Turkey

gsinemkeles@gmail.com,

birola@gazi.edu.tr

Abstract

In this study, we introduce the class of demi-weak almost limited operators on a Banach lattice as a generalization of weak almost limited operators defined by Elbour A., Machrafi N., and Moussa M. in 2015. Let E be a Banach lattice, an operator $T: E \rightarrow E$ is called a demi-weak almost limited operator if for every sequence $(x_n) \subset E$ and every pairwise disjoint sequence $(f_n) \subset E'$ whenever $x_n \xrightarrow{w} 0$, $f_n \xrightarrow{w^*} 0$ and $(f_n(x_n) - f_n(T(x_n))) \rightarrow 0$ implies $f_n(x_n) \rightarrow 0$. We examine the relationship between weak almost limited operators and demi-weak almost limited operators. In addition, we establish a characterization of demi-weak almost limited operators. Finally, we obtain some properties of the class of demi-weak almost limited operators.

Keywords: Weak almost limited operators; Demi-weak almost limited operators; Limited operators; Banach lattice.

REFERENCES

1. A. Elbour, N. Machrafi, M. Moussa, On the class of weak almost limited operators, Quaestiones Mathematicae, Vol:38, No:6, 817–827, 2015.
2. C.D. Aliprantis, O. Burkinshaw, Positive Operators, Berlin, Springer, 2006.
3. C.D. Aliprantis, O. Burkinshaw, Locally Solid Riesz Spaces, New York, Academic Press, 1978.
4. G.S. Keles, B. Altın, The Class of Demi-Order Norm Continuous Operators, Gazi University Journal of Science, Vol:36, No:4, 1693 - 1698, 2023.
5. G.S. Keles, B. Altın, The Class of Demi-Strongly Order Bounded Operators, Sakarya University Journal of Science, Vol:28, No:2, 364 - 370, 2024.
6. P. Meyer-Nieberg, Banach Lattices, Berlin, Springer, 1991.
7. W.V. Petryshyn, Construction of fixed points of demicompact mappings in Hilbert space, Journal of Mathematical Analysis and Applications, Vol:14, No:2, 276-284, 1966.

Synthesis and Characterization of a Series of Poly(2-hydroxypropyl acrylate)-based Copolymers

Bedrettin Savaş^{1*}, Efkân Çatıker², Temel Öztürk³, Abdullah Karanfil²

¹Kafkas University, Kars Vocational School, 36100 Kars, Turkey

²Ordu University, Faculty of Arts and Sciences, Department of Chemistry, 52200 Ordu, Turkey

³Giresun University, Faculty of Arts and Sciences, Department of Chemistry, 28200 Giresun, Turkey

bdrtn.svs.36@gmail.com, bedrettin.savas@kafkas.edu.tr

Abstract

Poly(2-hydroxypropyl acrylate) (PHPA) is a water-soluble and thermo-responsive polymer that exhibits a lower critical dissolution temperature [1]. In this study, poly- β -alanine oligomer with vinyl end-group was synthesized using acrylamide as a monomer, sodium tert-butoxide as an initiator, and hydroquinone as a radical polymerization inhibitor via hydrogen-transfer polymerization [2]. The macromonomer and HPA were polymerized using the "grafting through" strategy via free-radical polymerization to obtain graft copolymers P(HPA-g-BA) with various degrees of grafting [3]. Similarly, N-isobutoxymethyl acrylamide (IBMAAm) and N-methoxypropyl acrylamide (MPAAm) were used as co-monomers to obtain poly(HPA-co-IBMAAm) and poly(HPA-co-MPAAm) random copolymers, respectively. The structural and thermal characterization of the products was achieved using basic polymer characterization methods such as elemental analysis, FTIR, ¹H-NMR, GPC, DSC, and TGA.

Keywords: Poly(2-hydroxypropyl acrylate), copolymer, free-radical polymerization, thermo-responsive polymer.

This study was supported by the Scientific Research Projects Coordination Unit of Kafkas University (project number: 2024-FM-33).

REFERENCES

1. Eggenhuisen T.M., Becer C.R., Fijten M.W.M., Eckardt R., Hoogenboom R., Schubert U.S. (2008). Libraries of statistical hydroxypropyl acrylate containing copolymers with LCST properties prepared by NMP. *Macromolecules* 41:5132-5140. <https://doi.org/10.1021/ma800469p>.
2. Çolakoğlu G.N., Çatıker E., Öztürk T., Meyvacı E. (2022). Synthesis and characterization of brush-type poly- β -alanine-grafted polymethyl methacrylate using "grafting through" method. *Chemical Papers* 76:869–878 <https://doi.org/10.1007/s11696-021-01908-0>.
3. Savaş B., Karanfil A., Çatıker E., Öztürk T. (2025). Synthesis and characterization of a set of novel copolymers by using RAFT polymerization method and "grafting through" strategy. *International Journal of Polymer Analysis and Characterization* (accepted). DOI: 10.1080/1023666X.2025.2502819

THE PERFORMANCE ADVANTAGE OF THE LEVY-GTO HYBRID ALGORITHM COMPARED TO GTO

Bevtullah YAĞIZ¹, Dr. Öğr. Üyesi Erdal EKER², Sinan DENİZ³, Doç. Dr. Yener ALTUN⁴

¹ Institute of Science, University of Yüzüncü Yıl, Van, Turkey

² Department of Marketing and Advertising, Muş Alparslan University, Muş, Turkey

³ Institute of Science, University of Yüzüncü Yıl, Van, Turkey

⁴ Department of Business Administration, University of Yüzüncü Yıl, Van, Turkey

beytmat@gmail.com

e.eker@alparslan.edu.tr

sinan_deniz65@hotmail.com

yeneraltun@yyu.edu.tr

Abstract

The aim of this study is to experimentally evaluate the performance of swarm-based heuristic algorithms, which are used for problem optimization within the scope of machine learning models. In this context, the Gorilla Troops Optimizer (GTO) is examined in detail. To overcome the algorithm's fundamental limitation of getting trapped in local optima, a hybrid approach incorporating the Levy flight strategy is proposed. The effectiveness of the proposed method is demonstrated through experimental studies conducted on five test functions with varying characteristics. The results show that the hybrid approach provides a performance improvement ranging between 3% and 5%. The findings are also visually supported using box plots.

Keywords: Artificial learning; Metaheuristic algorithm; Gorilla troops algorithm; Levy flight.

REFERENCES

1. Abdollahzadeh, B., Soleimanian Gharehchopogh, F., Mirjalili, S., Artificial gorilla troops optimizer: a new nature-inspired metaheuristic algorithm for global optimization problems. *International Journal of Intelligent Systems*, 36(10), 5887-5958, 2021.
2. Kusuma, P. D., Hasibuan, F.C., Enriched Coati Osprey Algorithm: A Swarmbased Metaheuristic and Its Sensitivity Evaluation of Its Strategy. *IAENG International Journal of Applied Mathematics*, 54(2), 277-285, 2024.
3. Wong, W.K., Ming, C.I., A review on metaheuristic algorithms: recent trends, benchmarking and applications. In *2019 7th International Conference on Smart Computing & Communications (ICSCC)* (pp. 1-5). IEEE, 2019.

MATHEMATICAL MODELING OF CRIMEAN-CONGO HEMORRHAGIC FEVER SPREAD: AN APPROACH BASED ON THE SIRD MODEL

Yusuf Gürsoy¹ ; Nilüfer Topsakal² ; Figen Kangalgil³

¹ Sivas Cumhuriyet University Institute of Science, Department of Mathematics, Sivas, Türkiye

²Sivas Cumhuriyet University, Faculty of Science, Department of Mathematics

³Dokuz Eylül University, Bergama Vocational School, Department of Mathematics

yusufgursoy@cumhuriyet.edu.tr, ntopsakal@cumhuriyet.edu.tr, figen.kangalgil@deu.edu.tr

Abstract

Crimean-Congo Haemorrhagic Fever (CCHF) is a serious zoonotic disease transmitted to humans through tick bites and characterised by a high case fatality rate. As an endemic disease in Turkey, it constitutes an important public health problem. In this study, a compartmental SIRD (Susceptible-Infectious-Recovered-Dead) model was used to investigate the transmission dynamics of CCHF. The model was used to analyse epidemiological data reported at Sivas Cumhuriyet University Research Hospital between 2010 and 2024. The model is formulated as a system of ordinary differential equations (ODEs) with parameters such as transmission rate (β), recovery rate (γ) and mortality attributable to disease (μ) estimated using real-world data. The basic reproductive number (R_0) was analytically derived to serve as a threshold indicator for potential outbreak or eradication of the disease. Local stability analyses of both disease-free and endemic equilibrium points were performed by linearisation techniques using the Jacobian matrix. All simulations, parameter estimates and stability analyses were performed using MATLAB. The numerical results show the temporal progression of the disease under different initial conditions and parameter scenarios. The inclusion of the deceased in the SIRD framework facilitates a more realistic portrayal of diseases characterised by significant mortality. This study demonstrates that mathematical modelling, supported by computational tools such as MATLAB, offers valuable insights into the spread of infectious diseases and provides a quantitative basis for public health intervention planning.

Keywords: Crimean-Congo Hemorrhagic Fever; SIRD model; infectious disease modeling, MATLAB; Basic reproduction number (R_0); Stability analysis; Epidemiology

This work is supported by the Scientific Research Project Fund of Sivas Cumhuriyet University under the project number F-2025-735.

A NEUTRAL SYSTEM APPROACH TO ASYMPTOTIC STABILITY OF Q-FRACTIONAL SINGULAR DELAY SYSTEMS

Yener ALTUN¹, Şakir İŞLEYEN²

¹Department of Business Administration, University of Yüzüncü Yıl, Van, Turkey

²Department of Econometrics, University of Yüzüncü Yıl, Van, Turkey

yeneraltun@yyu.edu.tr

sakirisleyen@yyu.edu.tr

Abstract

This study is concerned with the problem of asymptotic stability in q -fractional singular delay systems. By representing the singular system in a neutral form and using an augmented Lyapunov–Krasovskii functional method, we obtain a new stability criterion in terms of a linear matrix inequality (LMI). The criterion is applicable for the stability test of both singular time-delay systems and neutral systems with constant time delays. The approach used in this study relies on directly calculating the quantum derivatives of the Lyapunov–Krasovskii functionals. Illustrative examples demonstrate the applicability of the theoretical results, the effectiveness of the method, and its merits.

Keywords: Asymptotic stability; Lyapunov–Krasovskii functional; LMI; q -fractional singular delay systems.

REFERENCES

1. Al-Salam W.A., Some fractional q -integrals and q -derivatives. Proc Edin Math Soc., 15: 135-140, 1969.
2. Liu, S., Zhou, X.F., Li, X., Jiang, W., Asymptotical stability of Riemann–Liouville fractional singular systems with multiple time-varying delays. Appl. Math. Lett., 65: 32–39, 2017.
3. Yang, W., Alsaedi, A., Hayat, T., Fardoun, H.M., Asymptotical stability analysis of Riemann-Liouville q -fractional neutral systems with mixed delays. Math. Meth. Appl. Sci., 42: 4876–4888, 2019.

GLOBAL STABILITY OF AN EXPONENTIAL DIFFERENCE EQUATION SYSTEM

D. YILDIZ, M. KARA, Y. YAZLIK

Abstract

In this study, we investigate the qualitative behavior of solutions to a discrete system of exponential-type difference equations:

$$\begin{cases} u_{n+1} = \frac{a_1 + b_1 e^{-v_n} + c_1 e^{-v_{n-1}}}{d_1 + p_1 v_n + q_1 v_{n-1}}, \\ v_{n+1} = \frac{a_2 + b_2 e^{-w_n} + c_2 e^{-w_{n-1}}}{d_2 + p_2 w_n + q_2 w_{n-1}}, \\ w_{n+1} = \frac{a_3 + b_3 e^{-u_n} + c_3 e^{-u_{n-1}}}{d_3 + p_3 u_n + q_3 u_{n-1}}, \end{cases}$$

for $n \in \mathbb{N}_0$, where the coefficients $a_j, b_j, c_j, d_j, p_j, q_j, j \in \{1,2,3\}$ are non-negative real numbers such that $a_j^2 + b_j^2 + c_j^2 \neq 0$ and $d_j^2 + p_j^2 + q_j^2 \neq 0, j \in \{1,2,3\}$. The initial conditions $u_{-s}, v_{-s}, w_{-s}, s \in \{-1,0\}$ are assumed to be positive real numbers. More specifically, we study the boundedness and persistence of positive solutions, the existence and uniqueness of a positive equilibrium point, the local and global behavior of this equilibrium, and the convergence rate of the positive solutions. Finally, several numerical examples are provided to support the theoretical findings.

Keywords: Boundedness, system of difference equations, local and global stability

REFERENCES

1. Abo-Zeid, Z. (2019). Behaviour of solution of a second order rational difference equation. *Mathematica Moravica*, 23(1), 11-25. 10.5937/MatMor1901011A
2. A.Khaliq, , M.Zubeir, ve , A. Q. Khan. (2020). Asymptotic behavior of the solution of difference equation system of exponential form. *Fractals*, 28(6), 1-20.
3. Y.Akrou, , M.Kara, , N.Touafek, ve Y.Yazlik, (2021). Solution formulas for some general systems of difference equations. *Mmn.*, 22(2), 529-555. 10.181514/MMN.2021.3365
4. F.Bozkurt, (2013). Stability analysis of a nonlinear difference equation. *International Journal Of Modern Nonlinear Theory And Application*, 2, 1-6. 10.4236/ijmnta.2013.21001
5. D.S.Dilip,T.Philip(2020) Global stability,periodicity and boundedness behavior of a difference equations. *Southeast Asian Bulletin Of Mathematics*, 44, 315-323.
6. M.Gümüs, ve Z.R.Abo, (2021). Qualitative study of a third order rational system of difference equations. *Mathematica Moravica*, 25, 81-97. 10.5937/MatMor210108G
7. M.Kara, (2023). Investigation of the global dynamics of two exponential-form difference equations system. *Electronic Resaerch Archive*, 31, 6697-6724. 10.3934/era.2023338.

IDEALS AND GRÖBNER BASES OF SOME PSEUDO-SYMMETRIC NUMERICAL SEMIGROUP FAMILIES

Meral Sürer ¹ and Mehmet Şirin Sezgin ²

¹ Department of Mathematics, University of Batman, Batman, Turkey

² Institute of Graduate Studies, University of Batman, Batman, Turkey

¹ meral.suer@batman.edu.tr,

² mehmetsezgin1984@gmail.com,

Abstract

In this work, we give the ideals of some pseudo-symmetric numerical semigroup families with multiplicity 4 and embedding dimension 3, and the Gröbner bases of these ideals.

Keywords: Gröbner basis; Ideals; Numerical semigroups; Pseudo-symmetric numerical semigroups.

REFERENCES

1. Adams, Loustaunau, An Introduction to Grobner Bases, American Mathematical Society, No. 3, 1994.
2. Ene, Herzog, Gröbner bases in commutative algebra, Graduate Studies in Mathematics 130, American Mathematical Society, 2012.
3. Rosales, Branco, Irreducible numerical semigroups, Pacific Journal of Mathematics, 209, 131–143, 2003.
4. Rosales, García-Sánchez, Numerical Semigroups, Springer, 2009.

*Abstract Submission should be prepared only **1 page**.

THE ANALYSIS OF LGS SUCCESS VIA ROUGH SET THEORY: THE CASE OF SİVEREK AND SAVUR

Berfin Karadağlı¹, Saliha Erboğa¹, Nurettin Bağirmaz²

¹The Institute for Graduate Educational Studies, Mardin Artuklu University, Mardin, TÜRKİYE

²Mardin Artuklu University, Mardin, TÜRKİYE

berfinkaradagli19@gmail.com,

salihaerboga850@gmail.com,

nurettinbagirmaz@artuklu.edu.tr,

Abstract

The aim of this presentation is to analyze the placement results of the 2025 High School Entrance Exam (LGS) candidates into various types of high schools using Rough Set Theory, considering both environmental and hereditary factors. In this context, personal, familial, and academic data were collected from 8th grade students in the Siverek and Savur districts. Based on the collected data, the most discriminative combinations of factors affecting students' LGS scores will be identified and analyzed within the framework of Rough Set Theory. As a result, the main factors influencing students' exam success will be examined in greater depth and presented to relevant governmental and academic stakeholders.

Keywords: Rough Set Theory; Success of LGS.

REFERENCES

1. Pawlak, Z. (1982). "Rough Sets", International Journal of Computer and Information Sciences, vol. 11, pp. 341–356.
2. Suraj, Z. (2004). "An Introduction to Rough Set Theory and Its Application". ICENCO'2004, Cairo, Egypt, December 27–30.
3. Yüce, G. S. (2021). Sosyal Medya Kullanım Alışkanlıklarının Kaba Kümeler Yardımıyla İncelenmesi. Unpublished MSc thesis, İnönü University, Malatya.
4. Suna, H. E., Tanberkan, H., Gür, B., Perc, M., et al. (2020). Akademik Başarının Yordayıcıları: Sosyoekonomik Düzey ve Okul Türü. Journal of Economy Culture and Society, (61), 41–64.

*Abstract Submission should be prepared only **1 page**.

GLOBAL EXISTENCE, DECAY AND NONEXISTENCE OF SOLUTIONS FOR LAMÉ TYPE EQUATION WITH A FRACTIONAL TIME DELAY TERM

Hasan KANDEMİR¹, Erhan PİŞKİN²

¹Dicle University, Institute of Natural and Applied Sciences, Diyarbakır, Türkiye

²Dicle University, Department of Mathematics, Diyarbakır, Türkiye

¹hasankan210@gmail.com, ²episkin@dicle.edu.tr

Abstract In this presentation, we consider the Lamé equation with a fractional type time delay term and a nonlinear source terms. We show the global existence and decay of the solutions under suitable conditions. Also, we prove the nonexistence of global solutions.

Keywords: Global existence, decay, nonexistence

REFERENCES

- 1) A. Benaissa, G. Soumia, Stability result of the Lamé system with a delay term in the internal fractional feedback, *Acta Universitatis Sapientiae* (2021): 336.
- 2) B. Mbodje, Wave energy decay under fractional derivative controls, *IMA Journal of Mathematical Control and Information*, 23.2 (2006): 237-257.
- 3) E. Pişkin, *Evolüsyon Denklemlerin Çözümlerinin Patlaması*, Pegem Akademi Yayıncılık, 2022.
- 4) E. Pişkin, *Evolüsyon Denklemlerin Çözümlerinin Azalması*, Pegem Akademi Yayıncılık, 2025.
- 5) A. Radhouane, A. Benaissa, A. Zraï, Blow-up and asymptotic behavior for a wave equation with a time delay condition of fractional type, *Rendiconti del Circolo Matematico di Palermo Series 2*, 70 (2021) 1061-1081.

FRACTIONAL HADAMARD'S TYPE INEQUALITIES FOR EXPONENTIALLY CO-ORDINATED CONVEX FUNCTIONS

Ali Yılmaz¹, Abdullah Akkurt¹ and Hüseyin Yıldırım¹

¹Department of Mathematics, Faculty of Sciences, Kahramanmaraş Sütçü İmam University, Kahramanmaraş, Turkey

yilmazali0556@gmail.com, abdullahmat@gmail.com, hyildir@ksu.edu.tr

Abstract

In this study, we introduce a novel definition of convexity in the context of exponential coordinates. Utilizing this new framework, we establish a series of fractional Hermite-Hadamard type inequalities. Furthermore, we provide several remarks to elucidate the connections between our findings and previously established results in the literature.

Keywords: Convex functions; co-ordinated convex mapping; Hermite-Hadamard inequality; fractional integral.

REFERENCES

1. Azpeitia, A.G., 1994. Convex functions and the Hadamard inequality. Rev. Colomb. Math. 28, 7--12.
2. Bakula, M.K., Pecaric J., 2004. Note on some Hadamard-type inequalities. J. Inequal. Pure Appl. Math. 5 (3), article 74.
3. Dragomir, S.S., 2001. On Hadamard's inequality for convex functions on the co-ordinates in a rectangle from the plane. Taiwanese J. Math. 4, 775--788.
4. Dragomir, S.S., Pearce, C.E.M., 2000. Selected Topics on Hermite--Hadamard Inequalities and Applications. RGMIA Monographs, Victoria University
5. Samko, S.G., Kilbas, A.A., Marichev, O.I., 1993. Fractional Integrals and Derivatives, Theory and Applications. Gordon and Breach, Yverdon et alibi.
6. Sarikaya, M.Z., 2014. On the Hermite--Hadamard-type inequalities for co-ordinated convex function via fractional integrals. Integr. Transf. Spec. Funct. 25, 134--147.
7. Sarikaya, M.Z., 2015. Some inequalities for differentiable co-ordinated convex mappings. Asian-Eur. J. Math. 8 (21 pages).

Evaluation of Biomechanical and Physiological Effects of Vehicle Vibrations on Pregnant Drivers

Özden Taşdemir¹, Muhammed Emin Kula², Ömer Gündoğdu³

^{1,2,3}Department of Mechanical Engineering, University of Atatürk, Erzurum, Turkey

oo.zdn.yldz@gmail.com¹, mekula27@gmail.com², omergun@atauni.edu.tr³

Abstract

The human body, which is highly sensitive to vibrations, can react differently to these effects according to different population groups (pregnant women, children, elderly people, etc.). In this study, a biodynamic model of a pregnant woman, a highly sensitive user group, was developed, and the vibrations affecting the targeted body parts were analyzed. While extant research in this field has predominantly focused on the effects of vibration on the dynamic systems of non-pregnant individuals, studies examining the impact of vertical vibrations on pregnant women, particularly in the sitting position under driving conditions, are severely limited. In this study, a biodynamic model of a pregnant woman, a highly sensitive user group, was developed and the vibrations affecting the targeted body parts were analyzed. In this context, the study investigates the impact of vibrations induced by speed bumps on highway safety, employing a half-vehicle model with a pregnant woman driver's seat as a case study. The analysis evaluated the forces acting upon the pregnant woman's lumbar region and the vertical accelerations experienced by the driver's head and fetus. The physical model of a pregnant woman is represented using mechanical system elements (spring, mass and damper). The model has eleven degrees of freedom and is mathematically expressed in terms of Newton-Euler and moment equations. The second-order linear ordinary differential equations with constant coefficients were transformed into state-space form and solved using MATLAB software. The results of the study are presented in graphical form, and the dynamic effects on the pregnant driver and the fetus are assessed in detail.

Keywords: Biodynamic model; Pregnant woman; Vibration analysis.

ON SOME PERFECT NUMERICAL SEMIGROUPS

Mehmet Sait ALAKUŞ¹

Department of Mathematics, University of Dicle, Diyarbakır, TURKEY,
alakusmehmetsait@gmail.com

Sedat İlhan²

Department of Mathematics, University of Dicle, Diyarbakır, TURKEY,
sedati@dicle.edu.tr

Abstract

A numerical semigroup consists of linear combinations of some natural numbers that are prime to each other. Numerical semigroups are particularly prominent with their applications in commutative algebra. However, perfect numerical semigroups are a subject worth examining. It is particularly important to express numerical semigroups in terms of type sequence elements. In previous studies, the structure of numerical semigroups with determine number four or less than four has been expressed in terms of elements of their type sequence. In this study, we will examine the structures of some perfect numerical semigroups with a number of determinants of five. Firstly, we will give the necessary and sufficient conditions for the type sequences of these numerical semigroups. Secondly, we will show that these numerical semigroups are perfect.

Keywords: Numerical semigroup, Frobenius number, isolated gaps, type sequence, perfect numerical semigroup, genus.

A COMPARATIVE STUDY OF ITERATIVE AND NON-ITERATIVE OPTIMIZATION METHODS FOR NONLINEAR LEAST SQUARES PROBLEMS

Sinan DENİZ¹, Beytullah YAĞIZ², Doç. Dr. Yener ALTUN³, Doç. Dr. Murat CANAYAZ⁴

¹ Institute of Science, University of Yüzüncü Yıl, Van, Turkey

² Institute of Science, University of Yüzüncü Yıl, Van, Turkey

³ Department of Business Administration, University of Yüzüncü Yıl, Van, Turkey

⁴ Department of Computer Engineering, Van Yuzuncu Yil University, Van, Turkey

sinan_deniz65@hotmail.com

beytmat@gmail.com

yeneraltun@yyu.edu.tr

mcanayaz@yyu.edu.tr

Abstract

In this study, the numerical characteristics of classical iterative methods such as Newton-Raphson, Gradient Descent, Gauss-Newton and Levenberg-Marquardt, which are widely used in solving nonlinear least squares problems, have been examined. These gradient-based methods are sensitive to initial values. In this regard, the logarithmic transformation method, used for determining initial parameter estimates is presented in detail. The performance of the classical methods has been compared with alternative optimization algorithms that do not require derivative information, including deterministic, heuristic, and evolutionary approaches. To support the comparison, a application involving the Gompertz function was conducted, and each method was analyzed in terms of convergence behavior, solution accuracy, and sensitivity to initial values.

Keywords: Newton-Raphson; Gradient Descent; Gauss-Newton; Levenberg-Marquardt; Deterministic; Heuristic; Evolutionary.

REFERENCES

1. Goldberg, D. E., *Genetic Algorithms in Search, Optimization, and Machine Learning*. Addison-Wesley, Reading, MA, 1989.
2. Marquardt, D. W., An algorithm for least-squares estimation of nonlinear parameters. *Journal of the Society for Industrial and Applied Mathematics*, 11(2), 431-441, 1963.
3. Yonar, A., Yonar, H., An Efficient Hybrid Algorithm with Particle Swarm Optimization and Nelder-Mead Algorithm for Parameter Estimation of Nonlinear Regression Modeling. *Gazi University Journal of Science*, 35(2), 716-729, 2022.

Strength Performance of Alkali Activated Mortars Produced with Waste Brick Powder Depending on NS/NH Ratio and Curing Temperature

Fahreddin Melih Dursun¹, Ali Öz^{1,2}

¹ Department of Civil Engineering, University of Atatürk, Erzurum, Turkey

²Narman Vocational School, Atatürk University, Erzurum, Turkey

fahreddinmelihdursun@gmail.com

Abstract

This study investigates the effects of waste brick powder (BP) use in alkali-activated composites (AAC) on mechanical performance and strength behavior under the sulfate effect. Sodium silicate/sodium hydroxide (NS/NH) ratios of 1.5, 2.0, 2.5 and 3.0 were determined and curing temperatures were applied at 40°C and 80°C in mixtures prepared by substituting 15% BP in GBFS-based binder system. Waste marble powder (WMP) was used as fine aggregate. Experimental findings revealed that the NS/NH ratio and curing temperature strongly affected the compressive strength. In particular, under the NS/NH=2.5 ratio and 80°C curing condition, the BP-based mixtures reached the highest compressive strength of 44.58 MPa. This was explained by the acceleration of geopolymerisation reactions and the formation of a denser C-A-S-H/N-A-S-H gel matrix. Under normal curing conditions (40°C), the compressive strength of the mixture with an NS/NH ratio of 2.5 was measured as 30.32 MPa, which is about 47% lower than the 80°C curing condition. This finding indicates high-temperature curing significantly increases reactivity and binder phase formation in BP-containing systems. Furthermore, regarding sulfate resistance, the compressive strength of the mixtures with BP exposed to MgSO₄ solution for 60 days decreased by less than 15% for the mixture with NS/NH = 2.5. This indicates that BP provides physical stability in the binder matrix formed after alkaline activation and develops resistance to sulfate ions. As a result, using BP offers significant potential for both sustainable building material production and durability. NS/NH=2.5 ratio and 80°C curing condition are recommended to achieve optimum performance in such systems.

Keywords: Sustainability, brick powder, Heat cure, Sulfate effect, NS/NH ratio

On Asymptotically Equivalent Sequences in Probability

Kübra Elif AKBAŞ¹ and Mahmut IŞIK²

¹Faculty of Medicine, Firat University, Elazığ, Turkey; keet@firat.edu.tr

²Faculty of Education, Harran University, Şanlıurfa, Turkey; misik63@yahoo.com

Abstract. In this study, we give some results related with asymptotically equivalent sequences in probability.

Keywords: Sequence of random variables, Asymptotical statistical equivalence.

REFERENCES

- [1] Akbaş, KE and Isık, M. On asymptotically λ -statistical equivalent sequences of order α in probability, Filomat 34(13) (2020), 4359-4365.172.
- [2] Akbaş, KE and Isık, M. Weighted Statistical Convergence in Probability, 8th International Conference on Computational Mathematics and Engineering Sciences 17-19 May 2024, Şanlıurfa - Türkiye.
- [3] Fast, H. Sur la convergence statistique, Colloq. Math. 2 (1951) 241-244.
- [4] Fridy, JA. On statistical convergence, Analysis 5(4) (1985), 301-313.
- [5] Ghosal, S. Weighted statistical convergence of order α and its applications, J. Egyptian Math. Soc. 24(1) (2016), 60-67.
- [7] Salat, T. On statistically convergent sequences of real numbers, Math. Slovaca 30 (1980), 139-150.
- [8] Schoenberg, IJ. The integrability of certain functions and related summability methods, Amer. Math. Monthly 66 (1959), 361-375.
- [9] Steinhaus, H. Sur la convergence ordinaire et la convergence asymptotique, Colloq. Math. 2 (1951), 73-74.

k -Lauricella Hypergeometric Functions

Ayşegül Kızılarşlan, Ayşegül Çetinkaya

Department of Mathematics, University of Kırşehir Ahi Evran, Kırşehir, Turkey

kizilarşlan.aysegul@ogr.ahievran.edu.tr, acetinkaya@ahievran.edu.tr

Abstract

The Pochhammer k -symbol, introduced by Diaz and Pariguan in 2007, has become one of the important symbols of the theory of special functions. There are many special functions defined using this symbol. In this study, k -Lauricella hypergeometric functions are defined by means of Pochhammer k -symbol. Also, some Euler type integral formulas are given.

Keywords: Pochhammer symbol; Integral representation; Lauricella hypergeometric functions.

REFERENCES

1. Diaz, R., Pariguan, E. (2007). On hypergeometric functions and Pochhammer k -symbol. *Divulgaciones Matemáticas*, 15(2), 179–192.
2. Exton, H. (1976). *Multiple hypergeometric functions and applications*. John Wiley & Sons, New York.
3. Halıcı, S., Çetinkaya, A. (2024). k -Srivastava hypergeometric functions and their integral representations, *Miskolc Mathematical Notes*, 25 (2), 685–697.
4. Hasanov, A., Srivastava, H. M. (2006). Some decomposition formulas associated with the Lauricella function F_A^n and other multiple hypergeometric functions, *Applied Mathematics Letters*, 113-121.
5. Kıymaz, I. O., Çetinkaya, A., Agarwal, P. (2017). A study on the k -generalizations of some known functions and fractional operators. *Journal of Inequalities and Special Functions*, 8(4), 31-41.
6. Manocha, H. L., Srivastava, H. M. (1984). *A Treatise on generating functions*. John Wiley & Sons, New York.
7. Mubeen, S., Rehman, A. (2014). A note on k -Gamma function and Pochhammer k -symbol. *J. Math. Sci*, 6(2), 93-107.
8. Mubeen, S., Habibullah, G. M. (2012). An integral representation of some k -hypergeometric functions. In *Int. Math. Forum* 7(4), 203-207.

AN ANALYSIS OF 9th GRADE MATHEMATICS TEXTBOOK PREPARED IN ACCORDANCE WITH TÜRKİYE CENTURY MAARİF MODEL CURRICULUM IN THE CONTEXT OF CONGRUENCE AND SIMILARITY

Ali İhsan Mut¹, Şeyda Bircan²

¹Department of Mathematics & Science Education, Dicle University, Diyarbakır, Turkey

aliihsan.mut@dicle.edu.tr,

²Bandırma District Directorate of National Education, Balıkesir, Turkey

seydaekmn@gmail.com

Abstract

Coxford (1991) highlights congruence and similarity as fundamental elements of Euclidean geometry. Çevik (2006) notes that congruence and similarity axioms and theorems are employed in the proofs of several theorems in plane geometry, such as the basic proportionality theorem, the sine theorem, and the angle bisector theorem, among others. As also emphasized by Çıldır (2007), despite the importance of the theme of congruence and similarity within the discipline of mathematics, inconsistencies in the terminology used to describe these concepts in textbooks and scholarly works are evident. In an effort to address these inconsistencies, mathematicians like Hilbert have sought to establish alternative axiom systems. This controversial situation in the history of mathematics, along with the differences between Hilbert and Euclid, has been reflected in scientific literature, textbooks, teachers, and, consequently, students as well. The aim of this study is to analyze the theme of congruence and similarity in terms of content structure, the terminology used, and the instructional approach as presented in the 9th grade mathematics textbook prepared in accordance with the Türkiye Century Maarif Model and implemented during the 2024–2025 academic year. The study adopts the document analysis method. As research material, the 9th grade mathematics textbook published within the scope of the Türkiye Century Maarif Model and introduced into the curriculum as of the 2024–2025 academic year was used. The data obtained were analyzed using content analysis methodology.

Keywords: Türkiye Century Maarif Model; Congruence and similarity; 9th grade mathematics textbook, Document analysis.

REFERENCES

1. Coxford Jr, A. F. (1991). Geometry from Multiple Perspectives. Curriculum and Evaluation Standards for School Mathematics Addenda Series, Grades 9-12. National Council of Teachers of Mathematics, Inc., 1906 Association Drive, Reston, VA 22091.

EXISTENCE AND UNIQUENESS OF SOLUTIONS FOR FRACTIONAL BOUNDARY VALUE PROBLEMS INVOLVING CAPUTO-HADAMARD FRACTIONAL DERIVATIVES

Dila Karataş¹ Fulya Yörük Deren¹

¹ Department of Mathematics, University of Ege, Izmir, Turkey

¹ Department of Mathematics, University of Ege, Izmir, Turkey

karatasdila01@gmail.com, fulya.yoruk@ege.edu.tr

Abstract: Fractional-order derivative models are gaining popularity over ordinary derivatives because they provide more accurate results in real-world applications. Researchers are increasingly studying fractional differential equations due to their effectiveness in fields like engineering, mathematics, physics, bioengineering, and applied sciences [1]. Nonlinear fractional boundary value problems are particularly important, as they better model complex phenomena. This study focuses on analyzing such nonlinear fractional boundary value problems. In this work, the following nonlinear fractional boundary value problem is considered with the help of Caputo-Hadamard fractional derivative operator.

$$-D^{\alpha}u(t) = f(t, u(t)), t \in (1, e), u'(1) = 0, \beta D^{\alpha-1}u(e) + u(\eta) = 0.$$

Here $f \in C([1, e] \times [0, \infty), [0, \infty))$, $1 < \alpha \leq 2$, $\beta > 0$, $1 < \eta < e$, D^{α} and $D^{\alpha-1}$ are Caputo-Hadamard fractional derivatives of order α and $\alpha-1$ respectively. Under some assumptions on the nonlinear term, the existence and uniqueness of positive solutions are obtained by certain fixed point theorems.

Keywords: Banach fixed point theorem, Caputo-Hadamard fractional derivative, fractional differential equations.

ACKNOWLEDGEMENT

This work was supported by TÜBİTAK (Scientific and Technological Research Council of Türkiye) under the 2209-A Project (Project No:1919B012336540)

REFERENCES

1. Kilbas, A. A., Srivastava, H. M., & Trujillo, J. (2006). Theory and Applications of Fractional Differential Equations. In: North-Holland Mathematics Studies, Vol. 204, Elsevier Science B. V.

An Experimental Study on the Durability of One-Part Geopolymer Foam Concrete Under NaCl Effect

İbrahim BOUBACR MOUNKAILA¹, Gökhan Kaplan¹

¹Department of Civil Engineering, University of Atatürk, Erzurum, Turkey

gkaplan@atauni.edu.tr

Abstract

In this study, the strength performance of geopolymer foam concretes produced for the development of environmentally friendly building materials was investigated under saline conditions (NaCl solution). Composites containing lime and pumice powder in different ratios (0%, 25%, 50%, 75%) were used as binder system and 10% and 20% sodium bicarbonate was preferred for activation. All samples were immersed in 5% NaCl solution after the 90-day curing period was completed and kept in this environment for 3 months. At the end of the curing period, the specimens were dried and their compressive strength was determined in accordance with ASTM C109 standard. According to the experimental results obtained under normal curing conditions, a significant improvement in compressive strength was observed as the pumice content increased. The highest strength was obtained in P50NC20 specimen with 10.22 MPa after 90 days of standard curing, which indicates that pumice increases the pozzolanic contribution and matrix integrity in the binder system. It was observed that contact with NaCl solution caused significant deterioration in the mechanical properties of geopolymer foam concretes. Mixtures containing only lime (P0NC10 and P0NC20) experienced strength losses of 76.24% and 73.05%, respectively, and very low residual strengths of 0.24-0.45 MPa were recorded. This was attributed to insufficient geopolymeric network formation and high porosity. In contrast, mixtures containing 25% and 50% pumice showed significantly better strength. In particular, the P50NC20 mix was the most resistant to NaCl environment, reaching a value of 7.89 MPa. In terms of relative strength losses, P50NC10 and P50NC20 mixtures retained 74.49% and 77.20% of their initial strength, respectively. These findings show that pumice powder provides significant contributions to the stabilization of the geopolymer matrix and that mechanical strength can be maintained under aggressive environmental conditions such as NaCl, especially with moderate pumice substitution. In this respect, the study offers a potential alternative for applications requiring chemical resistance such as infrastructure elements or structural elements that may be exposed to seawater.

Keywords: NaCl effect, Foam concrete, Sustainability, Pumice powder, Slaked lime.

Conceptual Perceptions of Prospective Teachers Regarding the Concept of "Unit" in Mathematics

Nejla GÜREFE¹

¹ Department of Mathematics Education, Faculty of Education, Mersin University, Mersin, Turkey

nejlagurefe@mersin.edu.tr

Abstract

Meaningful learning and internalization of the concept of measurement is largely associated with understanding the concept of unit. This concept establishes a direct bridge between the object or phenomenon to be measured and the measurement. It can be said that the knowledge of prospective teachers of the future regarding this concept, which is at the basis of many mathematical subjects such as arithmetic, algebra and measurement, is quite important. It has been claimed that when an understanding is carefully sought in a wrong answer given by a person, the logical side of that answer can be discovered. In this case, the ways in which a person learns mathematics and the obstacles that block these ways can be understood, and this can be a guide on how mathematical knowledge is produced, encoded in the mind and used. In this sense, it can be said that determining the mathematical understanding of the individual is an important guide for conceptual learning. The aim of this study is to determine the understanding of prospective mathematics teachers regarding the concept of "unit". The participants of this descriptive survey study were prospective teachers receiving education at a state university in the south of Turkey. Data were collected through documents and document analysis was used in its analysis. As a result of the findings, it was seen that the prospective teachers explained the concept of unit with different concepts such as "measurement", "the essence of the subject", "each part of the whole", "the smallest part of the multitude".

Keywords: Unit; Conception; Math Prospective Teachers

REFERENCES

1. Hiebert, J. (1981). Units of measure: Results and implications from national assessment. *Arithmetic Teacher*, 28(6), 38–43.
2. Webber, C. (2004). From errors to conceptions an approach to student diagnosis. In J. C. Lester, R. M. Vicari, & F. Paraguacu (Eds.), *Intelligent tutoring systems, lecture notes in computer science*. 3220, (pp 710–719). Springer, https://doi.org/10.1007/978-3-540-30139-4_124
3. Niss, M. (1999). Aspects of the nature and state of research in mathematics education. *Educational Studies in Mathematics*, 40(1), 1–24. <https://doi.org/10.1023/A:1003715913784>

ON THE BOUNDEDNESS AND STABILITY PROPERTIES OF SOLUTIONS FOR FOURTH ORDER NONLINEAR STOCHASTIC DIFFERENTIAL EQUATIONS WITH VARIABLE DELAYS

Zozan OKTAN¹ and Cemil TUNÇ²

Department of Mathematics, Faculty of Sciences, Van Yuzuncu Yil University, 65080, Van, Turkey

¹ [E-mail:zozanoktan34@gmail.com](mailto:zozanoktan34@gmail.com)

² [E-mail: cemtunc@yahoo.com](mailto:cemtunc@yahoo.com)

Abstract

This study addresses the analysis of fourth-order stochastic differential equations with variable delays. Two novel results are established, providing sufficient conditions for the stochastic asymptotic stability and stochastic boundedness of solutions. The proofs are developed through the application of the Lyapunov–Krasovskii functional technique. To substantiate the theoretical contributions, two illustrative examples are presented. In comparison with existing studies, the results obtained herein are original and offer a significant contribution to the qualitative theory of stochastic differential equations.

Keywords: Bounded variable delay, Fourth-order, LK-F, Nonlinear stochastic differential equation (SDE), Stability

REFERENCES

- [1] Arnold, L., 1974. *Stochastic differential equations: Theory and applications*, Jhon Wiley & Sons, 1974.
- [2] Abou-El-Ela, A. M.A. Sadek, A. I., Mahmoud, A. M. and Taie, R.O.A., On the stochastic stability and boundedness of solutions for stochastic delay differential equation of the second order, *Chin.J. Math.* Art. ID 358936, 8 PP, 2015.
- [3] Tunç, C., Oktan, Z., Improved new qualitative results on stochastic delay differential equations of second order. *Comput. Methods Differ. Equ.* 12(1),67-76, 2024. <https://doi.org/10.22034/cmde.2022.52821.2229>

ON THE STABILITY OF FIFTH-ORDER NONLINEAR STOCHASTIC DELAY DIFFERENTIAL EQUATIONS

Zozan OKTAN¹ and Cemil TUNÇ²

Department of Mathematics, Faculty of Sciences, Van Yuzuncu Yil University, 65080, Van, Turkey

¹[E-mail: zozanoktan34@gmail.com](mailto:zozanoktan34@gmail.com)

²[E-mail: cemtunc@yahoo.com](mailto:cemtunc@yahoo.com)

Abstract

In this work, we examine the stochastic asymptotic stability of solutions to fifth-order non-autonomous stochastic delay differential equations with multiple variable delays, utilizing the Lyapunov–Krasovskii functional method. A new theorem is established to characterize this qualitative property. To illustrate the effectiveness of the proposed approach, a detailed example is presented. The result obtained in this work offers a broader generalization compared to several existing findings in the current literature.

Keywords: Stochastic delay differential equations, asymptotically stability in probability, multiple variable delay, fifth-order, Lyapunov-Krasovskii functional

REFERENCES

- [1] Mahmoud, A. M., Bakhit, D. A. M. Study of the stability behaviour and the boundedness of solutions to a certain third-order differential equation with a retarded argument. *Annals of Applied Mathematics* 35.1 99-110, 2019.
- [2] Mao, X., *Stochastic differential equations and applications*. Elsevier, 2007.
- [3] Tunç, C., Oktan, Z., Improved new qualitative results on stochastic delay differential equations of second order. *Comput. Methods Differ. Equ.* 12(1),67-76, 2024. <https://doi.org/10.22034/cmde.2022.52821.2229>
- [4] Tunç, C., New results on the stability and boundedness of nonlinear differential equations of fifth order with multiple deviating arguments. *Bulletin of the Malaysian Mathematical Sciences Society*, 36(3), 671-682, 2013.
- [5] Tunç, C., Oktan, Z., Stability and boundedness of stochastic integro-delay differential equations. *J. Math. Anal.* 15, no. 5, 69-83, 2024.

Pell Statistical Convergence

Koray İbrahim Atabey¹, Mikail Et², Muhammed Çınar³

¹ Muş Nizamülmülk Girl Anatolia İmam Hatip High School, Muş, Turkey

²Department of Mathematics, University of Fırat, Elazığ, Turkey

³Department of Mathematics, University of Muş Alparslan, Muş, Turkey

korayatabey7@gmail.com,

mikail68@gmail.com,

muhammedcinar23@gmail.com

Abstract

The Pell matrix $P = (P_{nk})$ is used in this paper to introduce the concepts of Pell summability, Pell density function, and Pell statistical convergence. Additionally, we provide some inclusion relations regarding these concepts and strongly $P[p]$ -Cesaro summability.

Keywords: Pell analog; Pell numbers; Statistical Convergences.

REFERENCES

1. Hüseyin Aktuğlu, Şerife Bekar, On q -Cesaro matrix and q -statistical convergence, Journal of Computational and Applied Mathematics, 235(16), 4717–4723, 2011.
2. Koray İbrahim Atabey, Hemanta Kalita, Mikail Et, On Pell sequence spaces, Palestine Journal of Mathematics, (Inpress), 2025.
3. Koray İbrahim Atabey, Muhammed Çınar, Mikail Et, q -Fibonacci statistical convergence, Georgian Mathematical Journal, 2025.<https://doi.org/10.1515/gmj-2025-2022>.

NUMERICAL CALCULATIONS USING NEWMARK'S METHOD TO SOLVE A HYPERBOLIC PROBLEM

Adem Irmak¹, Arzu Aykut², Merve Özer³, Sıdıka Şule Şener Kılıç⁴

¹ Department of Mathematics, University of Atatürk, Erzurum, Turkey

² Department of Mathematics, University of Atatürk, Erzurum, Turkey

³ Department of Mathematics, University of Atatürk, Erzurum, Turkey

⁴ Department of Mathematics, University of Atatürk, Erzurum, Turkey

airmak@bayburt.edu.tr, aaykut@atauni.edu.tr, merve.ozel19@ogr.atauni.edu.tr,
senersule@atauni.edu.tr,

Abstract

In this study, the Newmark method is preferred for the solution of the differential equation that occurs together with the Galerkin method in the solution of hyperbolic problems. The weak solution of the hyperbolic equation is examined using the Galerkin and Newmark methods under different boundary conditions. This study is prepared to obtain numerical solutions for the weak solution of a hyperbolic problem with variable coefficient homogeneous boundary conditions.

Keywords: Newmark method, Hyperbolic equation, Galerkin Methods

REFERENCES

1. Bardella, L. and Genna, F., 2005. Newmark's Time Integration Method From the Discretization of Extended Functionals, *Journal of Applied Mechanics*, 72, 527-537.
2. Chiba, F., Kako, T., 1988, On the Stability and Convergence of Newmark's β method, *Institute of Statistical Mathematics Cooperative Research Report*, 110, 196–202.
3. Dupont, T., 1973, L2-Estimates For Galerkin Methods For Second Order Hyperbolic Equations, *SIAM Journal of Numerical Analysis*, 10, 5, 1973.
4. Farlow, S. 1982, *Partial Differential Equations for Scientists and Engineers*, Dover Publication New York
5. Fung, T.C., 1999, On The Accuracy of Galerkin Methods in the Time Domain, *Jornal of Vibration and Control*, 5, 155-174.
6. Grote, M.J., Schneebeli, A., 2006, Dominik Schötzau, Discontinuous Galerkin finite element method for the wave equation, *SIAM J. Numer. Anal.* 44, 6, 2408–2431.
7. Idesman, A. V., 2007, A new high-order accurate continuous Galerkin method for linear elastodynamics problems, *Comput Mech*, 40, 261–279.
8. Kreyszig, E., 1999, *Advanced Engineering Mathematics*, John Wiley&Sons, Inc., New York.

ON THE SEPARATION AXIOM OF TOPOLOGICAL SPACES M-SPACE AND T_M -SPACE

GHASSAN MOHAMMED ABDULLAH ALOMAISI¹, NURETTİN BAĞIRMAZ²

Abstract

In the study of topology, spaces are often classified based on their structural properties and the types of functions they support. One such interesting class is the **M-space**, which presents unique characteristics that differentiate it from more conventional topological spaces. A **M-space** is defined by a set of axioms that impose specific conditions on the space's topology, focusing on continuity, separation, and compactness properties. The **T_M -space**, a specific type of **M-space**, further refines these properties by introducing additional constraints that influence the relationships between points and open sets. This paper explores the fundamental aspects of both **M-spaces** and **T_M -spaces**, providing detailed examples and comparing them with more widely studied spaces such as Hausdorff spaces (T_2) and regular spaces (T_3). The study also highlights the relevance of **M-spaces** and **T_M spaces** in both theoretical and applied contexts, including their potential applications in modern fields such as quantum topology, data science, and network theory. The goal of this research is to advance the understanding of these spaces and their potential impact on the broader landscape of topology.

Keywords: Topology, Separated Axiom, M-Space, T_M -space

REFERENCES

- [1] Aslım, G., Genel Topoloji, Ege Üniv., Fan Fak., Yay . no: 109 izmir 1988.
- [2] Koçak, M., Genel Topolojiye giriş ve Çözümlü Alıştırmalar,
- [3] Cantor, G. (1874). "Über unendliche, lineare Punktmannigfaltigkeiten" (Mathematische Annalen)
- [6] Hausdorff, F. (1914). *Grundzüge der Mengenlehre.*,
- [7] Munkres, J. R. (2000). *Topology (2nd edition).*

¹ The institute for Graduate Educationl studes / Mardin Artuklu University/ Mardin/ Turkey

ghassan@artuklu.edu.tr

² Department of Mathematics, University of Mardin Artuklu, Turkey

nurettinbagirmaz@artuklu.edu.tr

ANALYZING A NONLINEAR SCHRÖDINGER TYPE EQUATION: DYNAMICAL ANALYSIS AND SOLITARY WAVE FEATURES

Elif Yaşar¹

¹Department of Mathematics, Faculty of Arts and Science, Bursa Uludag University,
Bursa, Turkey

elifyasar@uludag.edu.tr,

Abstract

This study explores exact soliton solutions of a nonlinear Schrödinger type equation, which governs femtosecond pulse propagation in optical fibers with higher-order dispersive and nonlinear effects. Using an advanced analytical approach, we derive bright and dark soliton solutions that exhibit amplitude and width variations under different nonlinear parameter regimes. The obtained solutions demonstrate how third-order dispersion and nonlinear terms influence pulse dynamics, offering potential applications in ultrafast optics and signal processing. Physical interpretations of these solitons are discussed and propagation characteristics in nonlinear media. The results complement existing theories on optical solitons [1-3] while providing new insights into ultrashort pulse control.

Keywords: a nonlinear Schrödinger type equation, soliton.

REFERENCES

1. M. J. Potasek, M. Tabor, Exact solutions for an extended nonlinear Schrödinger equation, Phys. Lett. A, 154, 449–452, 1991.
2. Y. Kodama, Optical solitons in a monomode fiber, J. Stat. Phys. 39, 597–614, 1985.
3. P. Kumari, R.K. Gupta, S. Kumar, K.S. Nisar, Doubly periodic wave structure of the modified Schrodinger equation with fractional temporal evolution, Results Phys., 33, 105128, 2022.

EULER METHOD IN NUMERICAL SOLUTION OF FRACTIONAL DIFFERENTIAL EQUATIONS

Ahu ERCAN¹, Zehra ÖZCANOĞLU¹

¹Department of Mathematics, University of Firat, Elazig, Turkey

zehraozcanoglu@gmail.com

Abstract

In this study, an approach based on Euler method is presented for the numerical solution of fractional order differential equations. In the study, fractional derivatives formulated under Caputo definition are taken into consideration, and how the classical Euler method is adapted and its applicability is detailed in this framework. In addition, the accuracy of the method are analyzed through sample problems. The findings show that Euler method can provide an effective starting point in solving fractional systems despite its simple structure.

Keywords: Fractional Differential Equations; Caputo Derivative; Euler Method.

REFERENCES

1. Diethelm, K., The analysis of fractional differential equations, Springer, Heidelberg, 2010.
2. Diethelm, K., An efficient parallel algorithm for the numerical solution of fractional differential equations, Fract. Calc. Appl. Anal. 84 (3) (2011), 475–490.
3. Tomasek, P. : On Euler Methods for Caputo Fractional Differential Equations. Archivum Mathematicum 59 (2023),287-294

SECOND HANKEL DETERMINANT FOR A SUBCLASS OF BI-UNIVALENT FUNCTIONS ASSOCIATED WITH GEGENBAUER POLYNOMIALS AND SUBORDINATION

Fatma SAĞSÖZ¹

¹ Department of Mathematics, Atatürk University, Erzurum, Turkey

faltuntas@atauni.edu.tr

Abstract

Recently, many upper bounds of second Hankel determinant $|a_2a_4 - a_3^2|$ have been investigated for various subclasses of bi-univalent functions. The main object of this work is to evaluate an upper bound for a novel subclass of bi-univalent functions associated with the Gegenbauer polynomials that satisfy subordination conditions on the open unit disk U . Some relevant remarks and observations are also presented.

Keywords: Analytic functions; Bi-univalent functions; Subordination; Gegenbauer polynomials; Second Hankel determinant.

REFERENCES

1. Abdelbaset Zeyani, Abdulmtalb Hussen, Second Hankel determinant for a bi-univalent function subclass based Gegenbauer (ultraspherical) polynomials, European Journal of Pure and Applied Mathematics, Vol:18, No:2, 1-17, 2025.
2. Christian Pommerenke, Univalent Functions, Vandenhoeck and Ruprecht, Göttingen, 1975.
3. James Waring Noonan, Derek Keith Thomas, On the second Hankel determinant of areally mean p -valent functions, Transactions of the American Mathematical Society, Vol: 223, 337–346, 1976.
4. Peter L. Duren, Univalent Functions, Grundlehren der Mathematischen Wissenschaften, Band 259, Springer-Verlag, New York, 1983.
5. Ulf Grenander, Gabor Szegő, Toeplitz Forms and Their Applications, California Monographs in Mathematical Sciences, University of California Press, Berkeley, 1958.

Production of Foam Concrete with Hasankale Pumice and Expanded Micronized Perlite

Bera Aydın¹, İsmail Yıldız²

¹Department of Civil Engineering, University of Ataturk, Erzurum, Turkey

bera.aydin14@ogr.atauni.edu.tr,

²Department of Civil Engineering, University of Ataturk, Erzurum, Turkey

ismail.yildiz@atauni.edu.tr

Abstract

In this study, the mechanical and physical properties of foam concrete, a type of lightweight concrete, were investigated. For this purpose, cement, pumice, expanded micronized perlite aggregates and foaming agent were used for foam concrete production. A total of 4 different concrete mixes were produced. In all 4 groups, the amount of foaming agent was kept constant. In the first group, cement, water and foaming agent were used. In the second group, cement, water, pumice and foaming agent were used. In the third group, cement, water, perlite and foaming agent were used. In the last group, cement, water, pumice, perlite and foaming agent were used. Density, compressive and flexural tests were performed on the produced samples. The unit weight of the fresh concrete mixtures was measured. After 28 days of curing under laboratory conditions, compressive and flexural strength tests were performed on the samples. As a result of the unit volume weight test, foam concrete samples between 720 kg/m³ and 1400 kg/m³ were produced. The lowest unit volume weight was obtained in the samples produced with perlite and the highest unit volume weight was obtained in the samples where pumice and perlite aggregates were used together. As a result of the 28-day compressive strength test, the lowest value was obtained in the mixture containing perlite with 6,33 MPa and the highest value was obtained in the normal foam concrete mixture without aggregates with 12,47 MPa. This is due to the fact that the compressive strength of cement is higher than the other aggregates. As a result of the 28-day flexural strength test, the lowest value was obtained with 2,04 MPa in the mixture where pumice and perlite were used together and the highest value was obtained with 3,0 MPa in the mixture containing pumice. As a result, the use of pumice and expanded perlite aggregates in foam concrete is used for a structure that is both lighter and has better thermal insulation properties rather than improving the mechanical properties of foam concrete.

Keywords: Foam Concrete; Pumice; Expanded Micronized Perlite; Compressive and Flexural Strength

ON GENERALIZED PROPORTIONAL FRACTIONAL INTEGRAL

Yasin Kaya¹

¹Department of Mathematics, University of Dicle, Diyarbakır, Turkey

ykaya@dicle.edu.tr,

Abstract

Fractional calculus operators deal with investigation and applications of integrals and derivatives of arbitrary (real or complex) order. There are many definitions of fractional integral and fractional derivatives of different types. Object of this talk is to present an introductory overview of the theory of an integral operator of fractional calculus known as generalized proportional fractional integral. We also show a fractional integral inequality whose proof based on techniques to the existing literature.

Keywords: Generalized proportional fractional integral; Inequality; Operator.

REFERENCES.

1. Aljaaidi, T. A., Pachpatte, D. B., Abdeljawad, T., Abdo, M. S., Almalahi, M. A., & Redhwan, S. S. (2021). Generalized proportional fractional integral Hermite–Hadamard’s inequalities. *Advances in Difference Equations*, 2021(1), 493.
2. Alzabut, J., Sudsutad, W., Kayar, Z., & Baghani, H. (2019). A new Gronwall–Bellman inequality in frame of generalized proportional fractional derivative. *Mathematics*, 7(8), 747.
3. Jarad, F., Abdeljawad, T., & Alzabut, J. (2017). Generalized fractional derivatives generated by a class of local proportional derivatives. *The European Physical Journal Special Topics*, 226, 3457-3471.
4. Neamah, M. K., & Ibrahim, A. (2021). Generalized proportional fractional integral inequalities for convex functions. *AIMS Mathematics*, 6(10), 10765-10777.
5. Rahman, G., Abdeljawad, T., Khan, A., & Nisar, K. S. (2019). Some fractional proportional integral inequalities. *Journal of Inequalities and Applications*, 2019, 1-13.

STRESS ANALYSIS OF A ROTATING FG POLAR ORTHOTROPIC DISK

Oğuzhan Demirel¹, Durmuş Yarımpaçlı¹

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

demirelemu@hotmail.com, durmusyarpabuc@osmaniye.edu.tr

Abstract

In this paper, the elastic analysis of a functionally graded polar orthotropic disk rotating with time-dependent angular velocity is considered. The material properties of the disk and thickness are exponentially graded in the radial direction. Under these conditions, a linear partial differential equation is obtained. The time dependence in the partial differential equation is eliminated by using the Laplace transform. Then, the boundary value problem in the radial direction is solved by the pseudospectral Chebyshev Method in the Laplace domain. The stresses in physical space are obtained by using the modified Durbin method. The effects of thickness, rotation and material properties on the stresses are analyzed.

Keywords: Elastic stresses, Polar orthotropic disk, Functionally graded materials, Rotation..

REFERENCES

1. U. Güven, A. Çelik and C. Baykara, On transverse vibrations of functionally graded polar orthotropic rotating solid disk with variable thickness and constant radial stress, J. of Reinforced Plastics and Composites, Vol:23, No:12, 1279-1284, 2004.
2. V. Yıldırım, Numerical/analytical solutions to the elastic response of arbitrarily functionally graded polar orthotropic rotating discs, J. of the Brazilian Society of Mechanical Sciences and Engineering, Vol:40, No:6, 320-341, 2018.
3. X. L. Peng and X. F. Li, Elastic analysis of rotating functionally graded polar orthotropic disks, International J. of Mechanical Sciences, Vol:60, No:1, 84-91, 2012.
4. L. Sondhi, A. K. Thawait, S. Sanyal and S. Bhowmick, Stress and deformation analysis of variable thickness clamped rotating disk of functionally graded orthotropic material, Materials Today: Proceedings, Vol:18, 4431-4440, 2019.
5. L. Sondhi, A. K. Thawait, S. Sanyal and S. Bhowmick, Stress and deformation analysis of functionally graded varying thickness profile orthotropic rotating disk, Materials Today: Proceedings, Vol:33, 5455-5460, 2020.

DEFORMATION ANALYSIS OF FG LONGITUDINAL RODS BY PSEUDOSPECTRAL CHEBYSHEV METHOD

Özlem Cerit¹, Durmuş Yarımpaşu¹

¹ Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

ozlemcerit08@gmail.com, durmusyarpasbu@osmaniye.edu.tr

Abstract

Deformation analysis of functionally graded longitudinal vibrated rod with variable cross-sectional area in axial direction is considered. It is assumed that the rod subjected to external harmonic excitation and material properties such as elastic modulus and density are graded in the axial direction by the Halpin-Tsai homogenization model. These conditions result in a partial differential equation with variable coefficients that is difficult to solve with conventional analytical methods. Under the Laplace transform, the partial differential equation is transformed into a time-independent boundary value problem in the longitudinal direction and solved by the pseudospectral Chebyshev Method. The displacements in physical space are obtained inverse modified Durbin Method. The effects of time dependent sinusoidal force on displacement distributions are discussed.

Keywords: Deformation analysis; Tip mass; Functionally graded materials; pseudospectral Chebyshev Method; Laplace transform, modified Durbin Method.

REFERENCES

1. Li, Q. S., Exact solutions for free longitudinal vibration of non-uniform rods. J. Sound Vib. 234(1):1–19, 2000.
2. Shatalov, M., Joubert, S., Fedotov, I., Lekalakala, M., Bidie, J., Dynamics of a longitudinally vibrating rod with variable length and cross-section, 24th International Congress on Sound and Vibration (ICSV24), London, 23-27 July 2017.
3. Yardimoglu, B., Aydin, L., Exact longitudinal vibration characteristics of rods with variable crosssections. Shock and Vibration 18: 555 -562, 2011.
4. Celebi, K., Keles, I., Tutuncu, N., Exact solutions for forced vibration of non-uniform rods by Laplace transformation. Gazi University Journal of Science 24(2): 347- 353, 2011.
5. Akgoz, B., Civalek, Ö., Longitudinal vibration analysis of strain gradient bars made of functionally graded materials (FGM). Comp.: Part B: Eng. 55: 263- 268, 2013.
6. Celebi, K., Yarımpaşu, D., Baran, T., Forced vibration analysis of inhomogeneous rods with non-uniform cross-section, Journal of Engineering Research, 6(3), 189-202, 2018.

THERMAL EFFICIENCY OF FGM ANNULAR FINS

Gazal Okşas¹, Ali Yıldırım², Durmuş Yarımabuç¹

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

²Osmaniye Vocational School, Osmaniye Korkut Ata University, Osmaniye, Turkey

gzloksas15@gmail.com, aliyildirim@osmaniye.edu.tr, durmusyarimpabuc@osmaniye.edu.tr

Abstract

The annular fins, which are used to enhance heat transfer from the cylindrical surface to the surrounding, are widely used too many applications such as heat exchangers and electronic cooling systems. The efficiency of annular fins is defined as the ratio of the heat transferred from a cylindrical surface through the fin to the heat transferred from the same fin with infinite thermal conductivity (the temperature everywhere on the fin being equal to the fin base temperature). The efficiency of annular fins varies depending on the fin material and geometry (shape, cross-section of the fin, and whether the fin is perpendicular or parallel to the surface to which the heat will be transferred). In this situation, fin efficiency can be increased not only with the fin material but also with changes in its geometry. The pseudospectral Chebyshev method is used in this study to numerically analyze the thermal efficiency of functionally graded variable cross-section annular fins. The material properties have been assumed to gradually vary in the radial direction in the fins, which is thought to have reached thermal equilibrium. The validation of the method has been ensured by comparing it with the results of an analytical study in the literature.

Keywords: Variable cross-section annular fin; Functionally graded material; Pseudospectral Chebyshev method

REFERENCES

1. Gardner, K. A., Efficiency of extended surface, Trans. ASME, 67, 621-631, 1945.
2. Kraus, A. D., Aziz, A., Welty, J. R., Extended surface heat transfer, John Wiley, New York, 2001.
3. Koizumi, M., FGM activities in Japan, Composites Part B, 28B, 1-4, 1997
4. Ullmann, A., Kalman, H., Efficiency and optimized dimensions of annular fins of different cross-section shapes, Int. J. Heat Mass Transfer, 32(6), 1105-1110.
5. Yıldırım, A., Yarımabuç, D., & Celebi, K. (2019). Thermal stress analysis of functionally graded annular fin. *Journal of Thermal stresses*, 42(4), 440-451.

THERMAL ANALYSIS OF FG POROUS FIN BY ADOMIAN DECOMPOSITION METHOD

Kerem Kaçar¹, Durmuş Yarımpaçlı¹

¹Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

proof56@hotmail.com, durmusyarpabuc@osmaniye.edu.tr

Abstract

In this paper, nonlinear thermal analysis of functionally graded convective-radiative porous longitudinal fins using Adomian decomposition method is discussed. The variation of material properties, which are generally neglected in functionally graded materials, with temperature is taken into account. The effects of the axial variation of the heat transfer coefficient as well as its dependence on temperature on the temperature distribution is discussed.

Keywords: Longitudinal porous fin, Heat conduction, Functionally graded materials, Adomian decomposition method.

REFERENCES

1. Adomian, G., A review of the decomposition method in applied mathematics, Journal of Mathematical Analysis and Applications, Vol:135, No:3, 501-544, 1988.
2. Keerthi, M.L., Gireesha, B.J., Analysis of entropy generation in a fully wet porous moving longitudinal fin exposed to convection and radiation: homogeneous and functionally graded materials, Journal of Thermal Analysis and Calorimetry, Vol:148, 6945-6957, 2023.
3. Ramesh, G.K., Manohar, G.R., Madhukesh, J.K., Venkatesh, P., Gireesha, B.J., Thermal aspects of a radiative-convective semi-spherical porous fin of functionally graded material, Eur.Phys.J.Plus, Vol:139, 97, 2024.
4. Riasat, S., Amna Huda, S., Insight into variable thermophysical features of the functionally graded porous annular fin influenced by internal heat and thermal radiation, Applied Thermal Engineering, Vol:253, 123-694, 2024.

APPLICATION OF INVERSE LAPLACE TRANSFORM METHODS ON HEAT CONDUCTION EQUATION

Zuhal Bilir¹, Mehmet Ali Sabihaoglu¹, Durmuş Yarımpanu¹

¹ Department of Mathematics, Osmaniye Korkut Ata University, Osmaniye, Turkey

zuhal.bilir1997@gmail.com, malisabihaoglu@gmail.com,
durmusyarimpabuc@osmaniye.edu.tr

Abstract

In this study, the one-dimensional heat conduction equation of the plate is solved using the combined method. The time dependence of the equation is eliminated using the Laplace transform. Then, the pseudospectral Chebyshev method is used to solve the boundary value problem in the Laplace space. The temperature distribution of the plate in the physical space is obtained by different inverse Laplace transform methods. A numerical simulation is performed to check the accuracy of the inverse Laplace transform methods.

Keywords: Inverse Laplace transform method, Heat conduction, Pseudospectral Chebyshev method.

REFERENCES

1. M. Cohen, Numerical Methods for Laplace Transform Inversion, Springer, Berlin, Germany 2007.
2. Belomestny, D., Mai, H., Schoenmakers, J., Generalized Post–Widder inversion formula with application to statistics, Journal of Mathematical Analysis and Applications, Vol:455, No:1, 89-104, 2017.
3. Sungur, B., Keles, I., Efficiency of the Gaver-Stehfest Method Transient Response of a Spiral Fin, European Mechanical Science, Vol:5, No:4, 194-200, 2020.
4. Fatoorehchi H., An accurate numerical method for inversion of Laplace transforms with applications in process dynamics and control, *Can J Chem Eng.*, Vol:99, 1374–1389, 2020.

PFAFFIAN FORMULATION OF NONLINEAR DIFFERENTIAL EQUATIONS

Barış Yapışkan¹

¹Department of Physics, Mimar Sinan Fine Arts University, Istanbul, Turkey

baris.yapiskan@msgsu.edu.tr,

Abstract

Nonlinear differential equations are fundamental in mathematically describing a diverse range of physical processes occurring in nature. Due to their intrinsic difficulty, the derivation of solutions for these equations remains a significant and actively investigated research area. Determinant-based methods [1-3] represent a prominent technique for deriving soliton solutions of nonlinear evolution equations. In this talk, we introduce the Pfaffian formulation [4] for nonlinear differential equations and discuss some recent applications of this approach.

Keywords: Soliton solutions; Pfaffian formulation; Nonlinear evolution equations.

REFERENCES

1. J. Satsuma, A Wronskian representation of N-soliton solutions of nonlinear evolution equations, J. Phys. Soc. Jpn., 46, 359-360, 1979.
2. N. C. Freeman, J. J. C. Nimmo, Soliton solutions of the Korteweg-de Vries and the Kadomtsev-Petviashvili equations: the Wronskian technique, Proc. R. Soc. London, A389, 319–329, 1983.
3. A. Nakamura, A bilinear N-soliton formula for the KP equation, J. Phys. Soc. Jpn., 58, 412-422, 1989
4. R. Hirota, Soliton solutions to BKP equation I, The Pfaffian technique, J. Phys. Soc. Jpn., 58, 2285-2296, 1989.

*Abstract Submission should be prepared only **1 page**.

ROBUSTNESS 2.0: A TRANSFORMATIONAL STORY BETWEEN ARTIFICIAL INTELLIGENCE AND STATISTICS

Esra Pamukçu¹

¹Department of Statistics, University of Firat, Elazig, Turkey

epamukcu@firat.edu.tr,

Abstract

Robust statistics has been a significant research area since the 1960s, offering methods designed to address challenges such as outliers, model misspecifications, and small sample sizes. Robust approaches developed by pioneers like Tukey, Huber, and Hampel have substantially enhanced the reliability of statistical analyses for decades. However, with the rise of machine learning and artificial intelligence, the advantages of big data and powerful algorithms overshadowed the popularity of robust methods. During this period, handling outliers became largely limited to data cleaning and regularization techniques.

Today, new challenges—including adversarial attacks, explainable artificial intelligence (XAI) requirements, and model reliability—have brought the concept of "robustness" back into focus. Nevertheless, existing literature often discusses "adversarial robustness" without connecting it to classical robust statistics, while statisticians have shown limited engagement with artificial intelligence applications. This situation has created an interdisciplinary methodological gap.

This study presents an exploratory review of the historical development of robust statistical approaches (Robustness 1.0), their transformation during the machine learning era (Robustness 1.5), and their new definitions and applications in the age of artificial intelligence (Robustness 2.0). This classification and the historical framework describing the evolution of robustness are, for the first time, introduced with these specific terms and conceptualized to contribute to the literature. Additionally, the future roles of robustness and proposed methodological directions for researchers will be discussed. The study aims to bridge the disciplines of statistics and artificial intelligence through a methodological framework grounded in the philosophy of robust thinking, offering a new conceptual perspective.

Keywords: Robustness; Artificial Intelligence; Statistics

ON GAUSSIAN CURVATURE OF THE CYCLIC SURFACES

Muhittin Evren Aydın,¹ Nesibe Tüzün Yılmaz²

¹Department of Mathematics, University of Firat, Elazığ, Turkey

²Institute of Natural and Applied Sciences, Firat University

meaydin@firat.edu.tr, nesibe_tuzun@windowslive.com

Abstract

In this talk, we analyze equations of Gauss curvature type for cyclic surfaces using the Mathematica program. In particular, we study such equations in the case of constant curvature.

Keywords: Gauss curvature; cyclic surface; rotational surface.

REFERENCES

1. R. Lopez, Cyclic hypersurfaces of constant Gauss curvature, Houston Journal of Mathematics, Vol:27, No:4, 799-805, 2001.
2. M. E. Aydın, R. Lopez, Translators of flows by powers of the Gauss curvature, Annali di Matematica Pura ed Applicata (1923 -), Vol:202, 235–251, 2023.
3. M. E. Aydın, R. Lopez, Rotational surfaces in $\mathbb{R}^4$ with new approaches and examples, International Electronic Journal of Geometry, Vol:17, No:1, 97–105, 2024.

NUMERICAL SOLUTION of GRLW EQUATION by CUBIC HERMITE B-SPLINE COLLOCATION METHOD

Sibel Özer¹, Yusuf Uçar¹, Halil Beyazıt¹

¹Department of Mathematics, University of Inonu, Malatya, Turkey

sibel.ozer@inonu.edu.tr, yusuf.ucar@inonu.edu.tr, halilbeyazt@gmail.com

Abstract

This study involves the numerical solution of the Generalized Regularized Long Wave (GRLW) equation, which has an important place in fluid dynamics. For this purpose, the Crank-Nicolson scheme is used for time discretization. The Rubin-Graves type linearization technique is used to deal with nonlinear terms. The space discretization is performed by cubic Hermite B-spline collocation method. To verify the accuracy of the proposed method, it is applied to some well-known test problems, the error norms and the conservation constants are calculated. The calculated error norms and conservation constants are presented in tables and compared with the work of previous researchers using the same parameters. In addition, the problem is presented visually by drawing graphs of the numerical results obtained. The tables and graphs show that the proposed method agrees with the analytical solutions of the test problems and preserves the conservation constants.

Keywords: GRLW equation; Collocation method; Cubic Hermite B-spline.

REFERENCES

1. T. Roshan, A Petrov-Galerkin method for solving the generalized regularized long wave (GRLW) equation, Computers and Mathematics with Applications, Vol:63, No:5, 943-956, 2012.
2. R. Mohammadi, Exponential B-spline collocation method for numerical solution of the generalized regularized long wave equation, Chinese Physics B, Vol:24, No:5, 050206, 2015.
3. H. Zeybek, S.B.G. Karakoç, A numerical investigation of the GRLW equation using lumped Galerkin approach with cubic B-spline. SpringerPlus, Vol:5, No:1, 199, 2016.
4. A. Kumari, V.K. Kukreja, Study of generalized regularized long wave equation via septic Hermite collocation method with Crank-Nicolson and SSP-RK43 schemes to capture the various solitons, Wave Motion, Vol:122, 103188, 2023.

DEVELOPMENT OF A DOMAIN-SPECIFIC AI PLATFORM BASED ON SMALL LANGUAGE MODELS

Sedat Karadağ¹, Senem Şahan Vahaplar¹, M. Hakkı Ersoy¹, Ahmet Feyzioğlu²

¹Experilabs (SahaBT Yazılım), R&D Center, İstanbul, Türkiye

²Department of Mechanical Engineering, Marmara University, İstanbul, Türkiye

senem.vahaplar@experilabs.com

Abstract

Large Language Models (LLM) have demonstrated remarkable success in natural language processing (NLP) tasks. However, their high computational requirements, prolonged training durations and data privacy concerns cause significant limitations for organizational adoption. These limitations are especially critical in regulated sectors such as finance, healthcare and government, where local data processing, compliance and customizability are essential. This study presents an AI platform based on domain-specific Small Language Models (SLM), enhanced with Retrieval-Augmented Generation (RAG) and optimized for Turkish NLP tasks. The proposed solution enables low-cost deployment, fast fine-tuning and high accuracy while reducing the need for technical expertise through a natural language query interface. The models are optimized using quantization, pruning, knowledge distillation techniques and include a custom tokenizer designed for Turkish morphological structures. Moreover, the platform's on-premise deployment capability, along with advanced data privacy techniques such as homomorphic encryption and differential privacy, ensures secure processing of sensitive data.

Keywords: Natural Language Processing (NLP); Small language models (SLM); Retrieval-augmented generation (RAG).

REFERENCES

1. Aykut Arda Akın, Mehmet Duran Akın, Zemberek: An Open Source NLP Framework for Turkic Languages, Structure, Vol: 10, 1-5, 2007.
2. Kadir Tohma, Yakup Kutlu, Challenges Encountered in Turkish Natural Language Processing Studies, Natural and Engineering Sciences, Vol: 5, No: 3, 204–211, 2020.
3. Fali Wang, Zhiwei Zhang, Xianren Zhang, Zongyu Wu, TzuHao Mo, Qiuha Lu, Wanjin Wang, Rui Li, Junjie Xu, Xianfeng Tang, Qi He, Yao Ma, Ming Huang, Suhan Wang. A Comprehensive Survey of Small Language Models in the Era of Large Language Models: Techniques, Enhancements, Applications, Collaboration with LLMs, and Trustworthiness. Journal of the ACM, Vol: 37, No: 4, Article 111, 2018.
4. Arkaitz Zubiaga, Natural Language Processing in the Era of Large Language Models, Frontiers in Artificial Intelligence, Vol: 6, 1350306, 2024.
5. Patrick Lewis, Ethan Perez, Aleksandra Piktus, Fabio Petroni, Vladimir Karpukhin, Naman Goyal, Heinrich Küttler, et al., Retrieval-Augmented Generation for Knowledge-Intensive NLP Tasks, 34th Conference on Neural Information Processing Systems (NeurIPS 2020), Vancouver, Canada, 2020.

INVESTIGATION OF WIND DIRECTIONS BY CIRCULAR DATA ANALYSIS

Yıldırım Demir¹, Çetin Görür², Saniye Demir³

¹Department of Statistics, University of Van Yüzüncü Yıl University, Van, Turkey

²Department of Econometrics, University of Van Yüzüncü Yıl University, Van, Turkey

³Department of Soil Science, University of Tokat Gaziosmanpaşa University, Tokat, Turkey

ydemir@yyu.edu.tr, cetingorur@yyu.edu.tr, saniye.140100@gmail.com

Abstract

In this study, the wind directions in Yozgat Basin were analysed with the circular data analysis method using daily wind direction data between 1969-2018. Unlike classical statistics methods, circular approaches take into account the angular nature of directional variables such as wind direction. The data set was divided into ten five-year periods and descriptive circular statistics (mean direction, resultant length, mean resultant length, circular variance, circular standard deviation and Von Mises concentration parameter (κ)) were calculated for each period. The findings show that wind directions are not randomly distributed throughout the year, but tend to cluster in certain directions during certain periods. Especially in the period 2009-2018, there is a strong directional concentration evidenced by increases in the mean resultant length and κ . Statistical tests such as Watson-Williams F, Concentration Homogeneity and Likelihood Ratio confirm that there are significant differences in mean direction and concentration parameter between periods ($p < 0.05$). These long-term changes in wind direction affect not only meteorological aspects, but also critical environmental and economic processes such as soil erosion, seeding, transport of air pollution, spread of forest fires and renewable energy potentials. This study highlights the value of circular statistics in environmental research and provides insights for regional planning and resource management.

Keywords: Environmental planning; Circular data analysis; Wind direction; Von Mises

REFERENCES

1. Demir, Y. *Examining Periodic Differences of Suicide Cases with Circular Data Analysis*. East J Med. 26(4), 580-588, 2021.
2. Fisher, N. I. *Statistical Analysis of Circular Data*. Cambridge University Press. 1993.
3. Jammalamadaka, S. R., SenGupta, A. *Topics in Circular Statistics*. World Sci. 2001
4. Mardia, K. V., Jupp, P. E. *Directional Statistics*. Wiley. 2000.

FINITE DIFFERENCE METHOD FOR LINEAR VOLTERRA INTEGRO-DIFFERENTIAL EQUATIONS

Zelal Temel¹

¹ Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

drzelaltemel@hotmail.com,

Abstract

We study a useful numerical method for the linear Volterra integro-differential equations with non-local conditions. The implicit difference principles for the differential component and the composite right-sided rectangle approach for the integral component are used to model the numerical solutions of the problem across a uniform mesh. The numerical approach's convergence and stability are next examined. The accuracy of the suggested approach is confirmed by the presented numerical experiments.

Keywords: Numerical method; Stability; Uniform mesh.

REFERENCES

1. Sumit Kumar, Sunil Kumar, Jesus Vigo-Aguiar, Analysis of a Nonlinear Singularly Perturbed Volterra Integro-Differential Equation, Journal of Computational and Applied Mathematics, 404, 113410, 2022.
2. Zelal Temel, Musa Cakir, A Robust Numerical Method for a Singularly Perturbed Semilinear Problem with Integral Boundary Conditions, Contemporary Mathematics, 5, 446-464, 2024.
3. Abdul Majid Wazwaz, Nonlinear Volterra Integro-Differential Equations: Linear and Nonlinear Integral Equations, Berlin: Springer, 2011.

Multi-layer Artificial Neural Networks in Optimization of Predictive Maintenance Processes

Okan Dardağan¹, Hakan Altunay¹

¹Department of Industrial Engineering, Süleyman Demirel University, Isparta, Turkey

okandardagan@gmail.com, hakanaltunay@sdu.edu.tr

Abstract

The prevention of breakdown and the optimization of maintenance processes are crucial for operational efficiency and cost effectiveness in industrial systems. This study proposes an Artificial Neural Network (ANN) based approach to model the complex and non-linear relationships within equipment sensor data, in contrast to classical predictive maintenance approaches. The proposed methodology is supported by a multi-layer ANN architecture. This architecture is designed to process time series data from the automotive industry for parameters that are fundamental to monitoring the health of industrial equipment, such as vibration, temperature and oil analysis. The ANN model provides high accuracy in failure prediction and flexibility in maintenance scheduling by learning the deterioration signals of the equipment. Preliminary results suggest the model outperforms conventional techniques in early-stage failure detection sensitivity and may substantially reduce maintenance costs. The results show that the ANN approach plays a critical role in the shift to reliable and predictable production systems by improving maintenance strategies.

Keywords: Predictive Maintenance; Artificial Neural Network; Artificial Intelligence; Industrial Engineering.

REFERENCES

1. Bangalore, P., Tjernberg, L. B., An artificial neural network approach for early fault detection of gearbox bearings. IEEE Transactions on Smart Grid, Vol:6 No:2, 980-987, 2015.
2. Rajpal, P. S., Shishodia, K. S., Sekhon, G. S., An artificial neural network for modeling reliability, availability and maintainability of a repairable system, Reliability Engineering & System Safety, Vol:91 No:7, 809-819, 2006.
3. Al-Naggar, Y.M., Jamil, N., Hassan, M.F., Yusoff, A.Z., Condition Monitoring Based on IoT for Predictive Maintenance of CNC Machines, Procedia CIRP, Vol:102, 314-318, Malaysia, 2021.
4. Kanawaday, A., Sane, A., Machine Learning for Predictive Maintenance of Industrial Machines Using IoT Sensor Data, 8th IEEE International Conference on Software Engineering and Service Science (ICSESS), 24-26 November, China, 87-90, 2017.

Analytical Solution of a Caputo Fractional-Order Computer Virus Propagation Model

Erkan Günerhan¹, Hatıra Günerhan², Ercan Çelik^{3,4}

¹Department of Computer engineering, Kafkas University, Kars, Turkey

erkan.gunerhan@gmail.com

²Department of Mathematics, Faculty of Education, Kafkas University, Kars, Turkey

gunerhanhatira@gmail.com,

³ Department of Applied Mathematics and Informatics, Kyrgyz-Turkish Manas University, Bishkek, Kyrgyzstan

⁴Department of Mathematics, Atatürk University, Erzurum, Turkey

ercan.celik@manas.edu.kg

Abstract

In this study, a proposed model describing the propagation of a computer virus in the network with antidote in a vulnerable system is analyzed. Using the Laplace Adomian Decomposition Method (LADM), the model's analytical and approximate-analytical solutions are determined. These solutions are found in the form of fast converging series that portray the system dynamics accurately. The efficiency of the method was tested and the validity of the introduced fractional-order model was proved through the numerical simulations.

Keywords: Caputo fractional derivative; Computer Virus Propagation; Laplace Adomian decomposition; Nonlinear system.

REFERENCES

1. Khanh NH, Huy NB. Stability analysis of a computer virus propagation model with antidote in vulnerable system. *Acta Mathematica Scientia* (2016) 36:49–61.
2. Yuan H, Chen G. Network virus-epidemic model with the point-to-group information propagation. *Appl Math Comput*, 2008, 206: 357–367.
3. Hua Y, Junjie W, Guoqing C. Infection function for virus propagation in computer network: An empirical study. *Tsinghua Science and Technology*, 2009, 14(5): 669–676.

NUMERICAL APPROACH OF SINGULARLY PERTURBED DELAY DIFFERENTIAL EQUATIONS WITH NONLOCAL CONDITIONS

Zelal Temel¹

¹ Department of Mathematics, University of Van Yuzuncu Yil, Van, Turkey

drzelaltemel@hotmail.com,

Abstract

The numerical solution of singularly perturbed delay differential equations with nonlocal conditions is presented in this study. The method's stability and convergence analysis are examined. The presented method is used to solve several numerical instances, display the problem solution graphs, and compare the calculated result with the exact solution.

Keywords: Singularly Perturbed Problem; Delay differential equations; Nonlocal Conditions.

REFERENCES

1. Mohan K. Kadalbajoo, Kapil K. Sharma, Numerical analysis of singularly perturbed delay differential equations with layer behavior, Applied Mathematics and Computation, 157.1,11-28, 2004.
2. Govindarao Lolugu, Jugal Mohapatra, A second-order numerical method for singularly perturbed delay parabolic partial differential equation, Engineering Computations, 36.2, 420-444, 2019.

On the Domain of 4-dimensional Catalan Matrix in the Space of Absolutely Summable Double Sequences

Sezer ERDEM¹

¹Department of Basic Engineering Sciences, Malatya Turgut Özal University, Malatya, Türkiye, sezer.erdem@ozal.edu.tr,

Serkan DEMİRİZ²

²Department of Mathematics, Tokat Gaziosmanpaşa University, Tokat, Türkiye, serkan.demiriz@gop.edu.tr,

Abstract

The aim of this study is to obtain a new double sequence space as the domain of the 4-dimensional Catalan matrix obtained by using Catalan numbers on the space $\mathcal{L}_u$ of absolutely summable double sequences, to investigate some algebraic and topological properties of this space, to calculate its α -, β (bp)- and λ -duals and to characterize matrix transformations from the newly obtained space to classical sequence spaces and vice versa.

Keywords: Catalan numbers; 4-dimensional Catalan matrix; Double sequence spaces.

REFERENCES

1. C.R. Adams, On non-factorable transformations of double sequences, Proc. Natl. Acad. Sci. USA, 19(5), 564567, 1933.
2. M. Ilkhan, A new conservative matrix derived by Catalan numbers and its matrix domain in the spaces c and c_0 , Linear and Multilinear Algebra, 68:2, 417434, 2020.
3. M. Ilkhan, E.E. Kara, Matrix transformations and compact operators on Catalan sequence spaces, J. Math.Anal.Appl., 498, 2021.
4. F. Moricz, B.E. Rhoades, Almost convergence of double sequences and strong regularity of summability matrices, Math. Proc. Cambridge Philos. Soc., 104(1988), 283294, 1988.
5. S. Erdem, S. Demiriz, 4-Dimensional Catalan Matrix and Its Associated Sequence Space, 10th International Eurasian Conference on Mathematical Sciences and Applications (IECMSA-2021), August 25-27, 2021, Sakarya- TURKEY.

An Analytical Study of a Fractional Differential Model for COVID-19

Hatıra Günerhan ¹

¹ Department of Mathematics, Faculty of Education, Kafkas University, Kars, Turkey

gunerhanhatira@gmail.com,

Abstract

This study investigates a nonlinear SITR system that models the dynamics of COVID-19 by incorporating fractional-order derivatives in the sense of the Caputo definition. The model categorizes the population into four compartments: susceptible (S1), infected (I), under treatment (T), and recovered (R). To analyze the fractional-order SITR system, the Laplace Adomian Decomposition Method (LADM) is employed. Approximate analytical solutions are obtained, demonstrating rapid convergence and effectively capturing the system's behavior. The accuracy and applicability of the method are assessed, confirming the validity of the proposed fractional-order model.

Keywords: Caputo fractional derivative; Laplace Adomian decomposition; Zika virus; Nonlinear system.

REFERENCES

1. Umar M, Sabir Z, Raja MAZ, Shoaib M, Gupta M, Sánchez YG. A Stochastic Intelligent Computing with Neuro-Evolution Heuristics for Nonlinear SITR System of Novel COVID-19 Dynamics. *Symmetry*. 2020; 12(10):1628.
2. Ahmed, Nauman, Amr Elsonbaty, Ali Raza, Muhammad Rafiq, and Waleed Adel. Numerical simulation and stability analysis of a novel reaction–diffusion COVID-19 model. *Nonlinear Dynamics* 106 (2021): 1293-1310.
3. Adel, W., Günerhan, H., Nisar, K.S. et al. Designing a novel fractional order mathematical model for COVID-19 incorporating lockdown measures. *Sci Rep* 14, 2926, 2024.
4. Khairulbahri, Muhamad. The SEIR model incorporating asymptomatic cases, behavioral measures, and lockdowns: Lesson learned from the COVID-19 flow in Sweden. *Biomedical Signal Processing and Control* 81 (2023): 104416.

Analytical Solution of a Mathematical Model for the Spread of the Zika Virus

Hatıra Günerhan ¹

¹ Department of Mathematics, Faculty of Education, Kafkas University, Kars, Turkey

gunerhanhatira@gmail.com

Abstract

In this study, a nonlinear SEIR system is analyzed to model the spread of the Zika virus, using a modified form of the Caputo fractional derivative. Analytical and approximate-analytical solutions of the proposed model are derived using the Laplace Adomian Decomposition Method (LADM). The solutions are presented as rapidly converging series. Approximate analytical solutions are obtained, demonstrating rapid convergence and accurately capturing the system's dynamics. The reliability of the method is verified, and the validity of the proposed fractional-order model is substantiated.

Keywords: Caputo fractional derivative; Laplace Adomian decomposition; Zika virus; Nonlinear system.

REFERENCES

1. Suantai S, Sabir Z, Raja MAZ, Cholanjiak W. Numerical Computation of SEIR Model for the Zika Virus Spreading. *Comput Mater Contin.* 2023;75(1):2155–2170.
2. S. Rezapour, H. Mohammadi and A. Jajarmi, A new mathematical model for Zika virus transmission, *Advances in Difference Equations*, vol. 2020, no. 1, pp.1–15, 2020.
3. Adel, W., Günerhan, H., Nisar, K.S. et al. Designing a novel fractional order mathematical model for COVID-19 incorporating lockdown measures. *Sci Rep* 14, 2926, 2024.
4. D.Musso, T. Nhan, E. Robin, C. Roche, D. Bierlaire et al., “Potential for Zika virus transmission through blood transfusion demonstrated during an outbreak in French Polynesia,” *Eurosurveillance*, vol. 19, no. 14, pp. 20761, 2014.

Applications of bifurcation analysis, sensitivity analysis and modified Kudryashov method

Arzu AKBULUT¹

¹Department of Mathematics, Bursa Uludağ University, Bursa, Turkey

arzuakbulut@firat.edu.tr,

Abstract

In this paper, we obtain the exact solution to a nonlinear differential equation that plays a crucial role in analyzing signal transmission and exploring nonlinear effects, making it indispensable for advancing various technological domains. Its significance spans optical communications, nonlinear phenomena, and other related fields, driving innovation and enhancing our ability to harness light for diverse applications. Then, we handled the bifurcations of the model and were interested in the sensitivity analysis.

Keywords: Modified Kudryashov method; Bifurcations; Sensitivity analysis.

REFERENCES

1. N. Gao, Y. Wang, Traveling wave solutions for the n-dimensional generalized Zakharov–Kuznetsov equation with variable coefficients, Optical and Quantum Electronics, Vol:53, No:58, 2021.
2. M. Gu, J. Li, F. Liu, Z. Li, C. Peng, Propagation of traveling wave solution of the strain wave equation in microcrystalline materials, Vol:22, No:1, 20240093, 2024.
3. M.E., Elbrolosy, A.A. Elmandouh, Dynamical behaviour of nondissipative double dispersive microstrain wave in the microstructured solids. European Physical Journal Plus, Vol:136, No:955, 2021.
4. A. Akbulut, A.H. Arnous, M.S. Hashemi, M. Mirzazadeh, Solitary waves for the generalized nonlinear wave equation in (3+1) dimensions with gas bubbles using the Nucci's reduction, enhanced and modified Kudryashov algorithms. Journal of Ocean Engineering and Science, 2022.(in press) <https://doi.org/10.1016/j.joes.2022.07.002>.

An artificial neural network procedure for the nonlinear typhoid fever disease system

Ayşe Nur Akkilic¹, Hasan Bulut¹, Zulqurnain Sabir²,

¹Department of Mathematics, Firat University, Elazığ, Turkey

²Department of Computer Science and Mathematics, Lebanese American University, Beirut, Lebanon

Email: akkilicaysenur@gmail.com

Abstract

The current investigations provide the numerical solutions of the nonlinear typhoid fever disease system by using an artificial neural network procedure. The model is categorized into four dynamics susceptible, exposed, infected, and recovered. The use of one class into another category makes the model nonlinear and numerical solutions of the model are achieved through the stochastic computing procedure, while the optimization is performed by using the Levenberg-Marquardt backpropagation (LMQBP) neural network.

Keywords: Typhoid fever disease; Neural network; Transfer function; Levenberg-Marquardt; numerical outputs.

REFERENCES

1. Mushayabasa, S., Bhunu, C.P. and Ngarakana-Gwasira, E.T., 2013. Mathematical analysis of a typhoid model with carriers, direct and indirect disease transmission. *International Journal of Mathematical Sciences and Engineering Applications*, 7(1), pp.79-90.
2. Ivanoff, B., Levine, M.M. and Lambert, P., 1994. Vaccination against typhoid fever: present status. *Bulletin of the World Health Organization*, 72(6), p.957.
3. Nsutebu, E.F., Martins, P. and Adigo, D., 2003. Prevalence of typhoid fever in febrile patients with symptoms clinically compatible with typhoid fever in Cameroon. *Tropical medicine & international health*, 8(6), pp.575-578.
4. Schemmer, A.K., 2012. Heterogeneity of inflammation and host metabolism in a typhoid fever model (Doctoral dissertation, University_of_Basel).

Prediction of Concrete Compressive Strength Using Mamdani-Type Fuzzy Inference System

Osman Tufan¹, Okan Dardağan¹, Hakan Altunay¹

¹Department of Industrial Engineering, Süleyman Demirel University, Isparta, Turkey

osmantufan.sdu@gmail.com, okandardagan@gmail.com, hakanaltunay@sdu.edu.tr

Abstract

As technology progresses, artificial intelligence (AI) techniques, which offer the potential for faster and more reliable results, have superseded conventional methodologies. Nowadays, AI techniques are commonly utilized for the prediction of concrete compressive strength. Predicting concrete compressive strength as accurately as possible by leveraging various parameters is of vital importance for numerous aspects, including primarily the construction of safe structures, as well as cost and time savings, and the support of sustainable building practices. Through AI-based prediction models, insights into concrete compressive strength can be obtained at very early stages, enabling the early detection of potential issues in material quality or mix design. As with many real-world problems, modelling and controlling concrete compressive strength to achieve the most accurate representation with experimental results is a very challenging and complex task. In this study, fuzzy logic approach is utilized to predict the 28-day concrete compressive strength. In the fuzzy logic model designed in Python, input parameters are determined as CaO, MgO, NaEq, LOI (Loss on Ignition), Blaine, 1-day, 2-day and 7-day compressive strengths. Trapezoidal and triangular membership functions are used to model the input and output parameters. Additionally, the Mamdani inference mechanism is used to evaluate the generated rule base. The centroid method is employed for defuzzification. The results demonstrate that the fuzzy logic approach can be effectively used to predict concrete compressive strength.

Keywords: Compressive Strength; Forecasting; Time Series; Piecewise Linear Fuzzy Number; Fuzzy Logic.

REFERENCES

1. Khademi, F., Akbari, M., Jamal, S. M., Nikoo, M., Multiple Linear Regression, Artificial Neural Network, and Fuzzy Logic Prediction of 28 Days Compressive Strength of Concrete, *Frontiers of Structural and Civil Engineering*, Vol:11, 90-99, 2017.
2. Cheng, M. Y., Chou, J. S., Roy, A. F., Wu, Y. W., High-Performance Concrete Compressive Strength Prediction using Time-Weighted Evolutionary Fuzzy Support Vector Machines Inference Model, *Automation in Construction*, Vol:28, 106-115, 2017.

Topp-Leone Power Garima Distribution: Properties and Applications

Ayşe Metin Karakaş¹ Ashhan Demir² Sinan Çalık³

¹Department of Statistics, University of Bitlis Eren, Elazig, Turkey

^{2,3}Department of Statistics, University of Firat, Elazig, Turkey

aysekarakas5767@gmail.com¹, a.bilici128@gmail.com², scalik@firat.edu.tr³

Abstract

This article introduces the Topp-Leone Power Garima Distribution (TLPGD), a novel continuous probability distribution derived by integrating the Topp-Leone generator with the Power Garima distribution. The proposed model enhances flexibility in modeling skewed and complex data across various domains. In this study, we explore the mathematical properties of the TLPGD distribution, including its probability density function (PDF), cumulative distribution function (CDF), moments, and entropy measures. Additionally, we discuss parameter estimation using maximum likelihood estimation (MLE) and demonstrate its applications through real-world datasets. The results indicate that the TLPGD distribution provides a superior fit compared to traditional models, making it a valuable tool for statisticians and data analysis. The TLPGD's probability density function (PDF) and cumulative distribution function (CDF) are derived, showcasing its ability to generalize the Power Garima distribution via the Topp-Leone transformation. Analytical expressions for moments, quantiles, entropy, and order statistics are provided. The TLPGD exhibits right-skewed, unimodal, and bathtub-shaped hazard rates, making it suitable for diverse real-world scenarios. The TLPGD offers a robust tool for modeling lifetime data in engineering, medicine, and environmental sciences, with potential extensions to bivariate and regression frameworks suggested for future research.

Keywords: Topp-Leone distribution, Power Garima distribution, hazard function, maximum likelihood estimation, reliability analysis

REFERENCES

1. Shanker, R. (2016). Garima distribution and its application to model behavioral science data. *Biometrics & Biostatistics International Journal*, 4(7), 1-9.
2. Topp, C.W.; Leone, F.C. A family of J-shaped frequency functions. *J. Am. Stat. Assoc.* 1955, 50, 209–219. [CrossRef]
3. Abebe, B., Tesfay, M., Eyob, T., & Shanker, R. (2019). A two-parameter power Rama distribution with properties and applications. *Biometrics and Biostatistics International Journal*, 8(1), 6-11.
4. Al-Shomrani, A.; Arif, O.; Hanif, S.; Shahbaz, M.Q.; Shawky, A. Topp–Leone Family of Distributions: Some Properties and Application. *Pak. J. Stat. Oper. Res.* 2016, 12, 443. [CrossRef]

Travelling-wave Solutions of Evolution Equations

Begüm Sağlam, Elif Nisa Sert, Zeliha Geçimli, Mazlum Gezer and Halis Yilmaz

Department of Mathematics, University of Dicle, Diyarbakır, Turkey

begumsaglam03@gmail.com, elifnisasert@gmail.com, zelihagecimli51@gmail.com,
mazlumgezer21@gmail.com, halisyilmaz@dicle.edu.tr

Abstract: Integrable systems and evolution equations are closely linked, especially in mathematical physics. Evolution equations describe how systems change over time, often appearing as partial differential equations (PDEs) or ordinary differential equations (ODEs). When an equation is integrable, it means that it can be solved exactly.

Many famous integrable systems are governed by evolution equations, such as the Korteweg–de Vries (KdV) equation and the nonlinear Schrödinger (NLS) equation. These equations often produce solitons—stable wave-like structures that maintain their shape over time due to the integrability of the system. Methods such as the inverse scattering technique, Hirota method, and Darboux transformations are commonly used to solve these equations and reveal their underlying structure.

Motivated by this, we present explicit traveling-wave solutions for the general KdV and NLS equations.

Keywords: Integrable systems; travelling-wave solutions; solitons.

REFERENCES

1. Ablowitz, M.J. and Clarkson, P.A., *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Vol.149, London Mathematical Society Lecture Note Series), Cambridge University Press (1991).
2. Drazin, P.G. and Johnson, R.S., *Solitons: An Introduction*, Cambridge University Press (1989).
3. Korteweg, D.J. and de Vries, G., *On the Change of Form of Long Waves Advancing in a Rectangular Canal, and on a New Type of Long Stationary Waves*, The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science, 39(240), 422–443 (1895).
4. Russell, J.S., *Report on waves*, Fourteenth meeting of the British Association for the Advancement of Science, 1844.

SOME SPECTRAL PROBLEMS OF DIFFUSION OPERATOR WITH HIGH ORDER APPROXIMATIONS

Mine Babaoglu

Department of Mathematics and Science Education, University of Kahramanmaras Sutcu
Imam, Kahramanmaras, Turkey

mnbabaoglu@gmail.com

Abstract

In this work, we obtain some spectral results of Diffusion operator with high order approximations. Also, we focus on the theorems which include the solution functions belong to the Paley-Wiener Space.

Keywords: Diffusion operator; Sampling theory.

REFERENCES

1. M. G. Gasymov, G. Sh. Guseinov, Determination of diffusion operator on spectral data, Dokl. Akad. Nauk Azerb. SSSR 37, 2, 19-23, 1981.
2. H. Koyunbakan, E. S. Panakhov, Half inverse problem for diffusion operators on the finite interval, J. Math. Anal. Appl., 326, 1024-1030, 2007.
3. A. I. Zayed, Advances in Shannon's Sampling Theory, CRC Press, Boca Raton, 1993.
4. B. M. Levitan, I. S. Sargsjan, Introduction to Spectral Theory: Selfadjoint Ordinary Differential Operators, American Mathematical Society, Providence, Rhode Island, 1975.
5. B. Chanane, Computing eigenvalues of regular Sturm-Liouville problems, Appl. Math. Lett., 12, 119-125, 1999.
6. B. Chanane, High Order Approximations of the Eigenvalues of Regular Sturm-Liouville Problems, J. Math. Anal. Appl., 226, 121-129, 1998.

New Lindley Distribution and Applications

Ayşe METİN KARAKAŞ¹, Fatma BULUT² and Murat KARAKAŞ³

¹ Department of Statistics, Bitlis Eren University, Bitlis, Turkey.

^{2,3} Department of Mathematics, Bitlis Eren University, Bitlis, Turkey.

akarakas@beu.edu.tr, fbulut@beu.edu.tr, mkarakas@beu.edu.tr

Abstract

In this paper, we investigated the new Lindley distribution a new, flexible statistical model for positively skewed and heavy-tailed data offering closed-form formulas, moment analysis, and strong real-world applicability across survival, reliability, and biomedical datasets.

Keywords: Lindley family; survival analysis; reliability data; hazard function.

REFERENCES

1. Ramos, M. A., Cordeiro, G. M., Marinho, P. D., Dias, C. B., Hamadani, G. G., The Zografos-Balakrishnan log-logistic distribution: properties and applications, Journal of Statistical Theory and Applications, Vol. 12, No. 3, 225–244, 2013.
2. Yakubu, A., Luguterah, A., Nasiru, S., On the odd inverse exponential class of distributions: properties, applications and cure fraction regression, Journal of Statistical Management Systems, Vol. 25, No. 4, 805–836, 2022.
3. Alizadeh, M., MirMostafaei, S. M. T. K., Altun, E., Ozel, G., Ahmadi, M. Kh., The odd log-logistic Marshall-Olkin power Lindley distribution: properties and applications, Journal of Statistical Management Systems, Vol. 20, 1065–1093, 2018.
4. Elbatal, I., Elgarhy, M., Transmuted quasi Lindley distribution: a generalization of the quasi Lindley distribution, International Journal of Pure and Applied Science and Technology, Vol. 18, 59–70, 2013.
5. Elgarhy, M., Elbatal, I., Diab, L. S., Hwas, H. K., Shawki, A. W., Transmuted generalized quasi Lindley distribution, International Journal of Science and Engineering Science, Vol. 1, 1–8, 2017.

COMPARISON OF SOLUTION METHODS IN NON-LINEAR EQUATIONS WITH ONE UNKNOWN

Cemil İNAN¹

¹Department of Mathematics, University of Mardin Artuklu, Mardin, Turkey

cemilinan@artuklut.edu.tr,

Abstract

In cases where solutions cannot be made with analytical methods, numerical methods are used to find the approximate roots of the equation. In this study, from the nonlinear equation solution methods with one unknown; closed interval and Open methods. These methods (strengths and weaknesses) were examined and compared.

Keywords: Numerical methods; Closed; Open approximate solutions.

REFERENCES

1. An Introduction to Numerical Analysis for Electrical and Computer Engineers, Christopher J. Zarowski, A JOHN
2. Elementary Numerical Analysis: An Algorithmic Approach, McGraw-Hill, 1980.
3. • An Introduction to Numerical Analysis, Endre S`uli and David F. Mayers, Cambridge University Press, 2003.

INVESTIGATION OF ITERATIVE APPROXIMATE SOLUTIONS OF LINEAR EQUATION SYSTEMS

Cemil İNAN¹

¹Department of Mathematics, University of Mardin Artuklu, Mardin, Turkey

cemilinan@artuklu.edu.tr,

Abstract

Solutions of problems encountered in engineering and applied science fields may not always be found with direct solution methods. The lack of analytical methods to solve the encountered problem has led to the use of numerical methods. Iterative methods are generally preferred for large linear equation systems. Among these are the “Jacobi iteration method”, the Gauss-Seidel iteration method” and the “(Successive Over Relaxation – SOR) method”. In this study, these methods are examined and their strengths are emphasized.

Keywords: Numerical Method, Iterative Approach, Error in Approximate

REFERENCES

1. Karacan,S.(2006)
.https://cdn.bartın.edu.tr/matematik/9dcf61c4011949c35216a4e035dc3c88/ders-notu.pdf
2. Bayıroğlu,H.2024
https://eelektrik.tr/wp-content/uploads/2024/11/EELEKTRİK-SAYISAL-COZUMLEME-01-EE.pdf

Regime-Based Modeling of USD/TL via Stochastic Differential Equations and Change Point Estimation

Sevda Ozdemir Calikusu¹, Fevzi Erdogan²

¹Department of Accounting and Tax, Van Yuzuncu Yil University, Van, Turkey

sevdaoazdemir@yyu.edu.tr

²Department of Econometrics, Van Yuzuncu Yil University, Van, Turkey

ferdogan@yyu.edu.tr

Abstract

This study examines the USD/TRY exchange rate over the period from April 30, 2015, to April 30, 2025, through a regime-based stochastic differential equation (SDE) modeling framework. Using change point estimation, we identify two structural breaks on November 10, 2021, and March 23, 2022, partitioning the data into three distinct regimes. For each regime, we apply tailored SDEs reflecting their unique statistical characteristics. In Regime 1 (pre-November 2021), a Cox-Ingersoll-Ross (CIR) process is adopted due to its suitability for modeling mean-reverting and strictly positive series with heteroskedasticity. Regime 2 spans a short and turbulent period with limited data, requiring the use of 4-hour interval GBM simulations to enhance sample density for valid calibration. This enriched data enables the application of the Heston stochastic volatility model. In Regime 3 (post-March 2022), characterized by moderately declining volatility, the Heston model continues to provide a flexible structure. The CIR model and Heston models are calibrated using Maximum Likelihood Estimation (MLE), which enables direct parameter inference under each regime's stochastic structure. By integrating regime-dependent dynamics and simulation-assisted calibration in short regimes, the approach achieves higher fidelity in capturing the evolving behavior of the exchange rate. Overall, the methodology emphasizes the significance of data-driven regime segmentation and stochastic modeling for financial time series in emerging markets.

Keywords: change point, Cox-Ingersoll-Rose model, exchange rate modeling, Heston model, stochastic differential equations, stochastic volatility, regime switching, simulation.

REFERENCES

1. Cox, J. C., Ingersoll, J. E., Ross, S. A., A Theory of the Term Structure of Interest Rates, *Econometrica*, 53(2), 385–407, 1985.
2. Heston, S. L., A Closed-Form Solution for Options with Stochastic Volatility with Applications to Bond and Currency Options. *Review of Financial Studies*, 6(2), 327–343, 1993.
3. Bai, J., Perron, P., Computation and Analysis of Multiple Structural Change Models. *Journal of Applied Econometrics*, 18(1), 1–22, 2003.

BLOW UP OF SOLUTIONS FOR A FRACTIONAL PETROVSKY EQUATION

Erkan Sancar¹ and Erhan Pişkin²

¹Institute of Natural and Applied Sciences, Dicle University, Diyarbakır, Turkey

²Dicle University, Department of Mathematics, Diyarbakır, Turkey

erkan_sancar@hotmail.com, episkin@dicle.edu.tr

Abstract

This presentation focuses on the fourth-order wave equation incorporating a fractional time delay. We provide preliminary details about the equation to establish the blow-up of its solutions.

Keywords: Blow up, Fractional, Fourth-order equation.

REFERENCES

1. R. Aounallah, A. Benaissa, A. Zraï, Blow-up and asymptotic behavior for a wave equation with a time delay condition of fractional type. *Rendiconti del Circolo Matematico di Palermo Series 2*, 70 (2021) 1061-1081.
2. V. Georgiev, G. Todorova, Existence of solutions of the wave equation with nonlinear damping and source terms. *J. Differ. Equ.* 109 (1994) 295-308.
3. E. Pişkin, T. Uysal, Blow up of the solutions for the Petrovsky equation with fractional damping terms, *Malaya Journal of Matematik*, 6 (2018) 85-90.
4. E. Pişkin, *Evolüsyon Denklemlerin Çözümlerinin Patlaması (Blow up of Solutions of Evolution Equations)*, Pegem Akademi Yayıncılık, 2022.

MATHEMATICAL FOUNDATIONS OF TECHNICAL INDICATORS USED IN FINANCIAL TIME SERIES

Tümer Akbayır

Department of Mathematics, University of Firat, Elazığ

tumerakbayir@gmail.com

Abstract

In this study, the mathematical background and models behind technical analysis indicators, which are widely used in crypto and stock exchange markets, will be analysed and their links with financial time series data, which are generally stochastic in nature, noisy and high volatility sequences, will be investigated. In addition, in this study, the effects of indicators on the decision-making process of buying or selling in money markets will be analysed.

REFERENCES

1. Timothy J. Mcintosh, Fundamental Analysis (2012) in The Sector Strategist. John Wiley & Sons, Ltd, 163-184. doi:10.1002/9781119205333.ch9.
2. Ed Ponsi, Technical Analysis and Chart Interpretations: A Comprehensive Guide to Understanding Established Trading Tactics for Ultimate Profit, 2016, Wiley Trading Series.

THE CURVES WITH MODIFIED ORTHOGONAL FRAME OF IN THE MINKOWSKI 3-SPACE

İremnur Yüzer Boynikoğlu¹ and Handan Öztekin¹

¹Department of Mathematics, University of Firat, Elazig, Turkiye

irem_nur_yuzer@hotmail.com and hbalgetir@firat.edu.tr

Abstract

In this study, the modified frames with both the non-zero curvature and the torsion of the non-unit speed timelike and spacelike curves in Minkowski 3-space E_1^3 are investigated. The relationships between the derivative vectors of the modified frames and the Frenet vectors or the vectors of the modified frames of the timelike and spacelike curves are given.

Keywords:: Minkowski Space; Modified Frame.

REFERENCES

1. Arıkan, M., Nurkan, S.K. (2020). Adjoint Curve According to Modified Orthogonal Frame with Torsion in 3-Space. Usak University Journal of Science and Natural Sciences. 2, 54 – 64.
2. Azak, A.Z. (2021). Involute-Evolute Curves According to Modified Orthogonal Frame. Journal of Science and Arts. 21(2), 385 – 394.
3. Bükcü, B., Karacan, M.K. (2016). On the modified orthogonal frame with curvature and torsion in 3-space. Mathematical Sciences and " Applications E-Notes. 4(1), 184 – 188.
4. Elsayied, H.K., Altaha, A.A., Elsharkawy, A. (2021). On Some Special Curves According to the Modified Orthogonal Frame in Minkowski 3-space E_1^3 . Kasmera, 49, 2-15.
5. Gür Mazlum, S. and Bektaş, M. (2022). On the modified orthogonal frames of the non-unit speed curves in Euclidean 3-Space E^3 , Turkish Journal of Science., 7(2), 58-74.
6. Lone, M.S., Es, H., Karacan, M.K., Bükcü, B. (2019). On some curves with modified orthogonal frame in Euclidean 3-Space. Iranian Journal of " Science and Technology, Transactions A: Science. 43(4), 1905 – 1916.
7. Uddin, S., Stankovic, M.S., Iqbal, M., Yadav, S.K., Aslam, M. (2022). Slant helices in Minkowski 3-space E_1^3 with Sasai's modified frame fields. Filomat 36, 151 – 164.

A Look at The Analysis of Coefficients of Univalent Functions with Artificial Intelligence

Hakan Faik Er (*), Ekrem Kadioğlu (**)

(*) Atatürk University, Institute of Science, Master's Student, Erzurum.

(**) Atatürk University, Faculty of Science, Department of Mathematics, Erzurum.

Abstract

With artificial intelligence, the information on some subjects can be accessed very easily. We will investigate the effectiveness of the artificial intelligence when the subjects become complex. For this purpose, we examined the competence of artificial intelligence on the subject of estimating the coefficients of the univalent functions.

Keywords: Univalent Functions; The coefficients of Univalent Functions; the artificial intelligence

References:

1. Duren P.L., 1983. Duren univalent functions. Springer-Verlag, New York.
2. Graham I. and Kohr G., 2003. Geometric function theory in one and higher dimensions.
3. Silverman H., 1975. Univalent functions with negative coefficients. Proceedings of the American Mathematical Society 51, no (1), 109-116.
4. Zorn P., 1986. The Bieberbach conjecture. Mathematics Magazine 59, no (3), 131-148.

DETERMINING THE SHORTEST ROUTE FOR AIR TRANSPORT AFTER A POSSIBLE FIRE IN MUĞLA PROVINCE USING ARTIFICIAL BEE COLONY ALGORITHM

Yasemin Demirel^{1,*}, Oya Mert Coşkun²

^{1*} Department of Management Information System, Fenerbahçe University, İstanbul, Türkiye, yasemin.bakir@fbu.edu.tr, ORCID: 0000-0001-8051-6012

² Department of Mathematics, Tekirdağ Namık Kemal University, Tekirdağ, Türkiye, oyamert@nku.edu.tr, ORCID: 0000-0002-8791-3341

Abstract

According to the official information on the OGM page, 360,556.20 hectares of land were burned and 54,201 fires broke out in Türkiye between 2000-2023. Of these fires, 4,622 were intentional, 25,183 were accident-negligence, 6,692 were natural, and the cause of 18,835 was unknown (OGM, 2023). In line with this data, it was aimed to determine the possible routes to be used during a fire in Muğla, the province with the highest probability of a fire. Reaching and responding to a fire in a short time reduces the amount of damage that may occur after the fire. In this study, it is considered to use the ABC Algorithm, which is an artificial bee colony algorithm, in the event of a possible Muğla fire. In the proposed study, the ABC algorithm was applied to reach the fire in the shortest time by air and the shortest route was determined.

Keywords: Artificial bee colony algorithm, Shortest Path, Optimization

REFERENCES

1. Karaboğa, D. 2004. Yapay Zeka Optimizasyon Algoritmaları, Atlas Yayın Dağıtım.
2. Orman Genel Müdürlüğü (OGM), (2023). Orman yangınları istatistikleri, <https://www.ogm.gov.tr/tr/e-kutuphane/resmi-istatistikler>
3. Yu, Yi & Wu, Yonggang & Liu, Xinglong. (2020). An enhanced artificial bee colony algorithm for numerical function optimization. IOP Conference Series: Materials Science and Engineering. 806. 012024. 10.1088/1757-899X/806/1/012024.
4. Devadason, J.R., Hepsiba, P.S. & Solomon, D.G. (2024). Case studies on the applications of the artificial bee colony algorithm. *Sādhanā* **49**, 152. <https://doi.org/10.1007/s12046-024-02498-9>

Optical solutions for Fokas system in optical fiber

Sibel Şehriban ATAŞ¹

¹ Department of Finance and Banking, University of Munzur, Tunceli, Turkey

sibelatas@munzur.edu.tr,

Abstract

The Modified Generalized Riccati Equation Mapping Method (M-G-REMM) is used in this work to examine the Fokas system, which models the distribution of unpredictable pulse in optical fibers. The equation that defines the system is transformed into a nonlinear ordinary differential equation (N-ODE) using a traveling wave transformation. Consequently, novel complex hyperbolic travelling wave solutions are obtained in multitude. As far as we are aware, no prior literature has reported on these solutions. Regarding the Fokas System, all of our findings have been confirmed using Mathematica and Maple. To enhance comprehension of the dynamic behavior of the current optical solutions with different time parameter values, graphical simulations of many graphs are provided. This makes (M-G-REMM) useful and practical for comprehending nonlinear issues in optic fiber, plasma physics, applied sciences.

Keywords: Modified Generalized Riccati Equation Mapping Method; Fokas system; Optical soliton solutions.

REFERENCES

- [1] Tarla, S., Ali, K. K., Sun, T. C., Yilmazer, R., & Osman, M. S., Nonlinear pulse propagation for novel optical solitons modeled by Fokas system in monomode optical fibers. *Results in Physics*, (2022), 36, 105381.
- [2] Naher, H., & Abdullah, F. A. (2012). New Traveling Wave Solutions by the Extended Generalized Riccati Equation Mapping Method of the (2+ 1)-Dimensional Evolution Equation. *Journal of Applied Mathematics*, 2012, (1), 486458.
- [3] Khater, M. M. (2021). Analytical simulations of the Fokas system; extension (2+ 1)-dimensional nonlinear Schrödinger equation. *International Journal of Modern Physics B*, 35(28), 2150286.
- [4] Baskonus, H. M., & Bulut, H. (2016). Exponential prototype structures for (2+ 1)-dimensional Boiti-Leon-Pempinelli systems in mathematical physics. *Waves in Random and Complex Media*, 26(2), 189-196.

Hybrid Jump-Heston and Bidirectional LSTM Model with Attraction Mechanism for S&P 500 Forecasting

Davood Ahmadian¹, Vahid Norouzi², Fevzi Erdogan³, Sevda Ozdemir Calikusu⁴

¹ Faculty of Mathematics, Statistics and Computer Science, University of Tabriz, Tabriz, Iran

d.ahmadian@tabrizu.ac.ir

² Faculty of Economics and Management, University of Tabriz, Tabriz, Iran

v.norouzi@tabrizu.ac.ir

³ Department of Econometrics, Faculty of Economics and Administrative Sciences, Van Yuzuncu Yil University, Van, Turkey

ferdogan@yyu.edu.tr

⁴ Department of Accounting and Tax, Ozalp Vocational School, Van Yuzuncu Yil University, Van, Turkey

sevdaoazdemir@yyu.edu.tr

Abstract

This study introduces a novel hybrid model for S&P 500 forecasting, integrating the Jump-diffusion Heston stochastic volatility model with an attention-enhanced Bidirectional LSTM (Bi-LSTM). The model leverages Jump-Heston for capturing stochastic volatility and market jumps alongside the Bi-LSTM with attention for temporal pattern learning and feature focus. Trained on daily S&P 500 data (2000-2024), preliminary results demonstrate superior forecasting performance over traditional models by effectively capturing market dynamics like volatility clustering and jump risks. This highlights its potential for risk management and algorithmic trading (Ge, 2025; Ouyang et al., 2024).

Keywords: LSTM, Heston, Jump-Diffusion, S&P500.

REFERENCES

1. Ouyang, Z., Lu, M., Ouyang, Z., Zhou, X., & Wang, R. (2024). A novel integrated method for improving the forecasting accuracy of crude oil: ESMD-CFastICA-BiLSTM-Attention. *Energy Economics*, 138, 107851.
2. Ge, Q. (2025). Enhancing stock market Forecasting: A hybrid model for accurate prediction of S&P 500 and CSI 300 future prices. *Expert Systems with Applications*, 260, 125380.

On a solvable difference equations system of second order its solutions are related to a generalized Mersenne sequence

Murad khan Hassani

¹Department of Mathematics, University of Nevsehir Haci Baktas Veli, Turkey

muradkhanhassani@gmail.com

Abstract

In this paper, we consider a class of two-dimensional nonlinear difference equations system of second order, which is a considerably extension of some recent results in the literature. Our main results show that class of system of difference equations is solvable in closed form theoretically. It is noteworthy that the solutions of aforementioned system are associated with generalized Mersenne numbers. The asymptotic behaviour of solution to aforementioned system of difference equations when $a=b$ and $p=0$ are also given. Finally, numerical examples are given to support the theoretical results presented in this paper.

Keywords: System of difference equations, solvable system, periodicity.

REFERENCES

Abo-Zeid R., On the solutions of a higher order difference equation, Georgian Mathematical Journal 2020; 27(2): 165-175.

THE APPROXIMATION OF FIXED POINT OF ENDRICHED ϕ -CONTRACTION MAPPINGS

Ümmügülsüm Günay¹ and Murat Özdemir²

^{1,2}Department of Mathematics, University of Ataturk, Erzurum, Turkey

ummugulsum.gunay18@ogr.atauni.edu.tr; mozdemir@atauni.edu.tr

Abstract

We introduce the class of enriched ϕ -contractions in Banach spaces as a natural generalization of ϕ -contractions and study the existence and approximation of the fixed points of mappings in this new class, which is shown to be an unsaturated class of mappings in the setting of a Banach space. We illustrated the usefulness of our fixed point results by studying the existence and uniqueness of the solutions of some second order (p,q) -difference equations with integral boundary value conditions.

Keywords: Banach space; enriched ϕ -contraction; enriched cyclic ϕ -contraction; fixed point; Maia type fixed point theorem; (p,q) -difference equation; integral boundary value condition

REFERENCES

1. Banach, S. Sur les opérations dans les ensembles abstraits et leurs applications aux équations intégrales. *Fundam. Math.* 1922, 3, 133–181. [CrossRef]
2. Agarwal, R.P.; O'Regan, D.; Sahu, D.R. *Fixed Point Theory for Lipschitzian-Type Mappings with Application; Topological Fixed Point Theory and Its Applications*; Springer: New York, NY, USA, 2009.
3. Agarwal, R.P.; Karapinar, E.; O'Regan, D.; Roldán-López-de-Hierro, A.F. *Fixed Point Theory in Metric Type Spaces*; Springer: Cham, Switzerland, 2015.
4. Berinde, V. *Iterative Approximation of Fixed Points*, 2nd ed.; *Lecture Notes in Mathematics*, 1912; Springer: Berlin, Germany, 2007.
5. Granas, A.; Dugundji, J. *Fixed Point Theory*; Springer: New York, NY, USA, 2003.
6. Kirk, W.A.; Sims, B. (Eds.) *Handbook of Metric Fixed Point Theory*; Kluwer Academic Publishers: Dordrecht, The Netherlands, 2001.
7. Kirk, W.; Shahzad, N. *Fixed Point Theory in Distance Spaces*; Springer: Cham, Switzerland, 2014.
8. Rus, I.A.; Petrusel, A.; Petrusel, G. *Fixed Point Theory*; Cluj University Press: Cluj-Napoca, Romania, 2008.
9. Caccioppoli, R. Un teorema generale sull'esistenza di elementi uniti in una trasformazione funzionale. *Rend. Accad. Lincei.* 1930, 11, 794–799.
10. Berinde, V.; Pacurar, M. Iterative approximation of fixed points of single-valued almost contractions. In *Fixed Point Theory and Graph Theory*; Elsevier/Academic Press: Amsterdam, The Netherlands, 2016; pp. 29–97.
11. Rus, I.A. *Generalized Contractions and Applications*; Cluj University Press: Cluj-Napoca, Romania, 2001.
12. Rus, I.A. Picard Operators and Applications. *Sci. Math. Jpn.* 2003, 58, 191–219.
13. Rus, I.A.; Serban, M.-A. Basic problems of the metric fixed point theory and the relevance of a metric fixed point theorem. *Carpathian J. Math.* 2013, 29, 239–258. [CrossRef]
14. Browder, F.E. On the convergence of successive approximations for nonlinear functional equations. *Indag. Math.* 1968, 30, 27–35. [CrossRef]

ON ASYMPTOTICALLY EQUIVALENT SEQUENCES IN PROBABILITY

Kübra Elif Akbaş¹, Mahmut Işık²

¹ Faculty of Medicine, Firat University, Elazığ, Türkiye

² Faculty of Education, Harran University, Şanlıurfa, Türkiye

misik63@yahoo.com

Abstract

In this study, we give some results related with asymptotically equivalent sequences in probability.

Keywords: Sequence of random variables; Asymptotical statistical equivalence.

REFERENCES

- [1]Akbaş, KE and Işık, M. Weighted Statistical Convergence in Probability, 8th International Conference on Computational Mathematics and Engineering Sciences 17--19 May 2024, Şanlıurfa/ Türkiye.
- [2]Akbaş, KE and Işık, M. Weighted Statistical Convergence of Order α in Probability, Proceedings of the Institute of Mathematics and Mechanics (To appear).
- [3]Braha, NL.; Srivastava, HM. and Et, M. Some weighted statistical convergence and associated Korovkin and Voronovskaya type theorems, J. Appl. Math. Comput. 65(1-2), 2021.

GENERALIZED BIVARIATE FIBONACCI AND LUCAS POLYNOMIALS BY HESSENBERG MATRICES

Yasemin Taşyurdu¹ and Beyza Aydan²

¹Department of Mathematics, Erzincan Binali Yıldırım University, Erzincan, Türkiye

²Department of Mathematics, Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, Türkiye

ytasyurdu@erzincan.edu.tr, beyzaaydan92@gmail.com

Abstract

In this paper, we give some Hessenberg matrices associated with generalized bivariate Fibonacci and Lucas polynomials, and then investigate the relationships between these matrices and polynomials. We show that determinants and permanents of these Hessenberg matrices are generalized bivariate Fibonacci and Lucas polynomials, which generalize the known results for Fibonacci, Lucas, Pell, Jacobsthal, Fermat, Chebyshev, Morgan-Voyce and Vieta polynomials in both univariate and bivariate forms.

Keywords: Bivariate Fibonacci Polynomial; Bivariate Lucas Polynomial; Determinant; Hessenberg Matrix; Permanent.

REFERENCES

1. Cahill, N. D., D'Errico, J. R., Narayan, D. A. Narayan, J. Y. 2002. Fibonacci Determinants, *College Mathematics Journal*, 33(3), 221-225.
2. Hoggatt Jr., V. E., Bicknell, M. 1973. Generalized Fibonacci Polynomials and Zeckendorf's Theorem, *The Fibonacci Quarterly*, 11(4), 399-419.
3. Kocer, E. G., Tuncel, S. 2016. Bivariate Fibonacci and Lucas Like Polynomials, *Gazi University Journal of Science*, 29(1), 109-113.
4. Ocal, A. A., Tuglu, N., Altinisik, E. 2005. On the Representation of k -Generalized Fibonacci and Lucas Numbers, *Applied Mathematics and Computation*, 170(1), 584-596.
5. Taşyurdu Y., Gültekin, İ. 2014. Determinantal and Permanent Representation of q -Fibonacci Polynomials, *Qscience Connect*, 25, 5p.
6. Taşyurdu, Y. 2018. On Generalized Fibonacci-Like Sequences by Hessenberg Matrices, *International Journal of Mathematics Trends and Technology*, 64(1), 51-58.

BI-PERIODIC GENERALIZED BIVARIATE FIBONACCI POLYNOMIALS

Yasemin Taşyurdu¹ and Beyza Aydan²

¹Department of Mathematics, Erzincan Binali Yıldırım University, Erzincan, Türkiye

²Department of Mathematics, Graduate School of Natural and Applied Sciences, Erzincan Binali Yıldırım University, Erzincan, Türkiye

ytasyurdu@erzincan.edu.tr, beyzaaydan92@gmail.com

Abstract

In this paper, we define bi-periodic generalized bivariate Fibonacci polynomials, which generalize Fibonacci, Pell, Jacobsthal polynomials and bi-periodic generalized Fibonacci polynomials that include bi-periodic Fibonacci, Pell and Jacobsthal polynomials, and other well-known generalized polynomials, in both univariate and bivariate forms. We obtain Binet formulas and generating functions of these polynomials, and then investigate their convergence properties. Also, we prove some new identities that are consistent with the known results of generalized bivariate Fibonacci polynomials.

Keywords: Bi-periodic Fibonacci polynomials; Bivariate Fibonacci polynomial, Generalized Fibonacci polynomial.

REFERENCES

1. Edson, M., Yayenie, O, 2009. A New Generalization of Fibonacci Sequence and Extended Binet's Formula. *Integers*. 9, 639-654.
2. Hoggatt Jr., V. E., Bicknell, M. 1973. Generalized Fibonacci Polynomials and Zeckendorf's Theorem, *The Fibonacci Quarterly*, 11(4), 399-419.
3. Kocer, E. G., Tuncel, S. 2016. Bivariate Fibonacci and Lucas Like Polynomials, *Gazi University Journal of Science*, 29(1), 109-113
4. Taşyurdu, Y. 2022. Bi-Periodic Generalized Fibonacci Polynomials, *Turkish Journal of Science*, 7(3), 157-167.
5. Yılmaz, N., Coşkun, A., Taşkara, 2017. N. On Properties of Bi-Periodic Fibonacci and Lucas polynomials. *AIP Conference Proceedings*, 1863, 310002.

Epidemiological Modeling with Seasonal Environmental Effects

İrem Akbulut Arık*, Seda İĞRET ARAZ*

*Department of Mathematics Education, Siirt University, Siirt, Turkey

iremakbulut@siirt.edu.tr, sedaaraz@siirt.edu.tr

Abstract

This study deal with a fractional-order epidemiological model incorporating fractional derivatives to capture memory effects inherent in disease transmission. To reflect environmental seasonality, almost periodic functions are introduced, allowing key parameters to vary over time. The existence and uniqueness of solutions are rigorously established. Numerical simulations are presented to illustrate the influence of fractional dynamics and seasonal fluctuations on disease behavior. The results highlight the significant impact of these factors on disease progression, providing a more realistic framework for understanding and managing epidemic outbreaks.

Keywords: Epidemiological modeling, fractional calculus, existence and uniqueness.

REFERENCES

1. Caputo M., Linear model of dissipation whose Q is almost frequency independent. II. Geophysical Journal International. 13 (5): 1967, 529-539.
2. İğret Araz S., Boulaaras S., Fractional modeling of gradual incorporation of infected prey into the predator-prey system with consideration of seasonality, hal-04929853.
3. Adewal, S.O., Adeniran, G.A., Olopade, I.A., Ajao, S.O., Mohammed, I.T.: Mathematical and sensitivity analysis of the dynamical spread of cholera. Int. J. Innov. Appl. Stud. 19(1), 46--54 (2017).
4. Huang, J.; Ruan, S.; Wu, X.; Zhou, X., Seasonal transmission dynamics of measles in China. Theory Biosci. 2018, 137, 185--195.
5. Rebelo, C.; Margheri, A.; Bacaër, N., Persistence in seasonally forced epidemiological models. J. Math. Biol. 2012, 64, 933--949.
6. Atangana A., İğret Araz S., Fractional Stochastic Differential Equations Applications to Covid-19 Modeling, Springer, 2022.

INVESTIGATING CROSSOVER BEHAVIORS IN CHAOTIC SYSTEMS

Seda İĞRET ARAZ¹

¹Department of Mathematics Education, Siirt University, Siirt, Turkey

sedaaraz@siirt.edu.tr,

Abstract

This work examines the concept of hybrid chaos, characterized by the integration of deterministic order and chaotic behavior within a unified framework. Employing a piecewise modeling approach, distinct chaotic systems are combined over defined intervals, with stochastic elements incorporated to enhance dynamical richness and adaptability. The proposed methodology offers a foundation for advancing hybrid chaotic system design, with potential applications in complex systems across various scientific domains.

Keywords: Crossover behaviors, hybrid chaotic attractors, stochastic approach.

REFERENCES

1. A. Akgul, I. Moroz, I. Pehlivan, S. Vaidyanathan, A new four-scroll chaotic attractor and its engineering applications, *Optik*, 127, (2016), 5491--5499.
2. Lorenz, Edward Norton, Deterministic nonperiodic flow, *Journal of the Atmospheric Sciences*. 20 (2), 1963, 130--141.
3. A. Arneodo, P. H. Coullet and E. A. Spiegel, The dynamics of triple convection, *Geophys. Astrophys. Fluid Dynamics*, 31, pp. 1-48, 1985.
4. Bouali S., A 3D Strange Attractor with a Distinctive Silhouette. The Butteráy Effect Revisited, 24th ABCM International Congress of Mechanical Engineering, (2013).
5. Atangana A., Igret Araz S., *Fractional Stochastic Differential Equations Applications to Covid-19 Modeling*, Springer, 2022.
6. Atangana A., Igret Araz S., *Piecewise differential equations: Theory, methods and applications*, *AIMS Mathematics* 8 (7), 2023.

EXAMINING THE GOLDEN RATIO-AESTHETIC RELATIONSHIP IN HISTORICAL BRIDGES IN DIYARBAKIR

Bilgesu ÇİFÇİ¹, Samet GÜNDÜZ²

¹ Bismil BİLSEM, Ministry of Education, Diyarbakır, Türkiye

² Bismil BİLSEM, Ministry of Education, Diyarbakır, Türkiye

bilgesuc8@gmail.com, asametgunduz@gmail.com

Abstract

The golden ratio is a special numerical proportion that is considered aesthetically pleasing and harmonious in both mathematics and art. The golden ratio, approximately equal to 1.618, appears widely in nature and everyday life, ranging from art to sculpture and graphic design. Throughout history, it has often been used in architecture to ensure aesthetic integrity and visual harmony. This research aims to examine the arches of historical bridges based on the concept of the golden ratio and evaluate these structures aesthetically. This study is significant in revealing the aesthetic balance in the design of historical bridges through the use of the golden ratio, thereby contributing to the preservation and evaluation of cultural heritage. The study was carried out on three historical bridges located in Diyarbakır, a city rich in historical and cultural heritage, hosting numerous civilizations throughout history. The presence of the golden ratio in the arches of the examined bridges was analyzed using span-to-height ratios. Deviations from the golden ratio were calculated, and these values were evaluated in terms of the golden ratio-aesthetic relationship. The results showed that the design principles used in bridge construction could be related to the golden ratio and aesthetic criteria. Thus, the importance of examining the aesthetic values of historical structures from a mathematical perspective was highlighted. Additionally, this study makes a significant contribution to the field by combining the aesthetic aspect of historical structures with mathematical analysis.

Keywords: Golden ratio; Aesthetic evaluation; Historic bridges.

REFERENCES

1. Akın, R. B. (2021). *Ottoman era mosque facades in Diyarbakir occupancyvacancy rates examination of the relationship between the golden ratio and aesthetics* (Publication No. 664813). [Master's thesis, Dicle University]. Turkish Council of Higher Education Thesis Center.
2. Yılmaz, E. M. (2017). *The relationship between golden ratio and aesthetics in madrasas of Seljuk period: The case of Konya*. (Publication No. 453524). [Master's thesis, Selçuk University]. Turkish Council of Higher Education Thesis Center.

**9TH INTERNATIONAL
CONFERENCE ON
COMPUTATIONAL
MATHEMATICS AND
ENGINEERING SCIENCES**

17 – 19 May 2025,
Diyarbakır – Türkiye

cmescongress.org

